

# A Versatile Growth Model with Statistically Stable Parameters

JON SCHNUTE

Department of Fisheries and Oceans, Resource Services Branch, Pacific Biological Station, Nanaimo, B.C. V9R 5K6

SCHNUTE, J. 1981. A versatile growth model with statistically stable parameters. Can. J. Fish. Aquat. Sci. 38: 1128–1140.

This paper presents a new comprehensive growth model which includes numerous historical models as special cases. The new model is derived from a concise biological principle which, unlike earlier theories, relates to growth *acceleration*. Properties of growth curves, such as asymptotic limits or inflection points, are shown to be incidental in this new context. Possible submodels include not only asymptotic growth (such as von Bertalanffy, Richards, Gompertz, or logistic growth) but also linear, quadratic, or exponential growth. By simple analysis of variance, the observed data can be used directly in deciding which type of model is most appropriate. The new model is cast in terms of parameters which have stable statistical estimates. From this perspective, it is shown how earlier formulations sometimes result in an endless computer search for optimal parameter estimates.

**Key words:** Growth, asymptotic growth, von Bertalanffy, Richards, Gompertz, logistic, length at age, weight at age, nonlinear parameter estimation

SCHNUTE, J. 1981. A versatile growth model with statistically stable parameters. Can. J. Fish. Aquat. Sci. 38: 1128–1140.

On trouvera dans l'article qui suit un modèle de croissance général nouveau incluant plusieurs anciens modèles comme cas spéciaux. Le nouveau modèle a été dérivé d'un principe biologique concis qui, contrairement aux théories antérieures, se rapporte à l'*accélération* de la croissance. Nous démontrons que les propriétés des courbes de croissance telles que limites asymptotiques ou points d'inflexion sont fortuites dans ce contexte nouveau. Des sous-modèles possibles incluent non seulement la croissance asymptotique (croissance de von Bertalanffy, Richards, Gompertz, ou logistique), mais également une croissance linéaire, quadratique ou exponentielle. On peut utiliser les données directement, par simple analyse de variance, afin de décider quel type de modèle est le plus approprié. Le nouveau modèle est présenté en termes de paramètres statistiquement stables. De ce point de vue, nous démontrons comment les formulations antérieures nécessitent parfois une recherche interminable par ordinateur en vue de trouver les estimations de paramètres optimales.

Received February 27, 1981

Accepted May 25, 1981

Reçu le 27 février 1981

Accepté le 25 mai 1981

THE biologist who wishes to find a curve describing size-at-age data invariably faces the problem of picking an appropriate model. Judging from the available literature, there is an extensive menu of models to contemplate, associated with such names or titles as Pütter, von Bertalanffy, Richards, Gompertz, or logistic. In addition to these models which all exhibit limiting, or asymptotic, growth, there are still others which do not, such as linear or exponential growth models. Ricker (1979) reviews much of the historical literature.

Even when a model is chosen, the biologist's problems are not over. There may also be a diverse suite of methods for estimating the parameters, such as a Walford (1946) plot or the procedures by Fabens (1965) or Allen (1966) to estimate certain von Bertalanffy parameters. Some methods fail to give all the parameters, and sometimes computer methods designed to estimate all parameters fail mysteriously. For example, it may happen that a parameter estimate slips gradually

towards infinity as the computer grinds on in endless pursuit.

If the parameters are actually found, at last, there still remains the problem of their interpretation and appropriateness. Consider, for example, generalized von Bertalanffy growth

$$(1) \quad Y(t) = y_{\infty}(1 - e^{-g(t-t_0)})^p,$$

Richards growth

$$(2) \quad Y(t) = y_{\infty}(1 + \frac{1}{p} e^{-g(t-t_0)})^{-p},$$

Gompertz growth

$$(3) \quad Y(t) = y_{\infty}e^{-e^{-g(t-t_0)}},$$

logistic growth (Richards growth with  $p = 1$ )

$$(4) \quad Y(t) = y_{\infty}(1 + e^{-g(t-t_0)})^{-1},$$

and linear growth

$$(5) \quad Y(t) = g(t - t_0),$$

where  $Y(t)$  represents fish size at age  $t$ , and  $y_\infty$ ,  $g$ ,  $p$ , and  $t_0$  are parameters, usually with  $y_\infty > 0$ ,  $g > 0$ , and  $p > 0$ . In (1) and (5) the parameter  $t_0$  represents a time when the growth curve crosses the  $t$ -axis. By contrast, in (2)–(4)  $t_0$  corresponds to an inflection point on the curve. Indeed, the curves (2) and (4) never cross the  $t$ -axis and neither does (2) if  $p > 0$ . Why should or shouldn't axis crossings and inflection points be important? Furthermore, if  $g > 0$ , then  $y_\infty$  in (1)–(4) is a theoretical limiting size. By contrast, in (5) there is no limiting size. Why should a limiting size be important or well defined? In addition, models (1) and (2) have four parameters, while (3) and (4) have three parameters, and (5) has only two. What is a reasonable number of parameters to use? The practitioner invariably faces such questions.

This paper presents a new comprehensive model which includes (1)–(5) as special cases and many other models as well. The model is motivated by a concise biological principle, and the four (or fewer) parameters in the model almost always have stable statistical estimates. In addition, the parameters have reasonable biological interpretations, and submodels correspond simply to limiting parameter values. These parametric properties allow direct use of the data in selecting an appropriate model systematically. The apparent problem of comparing parameters in models (1)–(5) disappears. Axis crossings, inflection points, and asymptotic size no longer play an essential or dominant role, although they can be identified easily if they occur. Finally, an example is given which illustrates how a data set can suggest Gompertz growth (3) as a limiting version of von Bertalanffy growth (1).

Throughout this paper, the model is formulated in the language of fish growth. Obviously, it need not be restricted to fish; it could apply to a life stanza for any organism. In addition, it might be used to describe the growth of a whole population. Such analogies are hardly new. For example, Schaefer (1954) used logistic growth as a model for tuna populations. Later, Pella and Tomlinson (1969) generalized this approach by employing, essentially, Richards growth. Still other applications are possible for the model here. It might, for instance, describe the total mortality of a cohort as a function of time or the mortality resulting from the sudden presence of toxic material. Each new application may require a shift in language, but the mathematics remains the same.

### Model Assumptions

As in equations (1)–(5), let  $Y(t)$  denote the size of a fish at age  $t$ . Here "size" can refer to standard length, hypural length, fork length, live weight, eviscerated weight, volume (measured by water displacement), or whatever standard measure of fish dimension the biologist seeks to describe. The rate of growth is then the derivative  $dY/dt$ , and

$$(6) \quad Z = \frac{1}{Y} \frac{dY}{dt}$$

represents the relative growth rate, that is, the rate of growth in size. Equivalently,  $Z$  represents the logarithmic

growth rate:

$$Z = \frac{d}{dt} (\log Y).$$

Just as one might study how the size  $Y(t)$  of a fish develops in relation to age, so one might also examine the development of the relative growth rate  $Z(t)$ . In particular, one might consider the seemingly abstract question: What is the relative growth rate of the relative growth rate, that is,

$$\frac{1}{Z} \frac{dZ}{dt} = ?$$

Perhaps the simplest possible assumption is that this quantity is a linear function of the relative growth rate:

$$(7) \quad \frac{1}{Z} \frac{dZ}{dt} = -(a + bZ),$$

where  $a$  and  $b$  are constants. For convenience, (7) is written with a minus sign on the right-hand side because the growth rate typically decreases. However,  $a$  and  $b$  should be considered unrestricted; either parameter can be positive, negative, or zero.

Conceptually, (7) represents a simple but significant departure from previous descriptions of the nature of growth. Traditionally, writers have focused on the growth rate  $dY/dt$ . [See Ricker's (1979) review or Fletcher (1973).] By contrast, (7) focuses on a rate of a rate, that is, *acceleration* in growth  $d^2Y/dt^2$ . A short calculation (given in Appendix B) shows that (6) and (7) together imply

$$(8) \quad \frac{d^2Y}{dt^2} = \frac{dY}{dt} [-a + (1 - b)Z].$$

Thus a fish whose growth conforms to (7) experiences growth acceleration which is proportional to the growth rate and to a linear function of the relative growth rate. Depending on the constants  $a$  and  $b$ , the animal's growth may be accelerated, decelerated, or both. In particular, the model includes possible S-shaped growth curves with a juvenile period of accelerated development followed later by decelerated growth during which the animal approaches a final limiting size.

For such S-curves, the parameters  $a$  and  $b$  have a rather simple significance. Associated with the S, there are two heights: that of the bottom arc and that of the full S. These correspond to the size when growth acceleration ends and to the final limiting size. There is also a width associated with the bottom arc of the S. This is the time spent during the accelerated growth phase. The dimensionless parameter  $b$  relates to the ratio of the two heights, or sizes, just mentioned. The parameter  $a$  has units  $\text{time}^{-1}$ . Its reciprocal  $1/a$  relates to the width, or time, described above. The precise mathematical relationships among  $a$ ,  $b$ , and S-curve characteristics are given later.

### Model Equations

I do not wish to propose (7) as a new esoteric principle which somehow underlies the true subtleties of biological

growth. Certainly the equation affords grounds for lengthy philosophical speculation, but the history of this problem suggests that such speculations have little practical value. The main purpose of (7) here is to provide a simple foundation for constructing a unified approach to a menu of growth models including (1)–(5) cited earlier. To achieve this goal, it is necessary first to incorporate (7) into a formal mathematical problem. Begin by rewriting (6) and (7), respectively, as follows:

$$(9) \quad \frac{dY}{dt} = YZ,$$

$$(10) \quad \frac{dZ}{dt} = -Z(a + bZ).$$

Equations (9) and (10) together comprise a pair of differential equations for the two functions  $Y(t)$  and  $Z(t)$ . They can be solved if suitable end conditions are specified. For example, it is enough to require that the size take two specific values at two particular times, that is,

$$(11) \quad Y(\tau_1) = y_1,$$

$$(12) \quad Y(\tau_2) = y_2.$$

It is also enough to specify both  $Y$  and  $Z$  at one particular time,

$$(13) \quad Y(\tau_1) = y_1,$$

$$(14) \quad Z(\tau_1) = z_1.$$

The formal development of the previous paragraph can be stated informally in the language of biology. Let  $\tau_1$  and  $\tau_2$  be two particular ages in the life of a fish. For example,  $\tau_1$  might be the age of a young fish and  $\tau_2$  the age of an old fish. Then, given the growth law (7), the fish's entire growth curve  $Y(t)$  is determined from its sizes  $y_1$  and  $y_2$  at ages  $\tau_1$  and  $\tau_2$ . The resulting curve depends on the four parameters in (9)–(12), namely  $a$ ,  $b$ ,  $y_1$ , and  $y_2$ . The two times  $\tau_1$  and  $\tau_2$  are regarded as chosen and fixed by the biologist. Similarly, given the growth law (7), the fish's entire growth curve  $Y(t)$  is determined by its size  $y_1$  and relative growth rate  $z_1$  at a particular time  $\tau_1$ . In this case the curve  $Y(t)$  depends on the four parameters in (9)–(10) and (13)–(14), namely  $a$ ,  $b$ ,  $y_1$ , and  $z_1$ .

Mathematically, (9)–(12) is a *two-point boundary value problem*. For convenience, call it Problem (I). Also, (9)–(10) and (13)–(14) is an *initial value problem*. Call it Problem (II). The solutions to both Problems (I) and (II) for both  $Y$  and  $Z$  are obtained in Appendix A. It turns out that the solution to (II) involves formulas which look a bit more compact than the solution to (I). However, for reasons which will be made clear in the statistical analysis later, the main emphasis in this paper is on the solution  $Y(t)$  to Problem (I). The form of the solution depends on whether or not the parameters  $a$  and  $b$  are zero. Explicitly, the solution in the four possible cases turns out to be:

Case 1:  $a \neq 0$ ,  $b \neq 0$

$$(15) \quad Y(t) = \left[ y_1^b + (y_2^b - y_1^b) \frac{1 - e^{-a(t-\tau_1)}}{1 - e^{-a(\tau_2-\tau_1)}} \right]^{1/b},$$

Case 2:  $a \neq 0$ ,  $b = 0$

$$(16) \quad Y(t) = y_1 \exp \left[ \log(y_2/y_1) \frac{1 - e^{-a(t-\tau_1)}}{1 - e^{-a(\tau_2-\tau_1)}} \right],$$

Case 3:  $a = 0$ ,  $b \neq 0$

$$(17) \quad Y(t) = \left[ y_1^b + (y_2^b - y_1^b) \frac{t - \tau_1}{\tau_2 - \tau_1} \right]^{1/b},$$

Case 4:  $a = 0$ ,  $b = 0$

$$(18) \quad Y(t) = y_1 \exp \left[ \log(y_2/y_1) \frac{t - \tau_1}{\tau_2 - \tau_1} \right].$$

It is assumed throughout this paper that  $y_2 > y_1 > 0$  and  $\tau_2 > \tau_1$ . Thus, the model describes growth, as opposed to shrinkage, in fish size.

There are four basic facts to realize about equations (15)–(18). First, they result from solving Problem (I), that is, equations (9)–(12). In fact, it is reasonably easy to see at once that (11) and (12) are satisfied. Simply substitute  $t = \tau_1$  or  $t = \tau_2$  in the right-hand side of any equation (15)–(18) and note that the expression collapses to  $y_1$  or  $y_2$ . Second, even if the formulas (15)–(18) look a bit complicated and unfamiliar, they determine a growth curve  $Y(t)$  from the four parameters  $a$ ,  $b$ ,  $y_1$ , and  $y_2$ . (Remember that  $\tau_1$  and  $\tau_2$  are ages fixed by the biologist for the species in question; typically  $\tau_1$  and  $\tau_2$  are young and old ages for a fish.) Third, although (15)–(18) look like four distinct formulas for  $Y(t)$ , in a numerical sense they are all one formula. If, for example,  $b$  is allowed to approach 0 in (15), then the limiting value of  $Y(t)$  is given by (16). Indeed, (15) is undefined when  $b = 0$ . Fourth, the formulas (15)–(18) incorporate all the models (1)–(5) and many others as well. Table 1 lists a number of special cases.

To see how the formulas (15)–(18) do indeed translate to models like (1)–(5), it is necessary to do a little algebraic tinkering in each case. For example, it turns out that equations (1) and (15) both have the general form

$$Y(t) = (\alpha + \beta e^{\gamma t})^\delta$$

for constants  $\alpha$ ,  $\beta$ ,  $\gamma$ ,  $\delta$ . Consequently, (1) and (15) represent the same curve with different sets of four parameters. In particular, (1) uses  $y_\infty$ ,  $g$ ,  $t_0$ , and  $p$ , while (15) uses  $a$ ,  $b$ ,  $y_1$ , and  $y_2$ . Each of these parameter sets can easily be transformed to the other. For example, in this case  $g = a$  and  $p = 1/b$ . Expressions for  $y_\infty$  and  $t_0$  are given in the next section. Incidentally, the von Bertalanffy growth law (1) with  $p = 1$  was rewritten parametrically in a form very similar to (15) with  $b = 1$  by Schnute and Fournier (1980). They used this form to obtain statistically stable estimates for mean length at age from length–frequency data.

In spite of the similarities, there is a very important difference between (15)–(18) and a list of models like (1)–(5). As stated above, (15)–(18) really constitute a single equation

TABLE 1. Special cases of the general model (15)–(18). For a historical review of model origins, see Ricker (1979). It is assumed in (15)–(18) that  $\tau_2 > \tau_1$  and  $y_2 > y_1 > 0$ .

Choice of $a, b$	Model title(s)	Equation
$a > 0, b > 0$	Generalized von Bertalanffy	(1)
$a > 0, b = 1$	Pütter No. 1	(1) with $p = 1$
$a > 0, b = \frac{1}{2}$	Specialized von Bertalanffy	(1) with $p = 3$
$a > 0, b = 0$	Pütter No. 2	(3)
$a > 0, b = 0$	Gompertz	(3)
$a > 0, b < 0$	Richards	(2)
$a > 0, b = -1$	Logistic	(4)
$a = 0, b = 1$	Linear	(5)
$a = 0, b = \frac{1}{2}$	Quadratic	(19)
$a = 0, b = 0$	$t$ th power	(20)
$a < 0, b = 1$	Exponential	(21)

expressed in several limiting forms. By contrast, (1)–(5) are separate equations, and it is impossible on the surface to see how they relate continuously to each other. For example, (1) always crosses the  $t$ -axis at  $t = t_0$ , while (3) never crosses that axis. It simply isn't clear how to manipulate the four parameters in (1) to obtain (3) as a limiting three-parameter model. This difficulty has practical consequences, illustrated numerically later.

Another new feature in the system (15)–(18) is that it includes more models than those considered relevant historically. Authors have primarily focused on models in which growth is asymptotic, that is, models which imply a limiting animal size. Yet the attitude toward such models is, at best, hesitant. Ricker (1979) includes a section titled "Asymptotic growth: is it real?" Knight (1968) writes about "Asymptotic growth: an example of nonsense disguised as mathematics." Equations (15)–(18) allow this whole debate to be resolved simply, because they include non-asymptotic models. This gives the actual size-at-age data the freedom to select the most appropriate model, asymptotic or not. The parameter estimates summarize the data's choice. For example, when  $a = 0$  and  $b = \frac{1}{2}$ , the model takes the form

$$(19) \quad Y(t) = (\alpha + \beta t)^2$$

with  $\beta > 0$ , that is, quadratic growth. If  $a = 0$  and  $b = 0$ , the model becomes essentially

$$(20) \quad Y(t) = \alpha \beta^t$$

with  $\alpha > 0$  and  $\beta > 0$ , that is,  $t$ -th power growth. As a final example, if  $a < 0$  and  $b = 1$ , the model reduces to the exponential growth equation

$$(21) \quad Y(t) = \alpha + \beta e^{\gamma t}$$

with  $\beta > 0$  and  $\gamma > 0$ .

### Model Parameters

The model (15)–(18) may or may not define a curve which crosses the  $y$ -axis, has an inflection point, or exhibits asymptotic behavior. If a particular curve has these properties, then

parameters can be associated with them. Table 2 lists notation for such parameters, along with others pertinent to the model. As formulated here, the model is cast in terms of two specified parameters,  $\tau_1$  and  $\tau_2$ , and four free parameters,  $a, b, y_1$ , and  $y_2$ . The remaining parameters in Table 2, if they exist, can be expressed in terms of these quantities. (See Appendix B for the required mathematical derivations.) For example, the relative growth rates  $z_1$  and  $z_2$  at times  $\tau_1$  and  $\tau_2$ , turn out to be

$$(22) \quad z_1 = \frac{de^{-a\tau_1}}{cy_1^b},$$

$$(23) \quad z_2 = \frac{de^{-a\tau_2}}{cy_2^b}$$

where

$$c = \begin{cases} (e^{-a\tau_2} - e^{-a\tau_1})/a, & a \neq 0, \\ \tau_2 - \tau_1, & a = 0, \end{cases}$$

$$d = \begin{cases} (y_2^b - y_1^b)/b, & b \neq 0, \\ \log(y_2/y_1), & b = 0. \end{cases}$$

The parameters  $c$  and  $d$  are introduced in (22) and (23) to simplify the treatment of the four cases resulting from whether or not  $a$  or  $b$  or both are 0.

If  $b = 0$ , then it can be shown (Appendix B) that there is no age  $\tau_0$  corresponding to a projected size 0. Similarly, if  $a = 0$ , then the growth curve is not asymptotic and has no inflection point. When they exist, the parameters appropriate to size 0, asymptotic size, and inflected growth are (Appendix B):

$$(24) \quad \tau_0 = \begin{cases} \tau_1 + \tau_2 - \frac{1}{a} \log \left[ \frac{e^{a\tau_2} y_2^b - e^{a\tau_1} y_1^b}{y_2^b - y_1^b} \right]; & a \neq 0, b \neq 0; \\ \tau_1 + \tau_2 - \frac{\tau_2 y_2^b - \tau_1 y_1^b}{y_2^b - y_1^b}; & a = 0, b \neq 0; \end{cases}$$

$$(25) \quad y_\infty = \begin{cases} \left[ \frac{e^{a\tau_2} y_2^b - e^{a\tau_1} y_1^b}{e^{a\tau_2} - e^{a\tau_1}} \right]^{1/b}; & a \neq 0, b \neq 0; \\ \exp \left( \frac{e^{a\tau_2} \log y_2 - e^{a\tau_1} \log y_1}{e^{a\tau_2} - e^{a\tau_1}} \right); & a \neq 0, b = 0; \end{cases}$$

$$(26) \quad \tau^* = \begin{cases} \tau_1 + \tau_2 - \frac{1}{a} \log \left[ \frac{b(e^{a\tau_2} y_2^b - e^{a\tau_1} y_1^b)}{y_2^b - y_1^b} \right]; & a \neq 0, b \neq 0; \\ \tau_1 + \tau_2 - \frac{1}{a} \log \left[ \frac{e^{a\tau_2} - e^{a\tau_1}}{\log(y_2/y_1)} \right]; & a \neq 0, b = 0; \end{cases}$$

$$(27) \quad y^* = \begin{cases} \left[ \frac{(1-b)(e^{a\tau_2} y_2^b - e^{a\tau_1} y_1^b)}{e^{a\tau_2} - e^{a\tau_1}} \right]^{1/b}; & a \neq 0, b \neq 0; \\ \exp \left( \frac{(e^{a\tau_2} \log y_2 - e^{a\tau_1} \log y_1)}{e^{a\tau_2} - e^{a\tau_1}} - 1 \right); & a \neq 0, b = 0; \end{cases}$$

TABLE 2. A list of possible parameters associated with the model (15)–(18). The final five parameters may not always exist. When  $\tau_0$  and  $y_\infty$  exist, they may not have the significance shown here.

Parameter	Defining equation	Units	Description
$\tau_1$	—	Time	First specified age
$\tau_2$	—	Time	Second specified age
$a$	See (7)	Time <sup>-1</sup>	Constant relative rate of relative growth rate
$b$	See (7)	Dimensionless	Incremental relative rate of relative growth rate
$y_1$	$y_1=Y(\tau_1)$	Size	Size at age $\tau_1$
$y_2$	$y_2=Y(\tau_2)$	Size	Size at age $\tau_2$
$z_1$	$z_1=Z(\tau_1)$	Time <sup>-1</sup>	Relative growth rate at age $\tau_1$
$z_2$	$z_2=Z(\tau_2)$	Time <sup>-1</sup>	Relative growth rate at age $\tau_2$
$\tau_0$	$Y(\tau_0)=0$	Time	Age of theoretical zero size
$\tau^*$	$\frac{d^2Y}{dt^2}(\tau^*)=0$	Time	Age of growth inflection
$y^*$	$y^*=Y(\tau^*)$	Size	Size at age of inflection
$z^*$	$z^*=Z(\tau^*)$	Time <sup>-1</sup>	Relative growth rate at age of inflection
$y_\infty$	$\lim_{t \rightarrow \infty} Y(t)$	Size	Asymptotic size

(28)  $z^* = \frac{a}{1 - b}$ .

As in earlier equations, the expression pairs (24)–(27) each comprise one continuous numerical result; that is, the second formula simply gives the limiting form of the first when one parameter vanishes. Even if  $a \neq 0$  and  $b \neq 0$ , it may still happen that some of the parameters (24)–(27) fail to exist. For example, if the argument of the logarithm, i.e. the expression in brackets, is negative in (24), then  $\tau_0$  is undefined. It may also happen that  $\tau_0$  and  $y_\infty$  do not have their usual significance related to size 0 and asymptotic size. All such difficulties are put into complete perspective in the next section.

Equations (22)–(28), although perhaps slightly complex, achieve the main purpose of this section: to express the last seven parameters of Table 2 in terms of the first six. The final equation (28) stands out in this list because of its simple form. Notice that the relative growth rate  $z^*$  at an inflection point depends *only* on  $a$  and  $b$ , not on  $\tau_1$ ,  $\tau_2$ ,  $y_1$ , or  $y_2$ . In essence  $z^*$  is determined by the differential equation (7) alone, regardless of the boundary conditions. Two other interesting parameters also have this property, namely, (i) the time interval  $\tau^* - \tau_0$  between the projected age at size 0 and the age of growth inflection and (ii) the fraction  $y^*/y_\infty$  of asymptotic size achieved by the age of growth inflection. Explicitly, in terms of  $a$  and  $b$ , these parameters are (Appendix B):

(29)  $\tau^* - \tau_0 = \frac{1}{a} \log \left( \frac{1}{b} \right)$ ,

(30)  $y^*/y_\infty = (1 - b)^{1/b}$ .

Equations (29) and (30) give full mathematical expression to the earlier discussion of S-curve characteristics. Notice that  $b$  determines the ratio  $y^*/y_\infty$ , which is just the ratio of heights

of the lower S-arc and the full S. Similarly, once  $b$  is fixed, then  $\frac{1}{a}$  determines the width  $\tau^* - \tau_0$  of the lower S-arc.

As suggested in the previous section, the formulas (24)–(26) lead to a simple practical result. Suppose that, given two specified ages  $\tau_1$  and  $\tau_2$ , the four parameters  $a$ ,  $b$ ,  $y_1$ , and  $y_2$  are found appropriate to a growth model (15)–(18). Then, provided they exist, the traditional parameters  $\tau_0$ ,  $y_\infty$ , and  $\tau^*$  can be found at once from (24)–(26). The curve can then be expressed equivalently in either von Bertalanffy form

(31)  $Y(t) = y_\infty(1 - e^{-a(t-\tau_0)})^{1/b}$ ,

or Richards form

(32)  $Y(t) = y_\infty(1 - b e^{-a(t-\tau^*)})^{1/b}$ ,

where (31) and (32) are counterparts to (1) and (2), respectively, in the notation of Table 2. Consequently, in some cases, alternative sets of four parameters, such as  $(y_\infty, \tau_0, a, b)$  or  $(y_\infty, \tau^*, a, b)$ , can also be used to describe the curve. There is, nevertheless, a compelling reason to focus primarily on  $(y_1, y_2, a, b)$ . These parameters exist for any curve in the family, while other parameters, such as  $y_\infty$ ,  $\tau_0$ , and  $\tau^*$ , may not.

Historically, it is possible to see how first-order differential equations gave a limited view of available growth curves. Both (31) and (32) satisfy

(33)  $\frac{dY}{dt} = \frac{a}{b} \left[ \left( \frac{y_\infty}{Y} \right)^b - 1 \right] Y$ .

This first-order three-parameter differential equation (proved in Appendix B) makes it seem as if  $y_\infty$  needs to be a model parameter. By contrast, the second-order two-parameter dif-

Can. J. Fish. Aquat. Sci. Downloaded from www.nrcresearchpress.com by ARLIS - Alaska Resources Lib & Info Services on 09/17/18  
For personal use only.




FIG. 1. Set of eight regions in the  $a, b$ -plane defined by four lines. The solid lines correspond to the  $a$ -axis and  $b$ -axis. The broken line parallel to the  $a$ -axis is defined by  $b = 1$ . The diagonal broken line with negative slope corresponds to equation (34) in the text.

Differential equation (7) or (8) avoids this impression and leads one to consider the larger family of curves (15)–(18).

### Model Properties

The parameters  $\tau_0$ ,  $y_\infty$ ,  $\tau^*$ , and  $y^*$  are defined in (24)–(27) only when the expressions in square brackets are positive. As Appendix C shows, this seemingly troublesome problem works out, in fact, to be a major clue for understanding the various faces of the model (15)–(18). There are eight possible characteristic shapes for growth curves. These depend primarily on the two parameters  $a$  and  $b$ . Figure 1 shows the  $a, b$ -plane divided into eight regions by four lines: the  $a$ -axis, the  $b$ -axis, the line  $b = 1$ , and the sloping line

$$(4) \quad b = - \frac{\tau_2 - \tau_1}{\log(y_2/y_1)} a.$$

Notice that only the fourth line (34) involves the remaining parameters  $\tau_1$ ,  $\tau_2$ ,  $y_1$ , and  $y_2$ . It is guaranteed to have negative slope by the assumptions that  $\tau_2 > \tau_1$  and  $y_2 > y_1 > 0$ . Figure 2 displays a collection of eight typical growth curves. A simple rule determines which curve fits a particular parameter set  $(\tau_1, \tau_2, y_1, y_2, a, b)$ . First determine the number of the region in Fig. 1 where the pair  $(a, b)$  lies; then find the curve numbered in Fig. 2. Appendix C gives mathematical justification for this correspondence between Fig. 1 and Fig. 2. Notice that the curves in Fig. 2 show which parameters  $\tau_0$ ,  $y_\infty$ ,  $\tau^*$ ,  $y^*$  are defined for each case.

It may happen that the point  $(a, b)$  lies on a boundary between two regions in Fig. 1, or even on one of the three vertices where the boundary lines meet. In such cases, the appropriate characteristic curve in Fig. 2 may be unclear. This section concludes with a detailed description of all the various possibilities.

One general principle should be considered whenever the model (15)–(18) is applied: an empirical curve can represent size-at-age data only for those ages with known size. A curve may be quite unrealistic when applied to an entire fish life, yet may give a completely adequate description of one life phase. For example, models which predict eventual growth to infin-




FIG. 2. Set of eight characteristic growth curves appropriate to a parameter pair  $(a, b)$  in each of the eight regions shown in Fig. 1. The curves all represent plots of size  $Y$  on the vertical scale against age  $t$  on the horizontal scale.

ite size are not necessarily inappropriate over a limited period of rapid growth, such as the larval stage.

Case 1:  $0 < a, 0 < b < 1$ .

This case represents the classical situation in which all parameters of Table 2 are defined. The curve is S-shaped and asymptotic with limiting size  $y_\infty$ . It has an inflection point  $(\tau^*, y^*)$ , and it crosses the  $t$ -axis at age  $\tau_0$ .

Case 2:  $0 < a, 1 \leq b$ ,

In this case the curve is asymptotic and crosses the  $t$ -axis, but it has no inflection point. On the boundary where  $b = 1$ , the model corresponds to the specialized von Bertalanffy curve (1) with  $p = 1$ . This three-parameter curve (also called a Pütter No. 1 curve by Ricker (1979)) is typically used in the particular case when "size" refers to "length." There is really no a priori reason to limit the model in this way. Analysis of variance (described later) can be used to decide when it is appropriate to restrict one or more parameter values.

Case 3:  $-b \log(y_2/y_1)/(\tau_2 - \tau_1) < a \leq 0, 1 < b$ .

When  $(a, b)$  lies in this region, the growth curve is a not asymptotic, but becomes unbounded. It crosses the  $t$ -axis at

age  $\tau_0$ , and an initial period of decelerated growth begins at this age. The curve continues later with an indefinite period of accelerated growth. Such curves may not occur commonly, but it is easy to imagine circumstances in which they might be appropriate. If a young fish experiences a difficult struggle for food until it reaches a critical size  $y^*$  (when things get increasingly better), then its growth pattern might follow a curve of this type.

As the point  $(a,b)$  approaches the boundary where  $a = 0$ , the final period of accelerated growth moves to the right. When  $a = 0$ , the curve represents unbounded decelerated growth. The line  $a = 0$  between regions 2 and 3 is also a limiting possibility for case 2, where  $y_\infty$  becomes infinite as  $a$  approaches 0.

Case 4:  $-b \log(y_2/y_1)/(\tau_2 - \tau_1) < a \leq 0, 0 \leq b \leq 1$ .

Here the corresponding curve starts on the  $t$ -axis and continues upward with unbounded accelerated growth. Such a curve might represent fish growth restricted, say, to the larval stage. The particular case  $(a,b) = (0,1)$  corresponds to linear growth; that is, the curve becomes a straight line (no acceleration) with positive slope. This point  $(0,1)$  is on the boundary of regions 1, 2, 3, and 4. The reader can perhaps see from Fig. 2 how a straight line could be a limiting possibility for each of the four relevant curves.

Case 5:  $a \leq -b \log(y_2/y_1)/(\tau_2 - \tau_1), 0 \leq b$ .

As in case 4, the curve here also represents unbounded accelerated growth. The difference is that the curve in this case cannot be extrapolated back to a theoretical age  $\tau_0$  corresponding to size 0. Instead, there is a theoretical minimum size  $y_\infty$ . Notice in Fig. 2 the unconventional significance of the parameter  $y_\infty$  here. Curves of this type might describe fish growth during a rapid developmental stage which begins slowly at size  $y_\infty$ .

As the point  $(a,b)$  moves to the boundary of this region, the parameter  $y_\infty$  moves towards 0. Thus, on the region boundary, the  $t$ -axis is a lower asymptote to the curve. This property also applies to the special point  $(a,b) = (0,0)$ , which lies on the boundary of regions 1, 4, 5, 6, 7, and 8. At this vertex, the model reduces to  $t$ -th power growth, equation (18) or (20).

Case 6:  $a < 0, b < 0$ .

Here the curve is similar to the previous case, but with a rather dramatic extra feature. Growth is now so rapid that the fish theoretically reaches infinite size by the finite age  $\tau_0$ . In this context both the parameters  $y_\infty$  and  $\tau_0$  have unconventional significance. Curves of this type might apply to a life stage where increased size continually makes the fish more accessible to an increased food supply. While it lasts, such a positive feedback effect could lead to extremely accelerated growth.

Case 7:  $0 \leq a \leq -b \log(y_2/y_1)/(\tau_2 - \tau_1), b < 0$ .

This case is almost the same as the previous one, except

that the lower asymptote is now the  $t$ -axis. As the parameter pair  $(a,b)$  approaches the sloping line (34), that is, the boundary between regions 7 and 8, the parameter  $\tau_0$  becomes infinite. In this limiting situation, the curve simply represents unbounded accelerating growth.

Case 8:  $-b \log(y_2/y_1)/(\tau_2 - \tau_1) < a, b \leq 0$ .

Here the curve is S-shaped once again. Unlike the classical situation of case 1, however, the curve does not extrapolate back to the  $t$ -axis. Instead, it has the  $t$ -axis as a lower asymptote. Such a curve might be appropriate if early fish growth is extremely gradual.

## Model Errors and Nonlinear Estimation

So far the model (15)–(18) has been posed deterministically, as if the data never deviated from the growth curve. Obviously this is unrealistic, and statistical assumptions are required to make the model workable. To be explicit, suppose that  $n$  fish are each sampled to determine an observed size  $\hat{Y}$  and age  $t$ . This gives  $n$  data points  $(t_i, \hat{Y}_i)$ ,  $i = 1, 2, \dots, n$ . For each age  $t_i$ , there is a corresponding size  $Y_i$  predicted by the model. Two common error assumptions are

$$(35) \quad \hat{Y}_i = Y_i + \sigma_1 \epsilon_i; \quad i = 1, \dots, n;$$

$$(36) \quad \hat{Y}_i = Y_i e^{\sigma_2 \epsilon_i}; \quad i = 1, \dots, n;$$

where the random variables  $\epsilon_i$  ( $i = 1, \dots, n$ ) are assumed to be independent and normal with mean 0 and variance 1. In (35) the errors are additive, and the parameter  $\sigma_1$ , which has units of size, measures the additive standard error in prediction. The second model (36) includes multiplicative errors. The dimensionless parameter  $\sigma_2$  measures the logarithmic standard predictive error.

In some cases the data collection process may suggest more complex models which admit errors in both the size  $Y$  and age  $t$ . Such models could easily be formulated, but their analysis would be a digression from the main purpose of this paper. Regardless of the model, the practitioner is ultimately faced with a nonlinear parameter estimation problem. For example, parameter estimates for (35) are obtained by minimizing the sum of squares

$$(37) \quad S_1(y_1, y_2, a, b) = \sum_{i=1}^n \left[ \hat{Y}_i - Y_i(y_1, y_2, a, b) \right]^2.$$

In (37)  $\tau_1$  and  $\tau_2$  are presumed fixed, so that the predicted sizes  $Y_i$  and the sum  $S_1$  depend on the remaining parameters. Similarly, estimates for the model (36) require minimizing

$$(38) \quad S_2(y_1, y_2, a, b) = \sum_{i=1}^n \left[ \log(\hat{Y}_i / Y_i(y_1, y_2, a, b)) \right]^2.$$

At this point, the practitioner often feels alienated from the analysis. If the sums  $S_1$  and  $S_2$  could be minimized by linear regression, then the estimation problem would be considered finished. Since the parameters enter the model nonlinearly,

the need to minimize  $S_1$  or  $S_2$  sometimes seems to be a tremendous technical impediment. Not surprisingly, the historical literature on this problem includes numerous intricate and specialized methods for estimating parameters in related models, like (1), (2), (3), or (4).

In my experience, there are two common misconceptions about nonlinear estimation. (I might add that I began work on nonlinear problems a few years ago with these misconceptions alive and well in my own mind.) The first is the belief that technical methods for finding a minimum are necessarily complex and difficult to use. The second is that, if such a method were easily available, then it would solve the problem completely, just as linear regression does. Reality often lies in the face of both beliefs. Modern computing technology makes it easy to interface with good minimization methods. Yet, when programs are implemented, mysterious problems may occur and the search for a minimum can seem endless.

As an explicit example, suppose that the practitioner wishes to describe a data set with model (1), generalized von Bertalanffy growth. The problem, apparently, is to estimate  $y_\infty$ ,  $g$ ,  $t_0$ , and  $p$ . This can be done quite easily, for example, with the aid of a general function minimization program written by Schnute (1981). (See Appendix D for further details.) The practitioner need only supply a few lines of code to define model (1), and the computer does the rest. In spite of this simple implementation, the method may fail, as illustrated in the next section. The difficulty lies not with the search method, but with the model posed. Here is the crux of the problem, typical in nonlinear estimation. It's easy enough to set up a minimizing scheme; the real problem centers on formulating the model in terms of estimable parameters. That is why the bulk of this paper is spent placing the model on a proper parametric foundation. Once this is accomplished, statistical questions can be resolved rather easily.

Unfortunately, the concept of "proper parameters" isn't easy to define in general. There is an art to choosing the right parameters for a particular nonlinear model. The growth model in this paper illustrates three possible guidelines for making that choice. First, the parameters should, if possible, relate directly to the data. Here  $y_1$  and  $y_2$  have this property because they correspond to actual points on the growth curve. Second, the parameters should be influenced only slightly by minor changes in the model. Here, for instance,  $z_1$  and  $z_2$  might be poor choices of parameters. Minor curve changes would influence the slopes  $z_1$  and  $z_2$  more than the positions  $y_1$  and  $y_2$ . Such reasoning led to the earlier decision to formulate the model here from problem (I), rather than (II).

The two guidelines just cited both relate to local dependence of the model on its parameters. The third guideline deals with global parameter variations. As the parameters tend to the limits of definition, they should not define a potentially useful curve. For instance, model (1) can tend to a perfectly reasonable Gompertz curve as  $t_0$  and  $p$  become infinitely large negative and positive, respectively, as illustrated in the next section. The model (15)–(18) does not suffer a similar defect. It can be shown that, as the point  $(a, b)$  moves to infinity in any direction, the corresponding curve in Fig. 2 becomes one with a sudden jump at one of the ages  $\tau_1$ ,

$\tau_2$ , or  $\tau^*$ . Such curves are unrealistic; consequently, any useful curve must indeed correspond to a finite point  $(a, b)$ .

## Example

In connection with his discussion of von Bertalanffy growth, Ricker (1975, p. 226) lists length and weight data for ciscoes (*Coregonus artedii*) from Vermilion Lake, Minnesota. The mean weights in grams for ages 2–11 are given, respectively, as 99, 193, 298, 383, 462, 477, 505, 525, 539, and 539. These means are based on different numbers of fish at each age, so parameter estimates might perhaps be found by minimizing a weighted sum of squares  $S_1$  or  $S_2$ . For the sake of simple illustration, however, consider the problem of fitting model (1) with additive errors (35) to these 10 data points. This can be accomplished, as described earlier, with the aid of software by Schnute (1981). (See Appendix D.)

Table 3 summarizes the results of the fitting process, which begins with a wild guess at point No. 1 ( $y_\infty = 550$ ,  $g = 0.5$ ,  $t_0 = 0$ ,  $p = 1$ ). This rather poor choice gives a large initial sum of squares  $S_1$ . For a start, model (1) is restricted to the special case  $p = 1$ , the Pütter No. 1 curve (Table 1). After about 160 calculations of  $S_1$ , the algorithm easily locates point No. 2, where  $S_1$  is minimized subject to the requirement that  $p = 1$ .

Traditionally, the Pütter No. 1 curve ( $p = 1$ ) is used to describe lengths, while weights are presumed to follow a Pütter No. 2 ( $p = 3$ ). The logical basis for this choice is that weight is volumetric and should therefore relate to length cubed. Point No. 3 in Table 3 shows the relevant parameter estimates with this assumption. Notice that the sum of squared model errors  $S_1$  at point No. 3 (615.3) is only about 35% of its value at point No. 2 (1755.3). Consequently, the three-parameter Pütter No. 2 curve really does fit the cisco weight data better than the three-parameter Pütter No. 1 curve.

It remains only to free the fourth parameter  $p$  to see how much the fit can be improved with the general model (1). Point No. 4 in Table 4 represents a failed attempt to do so. Once the algorithm is allowed to tamper freely with  $p$ , it succeeds in reducing  $S_1$  gradually by increasing  $p$  and decreasing  $t_0$  endlessly. Here this search is arbitrarily stopped at point No. 4 in Table 4, after several hundred calculations of  $S_1$ . Apparently, an optimum model (1) cannot be located.

TABLE 3. The sum of squares  $S_1$  for cisco weight data (see text) and model (1) at four points. Point No. 1 is a starting point for the search algorithm. Points 2 and 3 correspond to minima for  $S_1$  when  $p = 1$  and  $p = 3$ , respectively. Point No. 4 is reached by stopping the algorithm arbitrarily during an endless search for a general minimum.

	1	2	3	4
	$S_1$			
	171660	1755.3	615.3	429.0
$y_\infty$	550	584.0	555.5	548.1
$g$	0.5	0.3031	0.4574	0.5304
$t_0$	0	1.488	0.2746	-4.363
$p$	1	1	3	50.38



TABLE 4. The sum of squares  $S_1$  for cisco weight data (see text) and model (15)–(16) at six points. The first four points are taken from Table 3. Points 5 and 6 correspond to minima for  $S_1$  in (15) and (16), respectively. Parameters absent from this Table are not defined.

	1	2	3	4	5	6
	$S_1$					
	171660	1755.3	615.3	429.0	398.0	423.1
$y_1$	347.7	84.02	90.32	94.83	97.81	95.05
$y_2$	547.8	551.3	543.3	540.2	538.2	539.9
$a$	0.5	0.3031	0.4574	0.5304	0.5801	0.5356
$b$	1	1	0.3333	0.01985	-0.1890	0
$\tau_0$	0	1.488	0.2746	-4.363	—	—
$\tau^*$	0	1.488	2.676	3.027	3.218	3.046
$y_{\infty}$	550	584.0	555.5	548.1	544.1	547.6
$y^*$	0	0	164.6	199.6	217.7	201.4
$z^*$	—	—	0.6861	0.5411	0.4879	0.5356
$z_1$	0.2910	1.804	1.142	0.9469	0.8502	0.9380
$z_2$	0.002052	0.01796	0.01023	0.007731	0.006339	0.007560
Region	2	2	1	1	8	8

The first four columns of Table 4 present the same four points as Table 3, except that all parameters cited in Table 2 are listed. Throughout Table 4,  $\tau_1$  and  $\tau_2$  have been fixed at 2 and 11, respectively, the lowest and highest observed ages. Notice particularly the key parameters of this paper:  $y_1$ ,  $y_2$ ,  $a$ , and  $b$ . These reveal the essence of the apparent problems in Table 3. For example, the initial point No. 1 is obviously ill-chosen because the parameter  $y_1$  (weight at age 2) surely should be somewhere near 100 g, rather than 347.7 g as shown. (See the data at the start of this section.) More importantly, the convergence problem for model (1) at point No. 4 now comes into focus. Table 4 shows that  $b$  is nearly 0 at this point. In fact,  $p$  becomes positive infinite and  $t_0$  becomes negative infinite as  $b$  approaches 0. Using model (15), rather than model (1), the search algorithm easily finds point No. 5, where  $S_1$  is minimized for all four parameters in (15). Explicitly, starting at point No. 1, the algorithm requires fewer than 300 calculations of  $S_1$  to locate point No. 5. This point has the remarkable property that, while it determines just another curve (15), it corresponds to no curve at all of the form (1). This happens because  $t_0$  is not defined at point No. 5.

In terms of the regions in Fig. 1, the search algorithm starts at point No. 1 on the lower boundary of region 2. It then moves across region 1 to locate a minimum for  $S_1$  at point No. 5 in region 8. Consequently, when guided by model (1), the search is doomed to failure. Curves in region 8 can't be described by model (1) because  $t_0$  is not defined there. Consequently, a legendary piece of comic advice to the lost traveller applies literally in this case: "Come to think of it, sir, you can't get there from here." This difficulty epitomizes the importance of the third guideline for choosing parameters, discussed in the last section.

Table 4 also illustrates the other two guidelines. In connection with the first, notice that the estimates for  $y_1$  and  $y_2$  in Table 4 are reasonably close to the observed sizes 99 and 539 g, respectively, at ages 2 and 11. Also, the estimates for  $z_1$  and  $z_2$  vary more with small changes in  $S_1$  than  $y_1$  and  $y_2$ . This confirms parameter behavior anticipated by the second guideline.

The analysis does not stop with the determination of point No. 5. It may be that a three-parameter submodel could be used to describe the data adequately. Earlier discussion shows that, for two particular such curves, a Pütter No. 2 ( $b = \frac{1}{2}$ ) should be chosen in preference to a Pütter No. 1 ( $b = 1$ ). Yet point No. 5, the optimum four-parameter model, has a small negative value for  $b$ . This suggests considering a Gompertz model ( $b = 0$ , Table 1), defined in the notation of this paper by (16). Point No. 6 in Table 4 shows the parameter estimates in this case. Since  $S_1$  is even smaller at point No. 6 than at point No. 3, a Gompertz model works even better than a Pütter No. 2.

For the cisco weight data, then, the really interesting choice lies between the general four-parameter model (15) and the three-parameter Gompertz submodel (16). To make this choice, consider the variance ratio defined by the residual sum of squares for model (16) divided by the sum of squares for (15):

$$\frac{(423.1 - 398.0)/1}{398.0/6} = 0.378.$$

If the models were linear in their parameters, this statistic would be  $F$ -distributed with 1 and 6 df under the null hypothesis that  $b = 0$ . (Here 1 df comes from the single parameter  $b$ , and  $10 - 4$  (i.e. 6) df comes from the 10 data points and the four parameters  $y_1$ ,  $y_2$ ,  $a$ , and  $b$ .) Even for non-linear models, the above variance ratio is approximately  $F$ -distributed when sample sizes are large. Under this approximation, one would reject the null hypothesis at the 5% level if the ratio exceeds

$$F_{0.05}(1,6) = 3.78.$$

The much smaller observed ratio (0.378) suggests retaining the null hypothesis  $b = 0$ , so that a Gompertz model appears adequate for the data. Once again, this conclusion could not be reached in the context of model (1), because  $t_0$  is not defined when  $b = 0$ .

This example illustrates how the data can be used objectively in selecting an appropriate model. The general procedure is to begin by locating an optimal four-parameter model (15). Then one can inspect the parameter estimates for  $a$  and  $b$  in the context of the models listed in Table 1. It may happen that the estimated curve lies close to a three-parameter (or even two-parameter) submodel. In this case, one might estimate the parameters for the submodel and use analysis of variance to make a final choice. Since the latter analysis is approximate at best, this procedure should be considered an informal process, rather than a rigorous statistical analysis. Very often, the action of the search algorithm itself helps to suggest which parameters are well defined. For example, in Table 5, the estimates for  $y_1$ ,  $y_2$ , and  $a$  are similar at points 3, 4, 5, and 6, while the estimate for  $b$  varies considerably as the sum of squares  $S_1$  changes only slightly. This reinforces the idea of fixing  $b$  at some prescribed value (like 0) and determining the other three parameters accordingly.

The example here focuses on problems associated with models in which the parameter pair  $(a, b)$  lies near the  $a$ -axis in Fig. 1. As the example illustrates, such models lead to difficulties in the definition of  $\tau_0$ . Other examples can easily be found in which the data indicate a parameter pair  $(a, b)$  near the  $b$ -axis. In such cases there are similar problems with the definition of  $y_\infty$ . These problems are precisely the ones which have led writers like Knight (1968) and Ricker (1979) to question the validity of asymptotic growth. Using the right parameters, it would probably be possible to stir up controversy in the literature associated with the boundary between each of the 10 pairs of adjacent regions in Fig. 1. Here is fertile ground for 20 scientific papers, including one for each rebuttal.

### Acknowledgments

I am grateful to Dr W. E. Ricker for valuable advice on the history of growth models and to Prof Carl Walters for helpful suggestions on the presentation of this paper. Prof Walters was also instrumental in my acquisition of the Apple II microcomputer on which this work was carried out. Without that tool, I would not have made the discoveries described here, let alone completed the data analysis.

- ALLEN, K. R. 1966. A method of fitting growth curves of the von Bertalanffy type to observed data. *J. Fish. Res. Board Can.* 23: 163–179.
- FABENS, A. J. 1965. Properties and fitting of the von Bertalanffy growth curve. *Growth* 29: 265–289.
- FLETCHER, R. I. 1973. A synthesis of deterministic growth laws. Univ. Rhode Island Sch. Oceanogr. Kingston, RI.
- KNIGHT, W. 1968. Asymptotic growth: an example of nonsense disguised as mathematics. *J. Fish Res. Board Can.* 25: 1030–1307.
- NELDER, J. A., AND R. MEAD. 1965. A simplex method for function minimization. *Comput. J.* 7: 308–313.
- PELLA, J. J., AND P. K. TOMLINSON. 1969. A generalized stock production model. *Inter-Am. Trop. Tuna Comm. Bull.* 13: 421–496.
- RICKER, W. E. 1975. Computation and interpretation of biological statistics of fish populations. *Fish. Res. Board Can. Bull.* 191: 382 p.
1979. Growth rates and models. Chapter 11, *Fish physiology*, Vol. 8. Academic Press, New York, NY.
- SCHAEFER, M. B. 1954. Some aspects of the dynamics of populations important to the management of the commercial marine fish-

eries. *Inter-Am. Trop. Tuna Comm. Bull.* 1: 25–56.

- SCHNUTE, J. 1981. A manual for easy nonlinear parameter estimation in fishery research, with interactive microcomputer programs. *Can. Spec. Publ. Fish. Aquat. Sci.* 59. (In press)
- SCHNUTE, J., AND D. FOURNIER. 1980. A new approach to length–frequency analysis: growth structure. *Can. J. Fish. Aquat. Sci.* 37: 1337–1351.
- WALFORD, L. A. 1946. A new graphic method of describing the growth of animals. *Biol. Bull.* 90: 141–147.

### Appendix A. Solutions to Problems (I) and (II)

The solution  $Y$  to Problem (I) consists of equations (15)–(18). The corresponding solution  $Z$  is:

Case 1:  $a \neq 0, b \neq 0$

$$(A1) \quad Z(t) = \frac{a}{b} \frac{y_2^b - y_1^b}{y_2^{b\tau_1} [e^{a(t-\tau_1)} - 1] + y_1^{b\tau_2} [1 - e^{-a(t-\tau_2)}]},$$

Case 2:  $a \neq 0, b = 0$

$$(A2) \quad Z(t) = \frac{a \log(y_2/y_1) e^{-at}}{e^{-a\tau_1} - e^{-a\tau_2}},$$

Case 3:  $a = 0, b \neq 0$

$$(A3) \quad Z(t) = \frac{1}{b} \frac{y_2^b - y_1^b}{y_2^b(t - \tau_1) - y_1^b(t - \tau_2)},$$

Case 4:  $a = 0, b = 0$

$$(A4) \quad Z(t) = \frac{\log(y_2/y_1)}{\tau_2 - \tau_1}.$$

The solutions  $Y$  and  $Z$  to Problem (III) are given similarly in the equations:

Case 1:  $a \neq 0, b \neq 0$

$$(A5) \quad Y(t) = y_1 \left[ \frac{a + bz_1(1 - e^{-a(t-\tau_1)})^{1/b}}{a} \right],$$

$$(A6) \quad Z(t) = \frac{az_1 e^{-a(t-\tau_1)}}{a + bz_1(1 - e^{-a(t-\tau_1)})},$$

Case 2:  $a \neq 0, b = 0$

$$(A7) \quad Y(t) = y_1 \exp \left[ \frac{z_1(1 - e^{-a(t-\tau_1)})}{a} \right],$$

$$(A8) \quad Z(t) = z_1 e^{-a(t-\tau_1)},$$

Case 3:  $a = 0, b \neq 0$

$$(A9) \quad Y(t) = y_1 \left[ 1 + bz_1(t - \tau_1) \right]^{1/b},$$

$$(A10) \quad Z(t) = z_1/[1 + bz_1(t - \tau_1)],$$

Case 4:  $a = 0, b = 0$

$$(A11) \quad Y(t) = y_1 e^{z_1(t-\tau_1)},$$

$$(A12) \quad Z(t) = z_1.$$

To prove the 16 formulas (15)–(18) and (A1)–(A12), begin with problem (II) when  $a \neq 0$  and  $b \neq 0$ . Equation (10) can be written in the form

$$(A13) \quad \frac{1}{a} \left( \frac{b}{a + bZ} - \frac{1}{Z} \right) dZ = dt.$$

Integrating (A13) with the initial condition (14) gives

$$(A14) \quad \frac{1}{a} \log \left( \frac{a + bZ}{Z} \cdot \frac{z_1}{a + bz_1} \right) = t - \tau_1.$$

It is possible to solve for  $Z$  in (A14), and the result is (A6). Furthermore, it turns out that both the right and left sides of (A6) are logarithmic derivatives. Explicitly, (A6) can be written as

$$(A15) \quad \frac{d}{dt} \log Y = \frac{1}{b} \frac{d}{dt} \log \left[ a + bz_1(1 - e^{-a(t-\tau_1)}) \right].$$

Integrating (A15) with the initial condition (13) gives

$$(A16) \quad \log \left( \frac{Y}{y_1} \right) = \frac{1}{b} \log \left[ \frac{a + bz_1(1 - e^{-a(t-\tau_1)})}{a} \right].$$

Solving (A16) for  $Y$  gives (A5). This completes the proof of (A5)–(A6).

Equations (A7)–(A12) can now be obtained from (A5)–(A6) by taking limits as the parameters  $a$  or  $b$  or both tend to 0. The proofs depend on the two results

$$(A17) \quad \lim_{a \rightarrow 0} (1 - e^{-a\alpha})/a = \alpha,$$

$$(A18) \quad \lim_{b \rightarrow 0} (1 + \beta b)^{1/b} = e^\beta,$$

where  $\alpha$  is an expression independent of  $a$ , and  $\beta$  is similarly independent of  $b$ . For example, equations (A9)–(A10) follow from (A5)–(A6) with the aid of (A17). Similarly, (A7) follows from (A5) because of (A18). The transitions from (A6) to (A8) to (A12) are obvious, and (A11) follows from (A7) by (A17). This completes the proof of solutions (A5)–(A12) for Problem (II).

To obtain solutions to Problem (I), notice from (A5) and the end condition (12) that

$$(A19) \quad y_2 = y_1 \left[ \frac{a + bz_1(1 - e^{-a(\tau_2-\tau_1)})}{a} \right].$$

Solving (A19) for  $z_1$  gives

$$(A20) \quad z_1 = \frac{a(y_2^b - y_1^b)}{by_1^b(1 - e^{-a(\tau_2-\tau_1)})}.$$

Substituting (A20) into (A5) and (A6) gives the general solutions (15) and (A1) to Problem (I). The remaining equations (16)–(18) and (A2)–(A4) are limiting forms of (15) and (A1). The transitions require the use of

$$(A21) \quad \lim_{b \rightarrow 0} \left[ 1 + \beta(\gamma^b - 1) \right]^{1/b} = e^{\beta \log \gamma},$$

where  $\beta$  and  $\gamma$  are independent of  $b$ . A related limit, essentially a special case of (A21), is

$$(A22) \quad \lim_{b \rightarrow 0} \frac{y_2^b - y_1^b}{b} = \log(y_2/y_1).$$

Using (A17), (A21), and (A22), one can obtain (16)–(18) and (A2)–(A4) from (15) and (A1). This completes the derivation of solutions to Problem (I).

## Appendix B. Parameter Relationships

To obtain the expressions (22)–(23) for  $z_1$  and  $z_2$ , simply substitute  $t = \tau_1$  and  $t = \tau_2$  in (A1)–(A4). Also, if there is a time  $\tau_0$  when  $y(\tau_0) = 0$ , then by (15)

$$(B1) \quad y_1^b + (y_2^b - y_1^b) \frac{1 - e^{-a(\tau_0-\tau_1)}}{1 - e^{-a(\tau_2-\tau_1)}} = 0,$$

provided that  $b > 0$ . Solving (B1) for  $\tau_0$  gives the first expression (24). Similarly, the second expression (24) follows from (17). Notice that  $\tau_0$  in (24) is not defined for  $b = 0$  because, in either case, the fraction has denominator 0 and numerator nonzero.

From the definition (6) for  $Z$ , it follows by differentiation that

$$(B2) \quad \frac{dZ}{dt} = \frac{1}{Y} \frac{d^2Y}{dt^2} - Z^2.$$

Notice that equations (10) and (B2) both describe  $dZ/dt$ . Equating right-hand sides and solving for  $d^2Y/dt^2$  gives

$$(B3) \quad \frac{d^2Y}{dt^2} = YZ[(1 - b)Z - a].$$

Incidentally, by (9), this proves (8). Now if there is a time  $\tau^*$  when  $d^2Y/dt^2$  vanishes, it follows from (B3) that at least one of three possibilities holds:

$$(B4) \quad y^* = 0,$$

$$(B5) \quad z^* = 0,$$

$$(B6) \quad (1 - b)z^* - a = 0.$$

The possibility (B4) is eliminated by requiring a positive size  $y^*$ . Also (B5) can be eliminated because  $Z = 0$  gives a singu-

lar point for the system (9)–(10); consequently, if  $Z$  ever vanishes, then  $Y$  and  $Z$  must be constant. This leaves (B6) and proves (28).

Define the quantity

$$(B7) \quad k = Y^b (a + bZ).$$

Apparently  $k$  is a function of time  $t$ , since  $Y$  and  $Z$  are.

However, differentiating  $k$  gives

$$\begin{aligned} \frac{dk}{dt} &= bY^b \left[ (a + bZ) \frac{1}{Y} \frac{dY}{dt} + \frac{dZ}{dt} \right] \\ &= bY^b [Z(a + bZ) - Z(a + bZ)] \\ &= 0. \end{aligned}$$

Consequently,  $k$  is a constant; in particular

$$(B8) \quad k = y_1^b (a + bz_1).$$

Combining (B8) and (22) gives the alternative form

$$(B9) \quad k = a \frac{y_2^b e^{a\tau_2} - y_1^b e^{a\tau_1}}{e^{a\tau_2} - e^{a\tau_1}},$$

when  $a \neq 0$ . It follows from (B7) that

$$(B10) \quad Y = \left[ k/(a + bZ) \right]^{1/b},$$

$$(B11) \quad Z = (kY^{-b} - a)/b,$$

when  $b \neq 0$ . Equations (B10) and (B11) allow  $Y(t)$  and  $Z(t)$  to be expressed in terms of each other. Equivalently, these equations describe trajectories in the  $YZ$ -plane for the system (9)–(10).

In particular (B10) implies that

$$(B12) \quad y^* = [k/(a + bz^*)].$$

The expression (27) for  $y^*$  follows by substituting (28) and (B9) into (B12). As usual, the second expression (27) is a limiting form of the first, where, in this case, (A21) is helpful. Once  $y^*$  is found, then  $\tau^*$  can be determined from (15) by solving

$$y^* = \left[ y_1^b + (y_2^b - y_1^b) \frac{1 - e^{-a(\tau^* - \tau_1)}}{1 - e^{-a(\tau_2 - \tau_1)}} \right]^{1/b}$$

for  $\tau^*$ . This gives (26), where the second expression follows from the first by (A22).

Finally suppose that

$$y_\infty = \lim_{t \rightarrow \infty} Y(t)$$

exists. Then the limiting value of  $Z$  must be 0, since the growth rate must slow to a stop as the limiting size is reached. Consequently, from (B10)

$$(B13) \quad y_\infty = (k/a)^{1/b}.$$

The formula (25) follows from (B9) and (B13). This completes the proof of all parametric relationships (22)–(28). The final two equations (29) and (30) are simple consequences of (24)–(27). Furthermore, (33) can be proved by combining (B11), (B13), and (6).

## Appendix C. Curve Shapes

Substituting the expression (B11) for  $Z$  into (8) gives

$$(C1) \quad \frac{d^2 Y}{dt^2} = \frac{dY}{dt} \frac{a}{b} (k^* Y^{-b} - 1),$$

where the constant  $k^*$  is

$$(C2) \quad k^* = (1 - b) \frac{e^{a\tau_2} y_2^b - e^{a\tau_1} y_1^b}{e^{a\tau_2} - e^{a\tau_1}}.$$

It is shown in Appendix B that  $Z \neq 0$  on any curve; in fact, from the assumptions  $y_2 > y_1 > 0$  and  $\tau_2 > \tau_1$ , it can be proved that  $Z > 0$  on the entire curve. Consequently, for positive sizes  $Y > 0$ , it follows that  $dY/dt > 0$ . The sign of the curvature (C1), then, is determined by the two factors  $a/b$  and  $k^* Y^{-b} - 1$ . In particular, the curvature is independent of  $Y$  if  $k^* = 0$ . From (C2) this happens if either

$$(C3) \quad b = 1, \text{ or}$$

$$(C4) \quad e^{a\tau_2} y_2^b = e^{a\tau_1} y_1^b.$$

Equation (C4) is equivalent to (34). Consequently (C3) and (C4) define the two broken lines in Fig. 1.

To prove the correspondence between Fig. 1 and Fig. 2, it is necessary to answer the following two questions for each of the eight regions in Fig. 1:

- For what values of  $t$  is the expression (15) for  $Y$  defined?
- For what values of  $Y$  is the curvature (C1) positive, and for what values is it negative?

The answer to (a) hinges on solving the inequality

$$(C5) \quad y_1^b + (y_2^b - y_1^b) \frac{1 - e^{-a(t - \tau_1)}}{1 - e^{-a(\tau_2 - \tau_1)}} > 0$$

for  $t$ . Similarly, the answer to (b) involves solving the pair of inequalities

$$(C6) \quad \frac{a}{b} (k^* Y^{-b} - 1) \geq 0$$

for  $Y$ . Equivalent formulations for (C5) and (C6) are

$$(C7) \quad a \left[ e^{a\tau_2} y_2^b - e^{a\tau_1} y_1^b - (y_2^b - y_1^b) e^{a(\tau_1 + \tau_2 - t)} \right] > 0,$$

$$(C8) \quad b \left[ (1 - b) (e^{a\tau_2} y_2^b - e^{a\tau_1} y_1^b) - (e^{a\tau_2} - e^{a\tau_1}) Y^b \right] \geq 0.$$

The details of working through (C7)–(C8) in each of the eight regions are too lengthy to present here, but they are straightforward. For example, in region 8 it turns out that

$$a > 0,$$

$$e^{a\tau_2}y_2^b - e^{a\tau_1}y_1^b > 0,$$

$$y_2^b - y_1^b < 0.$$

These inequalities show that (C7) holds for any  $t$ ; consequently the curve in region 8 is defined for all time. By contrast, in regions 1, 2, 3, and 4 (C7) holds when  $t > t_0$ . Similarly (C7) is always true in region 5, but holds only for  $t < t_0$  in regions 6 and 7. The analysis of (C8) follows in much the same way. For example, in region 2 the inequalities

$$1 - b < 0,$$

$$e^{a\tau_2}y_2^b - e^{a\tau_1}y_1^b > 0,$$

$$e^{a\tau_2} - e^{a\tau_1} > 0,$$

guarantee that the left-hand side of (C8) is always negative. Consequently the curvature is always negative and the corresponding curve in Fig. 2 is convex downward.

To establish the role of  $y_\infty$  in Fig. 2, it is necessary to consider the limit of  $Y(t)$  in (15) as  $t \rightarrow -\infty$  or  $t \rightarrow +\infty$  whenever the curve is defined for arbitrarily large negative or positive  $t$ . After the relevant range for  $t$  is established from (C7), the required limits in (15) follow easily from the properties of  $e^{-at}$ .

## Appendix D. Computer Methods

The data analysis in this paper was done with the aid of a general function minimization program written in BASIC by Schnute (1981). The program employs the simplex search method of Nelder and Mead (1965). It will run on many computers, including the Apple II desk-top microcomputer for which it was developed. Schnute's (1981) documentation shows not only how to minimize the particular functions (37) and (38), but also how to employ this software in nonlinear estimation for any small model like the one discussed here. Typically, the practitioner needs to write just a few lines of code describing the model of interest and to key in the relevant data. Of course, the caveats in this paper still apply. For example, the model must be posed so that the search can proceed fruitfully.

## This article has been cited by:

1. Hubert Du Pontavice, Marine Randon, Sigrid Lehuta, Youen Vermard, Marie Savina-Rolland. 2018. Investigating spatial heterogeneity of von Bertalanffy growth parameters to inform the stock structuration of common sole, *Solea solea*, in the Eastern English Channel. *Fisheries Research* **207**, 28-36. [[Crossref](#)]
2. Luana Granada, Sofia Lopes, Sara C. Novais, Marco F.L. Lemos. 2018. Modelling integrated multi-trophic aquaculture: Optimizing a three trophic level system. *Aquaculture* **495**, 90-97. [[Crossref](#)]
3. Lisa E. Ailloud, Todd Gedamke, John M. Hoenig. 2018. Short-Term Pain and Long-Term Gain: Using Phased-In Minimum Size Limits to Rebuild Stocks-the Pacific Bluefin Tuna Example. *Transactions of the American Fisheries Society* **4**. . [[Crossref](#)]
4. René Zambrano, Gabriela Galindo-Cortes, E Alberto Aragón-Noriega. 2018. Comparison of growth pattern of male *Ucides occidentalis* (Ortmann, 1897) (Brachyura: Ocypodidae) based on a combination of commercial catches and non-commercial data. *Journal of Crustacean Biology* **38**:4, 429-434. [[Crossref](#)]
5. Airam-Nauzet Sarmiento-Lezcano, Raúl Triay-Portella, José Juan Castro, Uriel Rubio-Rodríguez, José G. Pajuelo. 2018. Age-based life-history parameters of the mesopelagic fish *Notoscopelus resplendens* (Richardson, 1845) in the Central Eastern Atlantic. *Fisheries Research* **204**, 412-423. [[Crossref](#)]
6. Steven H. Ferguson, Xinhua Zhu, Brent G. Young, David J. Yurkowski, Gregory W. Thiemann, Aaron T. Fisk, Derek C.G. Muir. 2018. Geographic variation in ringed seal (*Pusa hispida*) growth rate and body size. *Canadian Journal of Zoology* **96**:7, 649-659. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)] [[Supplemental Material](#)]
7. P. Charvet, F. M. Santana, K. L. De Lima, R. Lessa. 2018. Age and growth of the endemic Xingu River stingray *Potamotrygon leopoldi* validated using fluorescent dyes. *Journal of Fish Biology* **92**:6, 1985-1999. [[Crossref](#)]
8. Cristina Maria Iordan, Francesca Verones, Francesco Cherubini. 2018. Integrating impacts on climate change and biodiversity from forest harvest in Norway. *Ecological Indicators* **89**, 411-421. [[Crossref](#)]
9. Charles Breton, Pierre Blanchet, Ben Amor, Robert Beauregard, Wen-Shao Chang. 2018. Assessing the Climate Change Impacts of Biogenic Carbon in Buildings: A Critical Review of Two Main Dynamic Approaches. *Sustainability* **10**:6, 2020. [[Crossref](#)]
10. Xi He, John C. Field. 2018. Effects of recruitment variability and fishing history on estimation of stock-recruitment relationships: Two case studies from U.S. West Coast fisheries. *Fisheries Research* . [[Crossref](#)]
11. Sergio G. Castillo-Vargasmachuca, Jesús T. Ponce-Palafox, Eulalio Arámbul-Muñoz, Guillermo Rodríguez-Domínguez, Eugenio Alberto Aragón-Noriega. 2018. The spotted rose snapper ( *Lutjanus guttatus* Steindachner 1869) farmed in marine cages: review of growth models. *Reviews in Aquaculture* **10**:2, 376-384. [[Crossref](#)]
12. Perla J. Rivera-Velázquez, E. Alberto Aragón-Noriega, Guillermo Rodríguez-Domínguez, Raúl Pérez-González, Sergio G. Castillo-Vargasmachuca. 2018. Growth, maturity and

- mortality of the blue crab *Callinectes arcuatus* Ordway, 1863 (Decapoda, Portunidae) in a Mexican coastal lagoon. *Crustaceana* **91**:6, 659-675. [[Crossref](#)]
13. Matthew D. Taylor, Colin Creighton. 2018. Estimating the Potential Fishery Benefits from Targeted Habitat Repair: a Case Study of School Prawn (*Metapenaeus macleayi*) in the Lower Clarence River Estuary. *Wetlands* **11**. . [[Crossref](#)]
14. C  il  n Minto, John Hinde, Rui Coelho. 2018. Including unsexed individuals in sex-specific growth models. *Canadian Journal of Fisheries and Aquatic Sciences* **75**:2, 282-292. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
15. Panagiotis Tsilifis, William J. Browning, Thomas E. Wood, Paul K. Newton, Roger G. Ghanem. 2018. The Stochastic Quasi-chemical Model for Bacterial Growth: Variational Bayesian Parameter Update. *Journal of Nonlinear Science* **28**:1, 371-393. [[Crossref](#)]
16. A. M. Roschanski, P. Barre, A. Escobar-Guti  rrez, J. P. Sampoux, H. Muylle, I. Thomas, K. J. Dehmer, E. Willner. Patterns of Spring Growth and Phenology in Natural Populations of *Lolium perenne* Under Contrasting Field Conditions 14-19. [[Crossref](#)]
17. Kai Moriguchi. 2018. An approach for deriving growth equations for quantities exhibiting cumulative growth based on stochastic interpretation. *Physica A: Statistical Mechanics and its Applications* **490**, 1150-1163. [[Crossref](#)]
18. Derek G. Bolser, Arnaud Gr  ss, Mark A. Lopez, Erin M. Reed, Ismael Mascare  nas-Osorio, Brad E. Erisman. 2018. The influence of sample distribution on growth model output for a highly-exploited marine fish, the Gulf Corvina ( *Cynoscion othonopterus* ). *PeerJ* **6**, e5582. [[Crossref](#)]
19. B Lee, BQ Mann. 2017. Age and growth of narrow-barred Spanish mackerel *Scomberomorus commerson* in the coastal waters of southern Mozambique and KwaZulu-Natal, South Africa. *African Journal of Marine Science* **39**:4, 397-407. [[Crossref](#)]
20. H. Renard, D. Maro, S. Le Diz  s, A. Escobar-Guti  rrez, C. Voiseux, L. Solier, D. H  bert, M. Rozet, C. Cossonnet, R. Barillot. 2017. Tritium forms discrimination in ryegrass under constant tritium exposure: From seed germination to seedling autotrophy. *Journal of Environmental Radioactivity* **177**, 194-205. [[Crossref](#)]
21. E. Alberto Arag  n-Noriega, Jaime E. Mendivil-Mendoza, Edgar Alc  ntara-Razo, Wenceslao Valenzuela-Quin  n  n, Jos   A. F  lix-Ortiz. 2017. Multi-criteria approach to estimate the growth curve in the marine shrimp, *Penaeus vannamei* Boone, 1931 (Decapoda, Penaeidae). *Crustaceana* **90**:11-12, 1517-1531. [[Crossref](#)]
22. Denis G. Dumas, Daniel M. McNeish. 2017. Dynamic Measurement Modeling: Using Nonlinear Growth Models to Estimate Student Learning Capacity. *Educational Researcher* **46**:6, 284-292. [[Crossref](#)]
23. Elo  se C. Ashworth, S. Alex Hesp, Norman G. Hall. 2017. A new proportionality-based back-calculation approach, which employs traditional forms of growth equations, improves estimates of length at age. *Canadian Journal of Fisheries and Aquatic Sciences* **74**:7, 1088-1099. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)] [[Supplemental Material](#)]
24. Lisa E. Ailloud, Matthew V. Lauretta, Alex R. Hanke, Walter J. Golet, Robert J. Allman, Matthew R. Siskey, David H. Secor, John M. Hoenig. 2017. Improving growth estimates for Western Atlantic bluefin tuna using an integrated modeling approach. *Fisheries Research* **191**, 17-24. [[Crossref](#)]

25. . Age and Growth 89-110. [[Crossref](#)]
26. Alexander Barrios, Bruno Ernande, Kelig Mahé, Verena Trenkel, Marie-Joëlle Rochet. 2017. Utility of mixed effects models to inform the stock structure of whiting in the Northeast Atlantic Ocean. *Fisheries Research* **190**, 132-139. [[Crossref](#)]
27. Khayaletu Ntushelo. 2017. Effect of Salicylic Acid on the Growth and Chemical Responses of *Pectobacterium carotovorum* subsp. *carotovorum*. *Pakistan Journal of Biological Sciences* **20**:6, 278-288. [[Crossref](#)]
28. Eloïse C. Ashworth, Norman G. Hall, S. Alex Hesp, Peter G. Coulson, Ian C. Potter. 2017. Age and growth rate variation influence the functional relationship between somatic and otolith size. *Canadian Journal of Fisheries and Aquatic Sciences* **74**:5, 680-692. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)] [[Supplemental Material](#)]
29. Lucian Bentea, Murielle A. Watzky, Richard G. Finke. 2017. Sigmoidal Nucleation and Growth Curves Across Nature Fit by the Finke–Watzky Model of Slow Continuous Nucleation and Autocatalytic Growth: Explicit Formulas for the Lag and Growth Times Plus Other Key Insights. *The Journal of Physical Chemistry C* **121**:9, 5302-5312. [[Crossref](#)]
30. Terrance J. Quinn, Richard B. Deriso, Philip D. Ganz. 2017. Combining the Cohen-Fishman growth increment model with a Box-Cox transformation. *Fisheries Research* **187**, 11-21. [[Crossref](#)]
31. Cole C. Monnahan, James T. Thorson, Trevor A. Branch. 2017. Faster estimation of Bayesian models in ecology using Hamiltonian Monte Carlo. *Methods in Ecology and Evolution* **8**:3, 339-348. [[Crossref](#)]
32. Miguel Montoro Girona, Sergio Rossi, Jean-Martin Lussier, Denis Walsh, Hubert Morin. 2017. Understanding tree growth responses after partial cuttings: A new approach. *PLOS ONE* **12**:2, e0172653. [[Crossref](#)]
33. Christoph Fischer, Egbert Schönfelder. 2017. A modified growth function with interpretable parameters applied to the age–height relationship of individual trees. *Canadian Journal of Forest Research* **47**:2, 166-173. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
34. Peter D. Tonner, Cynthia L. Darnell, Barbara E. Engelhardt, Amy K. Schmid. 2017. Detecting differential growth of microbial populations with Gaussian process regression. *Genome Research* **27**:2, 320-333. [[Crossref](#)]
35. Daniel McNeish, Denis Dumas. 2017. Nonlinear Growth Models as Measurement Models: A Second-Order Growth Curve Model for Measuring Potential. *Multivariate Behavioral Research* **52**:1, 61-85. [[Crossref](#)]
36. Luis Diaz-Balteiro, Luiz C. E. Rodríguez. Influence of Carbon Sequestration in an Optimal Set of Coppice Rotations for Eucalyptus Plantations 187-204. [[Crossref](#)]
37. Mary E. Matta, Cindy A. Tribuzio, David A. Ebert, Kenneth J. Goldman, Christopher M. Gburski. Age and Growth of Elasmobranchs and Applications to Fisheries Management and Conservation in the Northeast Pacific Ocean 179-220. [[Crossref](#)]
38. Vincent Lugert, Jens Tetens, Georg Thaller, Carsten Schulz, Joachim Krieter. 2017. Finding suitable growth models for turbot (*Scophthalmus maximus* L.) in aquaculture 1 (length application). *Aquaculture Research* **48**:1, 24-36. [[Crossref](#)]



39. Eugenio Alberto Aragón-Noriega. 2016. Model Selection to Describe the Growth of the Squalid *Callista Megapitaria squalida* from the Eastern Gulf of California. *Journal of Shellfish Research* **35**:4, 747-755. [[Crossref](#)]
40. 2016. Individual growth estimation of *Ucides occidentalis* (Brachyura, Ocypodidae) in the Gulf of Guayaquil, Ecuador by indirect methods and multi-model selection. *Crustaceana* **89**:13, 1509-1524. [[Crossref](#)]
41. Ian Doonan, Kath Large, Alistair Dunn, Scott Rasmussen, Craig Marsh, Sophie Mormede. 2016. Casal2: New Zealand's integrated population modelling tool. *Fisheries Research* **183**, 498-505. [[Crossref](#)]
42. Philip M. Wilkinson, Thomas R. Rainwater, Allan R. Woodward, Erin H. Leone, Cameron Carter. 2016. Determinate Growth and Reproductive Lifespan in the American Alligator ( *Alligator mississippiensis* ): Evidence from Long-term Recaptures. *Copeia* **104**:4, 843-852. [[Crossref](#)]
43. Luana Granada, Nádia Sousa, Sofia Lopes, Marco F. L. Lemos. 2016. Is integrated multitrophic aquaculture the solution to the sectors' major challenges? - a review. *Reviews in Aquaculture* **8**:3, 283-300. [[Crossref](#)]
44. André E. Punt, Malcolm Haddon, Richard McGarvey. 2016. Estimating growth within size-structured fishery stock assessments: What is the state of the art and what does the future look like?. *Fisheries Research* **180**, 147-160. [[Crossref](#)]
45. R.I.C. Chris Francis. 2016. Growth in age-structured stock assessment models. *Fisheries Research* **180**, 77-86. [[Crossref](#)]
46. Kevin R. Piner, Hui-Hua Lee, Mark N. Maunder. 2016. Evaluation of using random-at-length observations and an equilibrium approximation of the population age structure in fitting the von Bertalanffy growth function. *Fisheries Research* **180**, 128-137. [[Crossref](#)]
47. Dale S. Kolody, J.Paige Eveson, Richard M. Hillary. 2016. Modelling growth in tuna RFMO stock assessments: Current approaches and challenges. *Fisheries Research* **180**, 177-193. [[Crossref](#)]
48. Brett T. van Poorten, Carl J. Walters. 2016. How can bioenergetics help us predict changes in fish growth patterns?. *Fisheries Research* **180**, 23-30. [[Crossref](#)]
49. Yi Xu, Steven L.H. Teo, Kevin R. Piner, Kuo-Shu Chen, R.J. David Wells. 2016. Using an approximate length-conditional approach to estimate von Bertalanffy growth parameters of North Pacific albacore (*Thunnus alalunga*). *Fisheries Research* **180**, 138-146. [[Crossref](#)]
50. Xi He, John C. Field, Donald E. Pearson, Lyndsey S. Lefebvre. 2016. Age sample sizes and their effects on growth estimation and stock assessment outputs: Three case studies from U.S. West Coast fisheries. *Fisheries Research* **180**, 92-102. [[Crossref](#)]
51. Carolina V. Minte-Vera, Mark N. Maunder, John M. Casselman, Steven E. Campana. 2016. Growth functions that incorporate the cost of reproduction. *Fisheries Research* **180**, 31-44. [[Crossref](#)]
52. Cole C. Monnahan, Kotaro Ono, Sean C. Anderson, Merrill B. Rudd, Allan C. Hicks, Felipe Hurtado-Ferro, Kelli F. Johnson, Peter T. Kuriyama, Roberto R. Licandeo, Christine C. Stawitz, Ian G. Taylor, Juan L. Valero. 2016. The effect of length bin width on growth

- estimation in integrated age-structured stock assessments. *Fisheries Research* **180**, 103-112. [[Crossref](#)]
53. Kai Lorenzen. 2016. Toward a new paradigm for growth modeling in fisheries stock assessments: Embracing plasticity and its consequences. *Fisheries Research* **180**, 4-22. [[Crossref](#)]
54. Ruben H. Roa-Ureta. 2016. Size-dependent mortality rate profiles. *Journal of Theoretical Biology* **402**, 107-116. [[Crossref](#)]
55. John M. Erhardt, Dennis L. Scarnecchia. 2016. Growth Model Selection and its Application for Characterizing Life History of a Migratory Bull Trout ( *Salvelinus confluentus* ) Population. *Northwest Science* **90**:3, 328-339. [[Crossref](#)]
56. R. Lessa, H. A. Andrade, K. L. De Lima, F. M. Santana. 2016. Age and growth of the midwater crocodile shark *Pseudocarcharias kamoharai*. *Journal of Fish Biology* **89**:1, 371-385. [[Crossref](#)]
57. Heather M. Christiansen, Steven E. Campana, Aaron T. Fisk, Jeremy Cliff, Sabine P. Wintner, Sheldon F. J. Dudley, Lisa A. Kerr, Nigel E. Hussey. 2016. Using bomb radiocarbon to estimate age and growth of the white shark, *Carcharodon carcharias*, from the southwestern Indian Ocean. *Marine Biology* **163**:6. . [[Crossref](#)]
58. Awatef Trabelsi, Andrzej Jaworski, Ewa Kamler, Jean-Noël Gardeur, Fabrice Teletchea, Habib Ayadi, Pascal Fontaine. 2016. The effect of hatching time on the bioenergetics of northern pike ( *Esox lucius* ) larvae from a single egg batch during the endogenous feeding period. *Fish Physiology and Biochemistry* **42**:2, 593-606. [[Crossref](#)]
59. A. Rotger, J.M. Igual, J.J. Smith, G. Tavecchia. 2016. Relative role of population density and climatic factors in shaping the body growth rate of Lilford's Wall Lizard ( *Podarcis lilfordi* ). *Canadian Journal of Zoology* **94**:3, 207-215. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)] [[Supplemental Material](#)]
60. Vincent Lugert, Georg Thaller, Jens Tetens, Carsten Schulz, Joachim Krieter. 2016. A review on fish growth calculation: multiple functions in fish production and their specific application. *Reviews in Aquaculture* **8**:1, 30-42. [[Crossref](#)]
61. Nancy L. Staub. 2016. The Age of Plethodontid Salamanders: A Short Review on Longevity. *Copeia* **104**:1, 118-123. [[Crossref](#)]
62. Iván Velázquez-Abunader, Karina Monsreal-Vela, Gaspar R. Poot-López. 2016. Model selection for determining the growth of juveniles and sub-adults of two species of shrimp (Decapoda, Penaeidae) in a tropical coastal lagoon. *Crustaceana* **89**:1, 29-45. [[Crossref](#)]
63. Benjamin Mos, Maria Byrne, Symon A. Dworjanyn. 2016. Biogenic acidification reduces sea urchin gonad growth and increases susceptibility of aquaculture to ocean acidification. *Marine Environmental Research* **113**, 39-48. [[Crossref](#)]
64. Carlos Humberto Hernández, Crisantema Hernández, Francisco Javier Martínez-Cordero, Nicolás Castañeda-Lomas, Guillermo Rodríguez-Domínguez, Albert G. J. Tacon, Eugenio Alberto Aragón-Noriega. 2016. Effect of Density at Harvest on the Growth Performance and Profitability of Hatchery-reared Spotted Rose Snapper, *Lutjanus guttatus* , Cultured in Floating Net Cages. *Journal of the World Aquaculture Society* **47**:1, 51-60. [[Crossref](#)]
65. Lisa J. Natanson, Li Ling Hamady, Brian J. Gervelis. 2016. Analysis of bomb radiocarbon data for common thresher sharks, *Alopias vulpinus*, in the northwestern Atlantic Ocean with revised growth curves. *Environmental Biology of Fishes* **99**:1, 39-47. [[Crossref](#)]

66. Arne Pommerening, Anders Muszta. 2016. Relative plant growth revisited: Towards a mathematical standardisation of separate approaches. *Ecological Modelling* **320**, 383-392. [[Crossref](#)]
67. R Petitet, L Avens, JC Castilhos, PG Kinas, L Bugoni. 2015. Age and growth of olive ridley sea turtles *Lepidochelys olivacea* in the main Brazilian nesting ground. *Marine Ecology Progress Series* **541**, 205-218. [[Crossref](#)]
68. . References 283-304. [[Crossref](#)]
69. Yi-Fu Wang, Tian-Xiang Yue, Zheng-Ping Du, Ming-Wei Zhao. 2015. Improving the accuracy of the height–diameter equation using the classified factors method. *Environmental Earth Sciences* **74**:8, 6471-6480. [[Crossref](#)]
70. Peter L. Horn. Southern hake ( *Merluccius australis* ) in New Zealand: biology, fisheries and stock assessment 101-125. [[Crossref](#)]
71. Christine C. Stawitz, Timothy E. Essington, Trevor A. Branch, Melissa A. Haltuch, Anne B. Hollowed, Paul D. Spencer. 2015. A state-space approach for detecting growth variation and application to North Pacific groundfish. *Canadian Journal of Fisheries and Aquatic Sciences* **72**:9, 1316-1328. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)] [[Supplemental Material](#)]
72. Roscinto Ian C. Lumbres, Young Jin Lee, Chung Weon Yun, Chang Duck Koo, Se Bin Kim, Yeong Mo Son, Kyeong Hak Lee, Hyun Kyu Won, Sung Cheol Jung, Yeon Ok Seo. 2015. DBH-height modeling and validation for *Acacia mangium* and *Eucalyptus pellita* in Korintiga Hutani Plantation, Kalimantan, Indonesia. *Forest Science and Technology* **11**:3, 119-125. [[Crossref](#)]
73. Esteban Gómez-García, Felipe Crecente-Campo, Marcos Barrio-Anta, Ulises Diéguez-Aranda. 2015. A disaggregated dynamic model for predicting volume, biomass and carbon stocks in even-aged pedunculate oak stands in Galicia (NW Spain). *European Journal of Forest Research* **134**:3, 569-583. [[Crossref](#)]
74. O. Tariche, J.G. Pajuelo, J.M. Lorenzo, A. Luque, J.A. Gonzalez. 2015. Age estimation and growth pattern of the grouper *Cephalopholis taeniops* (Epinephelidae) off the Cape Verde Archipelago, north-west Africa. *Journal of the Marine Biological Association of the United Kingdom* **95**:03, 599-609. [[Crossref](#)]
75. D.C. Camargo, F. Montoya, J.I. Córcoles, J.F. Ortega. 2015. Modeling the impacts of irrigation treatments on potato growth and development. *Agricultural Water Management* **150**, 119-128. [[Crossref](#)]
76. Alexandre M. Aires-da-Silva, Mark N. Maunder, Kurt M. Schaefer, Daniel W. Fuller. 2015. Improved growth estimates from integrated analysis of direct aging and tag–recapture data: An illustration with bigeye tuna (*Thunnus obesus*) of the eastern Pacific Ocean with implications for management. *Fisheries Research* **163**, 119-126. [[Crossref](#)]
77. Margit R. Wilhelm, Astrid Jarre, Coleen L. Moloney. 2015. Spawning and nursery areas, longitudinal and cross-shelf migrations of the *Merluccius capensis* stock in the northern Benguela. *Fisheries Oceanography* **24**, 31-45. [[Crossref](#)]
78. José Angel Hidalgo-De-La-Toba, Sergio Scarry González-peláez, Enrique Morales-Bojórquez, J. Jesús Bautista-Romero, Daniel Bernardo Lluch-Cota. 2015. Geoduck *Panopea generosa* Growth at Its Southern Distribution Limit in North America using a Multimodel Inference Approach. *Journal of Shellfish Research* **34**:1, 91-99. [[Crossref](#)]

79. Sergio Scarry González-Peláez, Enrique Morales-Bojórquez, Daniel Bernardo Lluch-Cota, Salvador Emilio Lluch-Cota, J. Jesús Bautista-Romero. 2015. Modeling Geoduck Growth: Multimodel Inference in *Panopea globosa* from the Southwestern Baja California Peninsula, Mexico. *Journal of Shellfish Research* **34**:1, 101-112. [[Crossref](#)]
80. Eugenio Alberto Aragón-Noriega, Luis Eduardo Calderon-Aguilera, Sergio Alejandro Pérez-Valencia. 2015. Modeling Growth of the Cortes Geoduck *Panopea globosa* from Unexploited and Exploited Beds in the Northern Gulf of California. *Journal of Shellfish Research* **34**:1, 119-127. [[Crossref](#)]
81. Enrique Morales-Bojórquez, Eugenio Alberto Aragón-Noriega, Hugo Aguirre-Villaseñor, Luis E. Calderon-Aguilera, Viridiana Y. Zepeda-Benitez. 2015. Selection of Models to Predict *Panopea globosa* Growth: Application of a Mixture Probability Distribution Function. *Journal of Shellfish Research* **34**:1, 129-136. [[Crossref](#)]
82. Florencia Doño, Santiago Montealegre-Quijano, Andrés Domingo, Paul G. Kinas. 2015. Bayesian age and growth analysis of the shortfin mako shark *Isurus oxyrinchus* in the Western South Atlantic Ocean using a flexible model. *Environmental Biology of Fishes* **98**:2, 517-533. [[Crossref](#)]
83. Chih-Shin Chen, Jing-Yu Chen, Chiao-Wen Lin. 2015. Variation in life-history traits for micro-cohorts of *Sepioteuthis lessoniana* in the waters off northern Taiwan. *Fisheries Science* **81**:1, 53-64. [[Crossref](#)]
84. A.K. Jaiswal, S. Jaiswal. Modelling the effects of natural antimicrobials as food preservatives 259-284. [[Crossref](#)]
85. D. Parker, A. J. Booth. 2015. Aspects of the biology and life history of largespot pompano, *Trachinotus botla*, in South Africa. *Marine and Freshwater Research* **66**:3, 247. [[Crossref](#)]
86. Lisa J. Natanson, Gregory B. Skomal. 2015. Age and growth of the white shark, *Carcharodon carcharias*, in the western North Atlantic Ocean. *Marine and Freshwater Research* **66**:5, 387. [[Crossref](#)]
87. R Sedmák, L Scheer. 2015. Properties and prediction accuracy of a sigmoid function of time-determinate growth. *iForest - Biogeosciences and Forestry* **8**:5, 631-637. [[Crossref](#)]
88. F. Dubeau, Y. Mir. 2014. Exponential Growth Model: From Horizontal to Linear Asymptote. *Communications in Statistics - Simulation and Computation* **43**:10, 2186-2204. [[Crossref](#)]
89. Michael Hellmair, Andrew P. Kinziger. 2014. Increased Extinction Potential of Insular Fish Populations with Reduced Life History Variation and Low Genetic Diversity. *PLoS ONE* **9**:11, e113139. [[Crossref](#)]
90. D. Biriá, E. Maghsoudi, R. Roostaazad. 2014. APPLICATION OF POWER LAW LOGISTIC MODEL TO GROWTH KINETICS OF *BACILLUS LICHENIFORMIS* MS3 ON A WATER-INSOLUBLE SUBSTRATE. *Chemical Engineering Communications* **201**:11, 1514-1525. [[Crossref](#)]
91. C. Shane Reese, Alyson G. Wilson. 2014. Discussion of 'Methods for planning repeated measures accelerated degradation tests'. *Applied Stochastic Models in Business and Industry* **30**:6, 674-676. [[Crossref](#)]

92. VY Zepeda-Benitez, E Morales-Bojórquez, J López-Martínez, A Hernández-Herrera. 2014. Growth model selection for the jumbo squid *Dosidicus gigas* from the Gulf of California, Mexico. *Aquatic Biology* **21**:3, 231-247. [[Crossref](#)]
93. Lisa J. Natanson, Douglas H. Adams, Megan V. Winton, Jasmine R. Maurer. 2014. Age and Growth of the Bull Shark in the Western North Atlantic Ocean. *Transactions of the American Fisheries Society* **143**:3, 732-743. [[Crossref](#)]
94. P. M. Kiffney, E. R. Buhle, S. M. Naman, G. R. Pess, R. S. Klett. 2014. Linking resource availability and habitat structure to stream organisms: an experimental and observational assessment. *Ecosphere* **5**:4, art39. [[Crossref](#)]
95. M. J. Kingsford, M. D. Finn, M. D. O'Callaghan, J. Atema, G. Gerlach. 2014. Planktonic Larval Duration, age and growth of *Ostorhinchus doederleini* (Pisces: Apogonidae) on the southern Great Barrier Reef, Australia. *Marine Biology* **161**:2, 245-259. [[Crossref](#)]
96. S. Dick, J. B. Shurin, E. B. Taylor. 2014. Replicate divergence between and within sounds in a marine fish: the copper rockfish ( *Sebastes caurinus* ). *Molecular Ecology* **23**:3, 575-590. [[Crossref](#)]
97. Richard S. McBride. The Continuing Role of Life History Parameters to Identify Stock Structure 77-107. [[Crossref](#)]
98. Daniel J. Stoessel. 2014. Age, growth, condition and reproduction of roach *Rutilus rutilus* (Teleostei : Cyprinidae), in south-eastern Australia. *Marine and Freshwater Research* **65**:3, 275. [[Crossref](#)]
99. Romain Barillot, Didier Combes, Sylvain Pineau, Pierre Huynh, Abraham J. Escobar-Gutiérrez. 2014. Comparison of the morphogenesis of three genotypes of pea (*Pisum sativum*) grown in pure stands and wheat-based intercrops. *AOB PLANTS* **6**. . [[Crossref](#)]
100. . REFERENCES 420-430. [[Crossref](#)]
101. Geoffrey Guest, Francesco Cherubini, Anders Hammer Strømman. 2013. Climate impact potential of utilizing forest residues for bioenergy in Norway. *Mitigation and Adaptation Strategies for Global Change* **18**:8, 1089-1108. [[Crossref](#)]
102. Troy A. Dexter, Michał Kowalewski. 2013. Jackknife-corrected parametric bootstrap estimates of growth rates in bivalve mollusks using nearest living relatives. *Theoretical Population Biology* **90**, 36-48. [[Crossref](#)]
103. Lisa J. Natanson, Brian J. Gervelis, Megan V. Winton, Li Ling Hamady, Simon J. B. Gulak, John K. Carlson. 2013. Validated age and growth estimates for *Carcharhinus obscurus* in the northwestern Atlantic Ocean, with pre- and post management growth comparisons. *Environmental Biology of Fishes* . [[Crossref](#)]
104. Megan V. Winton, Lisa J. Natanson, Jeff Kneebone, Gregor M. Cailliet, David A. Ebert. 2013. Life history of *Bathyrhaja trachura* from the eastern Bering Sea, with evidence of latitudinal variation in a deep-sea skate species. *Journal of the Marine Biological Association of the United Kingdom* 1-12. [[Crossref](#)]
105. Javier Castaño-Santamaría, Felipe Crecente-Campo, Juan Luis Fernández-Martínez, Marcos Barrio-Anta, José Ramón Obeso. 2013. Tree height prediction approaches for uneven-aged beech forests in northwestern Spain. *Forest Ecology and Management* **307**, 63-73. [[Crossref](#)]

106. M. R. Wilhelm, J.-P. Roux, C. L. Moloney, A. Jarre. 2013. Data from fur seal scats reveal when Namibian *Merluccius capensis* are hatched and how fast they grow. *ICES Journal of Marine Science* **70**:7, 1429-1438. [[Crossref](#)]
107. Felipe Crecente-Campo, José Javier Corral-Rivas, Benedicto Vargas-Larreta, Christian Wehenkel. 2013. Can random components explain differences in the height-diameter relationship in mixed uneven-aged stands?. *Annals of Forest Science* . [[Crossref](#)]
108. Ramazan Özçelik, Maria J. Diamantopoulou, Felipe Crecente-Campo, Unal Eler. 2013. Estimating Crimean juniper tree height using nonlinear regression and artificial neural network models. *Forest Ecology and Management* **306**, 52-60. [[Crossref](#)]
109. Bernard R. Parresol, Margaret S. Devall. 2013. Patterns of Diametric Growth in Stem-Analyzed Laurel Trees ( *Cordia alliodora* ) in a Panamanian Forest. *The Southwestern Naturalist* **58**:2, 170-178. [[Crossref](#)]
110. Richard D. Methot, Chantell R. Wetzel. 2013. Stock synthesis: A biological and statistical framework for fish stock assessment and fishery management. *Fisheries Research* **142**, 86-99. [[Crossref](#)]
111. Jason D. Baker, Thea C. Johanos, Tracy A. Wurth, Charles L. Littnan. 2013. Body growth in Hawaiian monk seals. *Marine Mammal Science* n/a-n/a. [[Crossref](#)]
112. Fay Helidoniotis, Malcolm Haddon. 2013. Growth Models for Fisheries: The Effect of Unbalanced Sampling Error On Model Selection, Parameter Estimation, and Biological Predictions. *Journal of Shellfish Research* **32**:1, 223-235. [[Crossref](#)]
113. François Dubeau, Youness Mir. 2013. Growth Models with Oblique Asymptote. *Mathematical Modelling and Analysis* **18**:2, 204-218. [[Crossref](#)]
114. Roscinto Ian C. Lumbres, Young Jin Lee, Feliciano G. Calora, Marissa R. Parao. 2013. Model fitting and validation of six height-DBH equations for *Pinus kesiya* Royle ex Gordon in Benguet Province, Philippines. *Forest Science and Technology* **9**:1, 45-50. [[Crossref](#)]
115. D Parker, AJ Booth, BQ Mann. 2013. A spatio-temporal assessment of the *Trachinotus botla* shore-fishery in KwaZulu-Natal, South Africa. *African Journal of Marine Science* **35**:1, 35-46. [[Crossref](#)]
116. Rosana Ouréns, Luis Flores, Luis Fernández, Juan Freire. 2013. Habitat and density-dependent growth of the sea urchin *Paracentrotus lividus* in Galicia (NW Spain). *Journal of Sea Research* **76**, 50-60. [[Crossref](#)]
117. Geoffrey Guest, Francesco Cherubini, Anders H. Strømman. 2013. Global Warming Potential of Carbon Dioxide Emissions from Biomass Stored in the Anthroposphere and Used for Bioenergy at End of Life. *Journal of Industrial Ecology* **17**:1, 20-30. [[Crossref](#)]
118. Thomas A. Ebert. Growth and Survival of Postsettlement Sea Urchins 83-117. [[Crossref](#)]
119. A. E. Punt, T. Huang, M. N. Maunder. 2013. Review of integrated size-structured models for stock assessment of hard-to-age crustacean and mollusc species. *ICES Journal of Marine Science* **70**:1, 16-33. [[Crossref](#)]
120. Jorge Alberto López-Rocha, Marcial Arellano—Martínez, Bertha Patricia Ceballos—Vázquez, Iván Velázquez—Abunader, Sheila Castellanos—Martínez, Esteban Torreblanca—Ramírez. 2012. Use of Length—Frequency Analysis for Growth Estimation of the California Two-



- Spotted Octopus *Octopus bimaculatus* Verrill 1883 of the Gulf of California. *Journal of Shellfish Research* **31**:4, 1173-1181. [[Crossref](#)]
121. R. Barillot, D. Combes, V. Chevalier, C. Fournier, A. J. Escobar-Gutierrez. 2012. How does pea architecture influence light sharing in virtual wheat-pea mixtures? A simulation study based on pea genotypes with contrasting architectures. *AoB Plants* **2012**:0, pls038-pls038. [[Crossref](#)]
122. Ryan M. Bright, Francesco Cherubini, Anders H. Strømman. 2012. Climate impacts of bioenergy: Inclusion of carbon cycle and albedo dynamics in life cycle impact assessment. *Environmental Impact Assessment Review* **37**, 2-11. [[Crossref](#)]
123. E. Lek, D. V. Fairclough, N. G. Hall, S. A. Hesp, I. C. Potter. 2012. Do the maximum sizes, ages and patterns of growth of three reef-dwelling labrid species at two latitudes differ in a manner conforming to the metabolic theory of ecology?. *Journal of Fish Biology* **81**:6, 1936-1962. [[Crossref](#)]
124. Francesco Cherubini, Geoffrey Guest, Anders H. Strømman. 2012. Application of probability distributions to the modeling of biogenic CO<sub>2</sub> fluxes in life cycle assessment. *GCB Bioenergy* **4**:6, 784-798. [[Crossref](#)]
125. Stephen J. Newman, Michael C. Mackie, Paul D. Lewis. 2012. Age-based demography and relative fisheries productivity of Spanish mackerel, *Scomberomorus commerson* (Lacepede) in Western Australia. *Fisheries Research* **129-130**, 46-60. [[Crossref](#)]
126. Peter G. Coulson, Ian C. Potter, Norman G. Hall. 2012. The biological characteristics of *Scorpius aequipinnis* (Kyphosidae), including relevant comparisons with those of other species and particularly of a heavily exploited congener. *Fisheries Research* **125-126**, 272-282. [[Crossref](#)]
127. Rolando Cruz-Vásquez, Guillermo Rodríguez-Domínguez, Edgar Alcántara-Razo, Eugenio Alberto Aragón-Noriega. 2012. Estimation of Individual Growth Parameters of the Cortes Geoduck *Panopea globosa* from the Central Gulf of California using a Multimodel Approach. *Journal of Shellfish Research* **31**:3, 725-732. [[Crossref](#)]
128. B. Rijal, A. R. Weiskittel, J. A. Kershaw. 2012. Development of regional height to diameter equations for 15 tree species in the North American Acadian Region. *Forestry* **85**:3, 379-390. [[Crossref](#)]
129. R Petitet, ER Secchi, L Avens, PG Kinas. 2012. Age and growth of loggerhead sea turtles in southern Brazil. *Marine Ecology Progress Series* **456**, 255-268. [[Crossref](#)]
130. KM Bailey, T Zhang, Ks Chan, SM Porter, AB Dougherty. 2012. Near real-time forecasting of recruitment from larval surveys: application to Alaska pollock. *Marine Ecology Progress Series* **452**, 205-217. [[Crossref](#)]
131. Yi-Jay Chang, Chi-Lu Sun, Yong Chen, Su-Zan Yeh. 2012. Modelling the growth of crustacean species. *Reviews in Fish Biology and Fisheries* **22**:1, 157-187. [[Crossref](#)]
132. Paula Ximena Hurtado, Sabine K. Schnabel, Alon Zaban, Merja Veteläinen, Elina Virtanen, Paul H. C. Eilers, Fred A. Eeuwijk, Richard G. F. Visser, Chris Maliepaard. 2012. Dynamics of senescence-related QTLs in potato. *Euphytica* **183**:3, 289-302. [[Crossref](#)]
133. YunSheng Wang, LiPing Gao, ZhengRong Wang, YaJun Liu, MeiLian Sun, DongQing Yang, ChaoLing Wei, Yu Shan, Tao Xia. 2012. Light-induced expression of genes involved in phenylpropanoid biosynthetic pathways in callus of tea (*Camellia sinensis* (L.) O. Kuntze). *Scientia Horticulturae* **133**, 72-83. [[Crossref](#)]

134. Francesco Cherubini, Anders H. Strømman, Edgar Hertwich. 2011. Effects of boreal forest management practices on the climate impact of CO<sub>2</sub> emissions from bioenergy. *Ecological Modelling* **223**:1, 59-66. [[Crossref](#)]
135. Roscinto Ian C. Lumbres, Young Jin Lee, Yeon Ok Seo, Sung Ho Kim, Jung Kee Choi, Woo Kyun Lee. 2011. Development and validation of nonlinear height-DBH models for major coniferous tree species in Korea. *Forest Science and Technology* **7**:3, 117-125. [[Crossref](#)]
136. . Bibliography 327-395. [[Crossref](#)]
137. Joseph M. O'Malley. 2011. Spatiotemporal variation in the population ecology of scaly slipper lobsters *Scyllarides squammosus* in the Northwestern Hawaiian Islands. *Marine Biology* **158**:8, 1887-1901. [[Crossref](#)]
138. Genoveva Cerdaneres-Ladrón De Guevara, Enrique Morales-Bojórquez, Rubén Rodríguez-Sánchez. 2011. Age and growth of the sailfish *Istiophorus platypterus* (Istiophoridae) in the Gulf of Tehuantepec, Mexico. *Marine Biology Research* **7**:5, 488-499. [[Crossref](#)]
139. A. Baer, C. Schulz, I. Traulsen, J. Krieter. 2011. Analysing the growth of turbot (*Psetta maxima*) in a commercial recirculation system with the use of three different growth models. *Aquaculture International* **19**:3, 497-511. [[Crossref](#)]
140. Rodrigo R. Barreto, Rosangela P. Lessa, Fabio H. Hazin, Francisco M. Santana. 2011. Age and growth of the blacknose shark, *Carcharhinus acronotus* (Poey, 1860) off the northeastern Brazilian Coast. *Fisheries Research* **110**:1, 170-176. [[Crossref](#)]
141. Joana Amaral Paulo, José Tomé, Margarida Tomé. 2011. Nonlinear fixed and random generalized height-diameter models for Portuguese cork oak stands. *Annals of Forest Science* **68**:2, 295-309. [[Crossref](#)]
142. Xinjun Chen, Huajie Lu, Bilin Liu, Yong Chen. 2011. Age, growth and population structure of jumbo flying squid, *Dosidicus gigas*, based on statolith microstructure off the Exclusive Economic Zone of Chilean waters. *Journal of the Marine Biological Association of the United Kingdom* **91**:01, 229-235. [[Crossref](#)]
143. Even Tjørve, Kathleen M.C. Tjørve. 2010. A unified approach to the Richards-model family for use in growth analyses: Why we need only two model forms. *Journal of Theoretical Biology* **267**:3, 417-425. [[Crossref](#)]
144. Juan Carlos Camargo García, Christoph Kleinn. 2010. Length curves and volume functions for guadua bamboo (*Guadua angustifolia* Kunth) for the coffee region of Colombia. *European Journal of Forest Research* **129**:6, 1213-1222. [[Crossref](#)]
145. L. F. Arauz, K. N. Neufeld, A. L. Lloyd, P. S. Ojiambo. 2010. Quantitative Models for Germination and Infection of *Pseudoperonospora cubensis* in Response to Temperature and Duration of Leaf Wetness. *Phytopathology* **100**:9, 959-967. [[Crossref](#)]
146. S. Cossington, S. A. Hesp, N. G. Hall, I. C. Potter. 2010. Growth and reproductive biology of the foxfish *Bodianus frenchii*, a very long-lived and monandric protogynous hermaphroditic labrid. *Journal of Fish Biology* no-no. [[Crossref](#)]
147. Luis Flores, Billy Ernst, Ana M. Parma. 2010. Growth pattern of the sea urchin, *Loxechinus albus* (Molina, 1782) in southern Chile: evaluation of growth models. *Marine Biology* **157**:5, 967-977. [[Crossref](#)]



148. G. G. Thorarinsdottir, L. Jacobson, S. A. Ragnarsson, E. G. Garcia, K. Gunnarsson. 2010. Capture efficiency and size selectivity of hydraulic clam dredges used in fishing for ocean quahogs (*Arctica islandica*): simultaneous estimation in the SELECT model. *ICES Journal of Marine Science* **67**:2, 345-354. [[Crossref](#)]
149. Felipe Crecente-Campo, Margarida Tomé, Paula Soares, Ulises Diéguez-Aranda. 2010. A generalized nonlinear mixed-effects height-diameter model for *Eucalyptus globulus* L. in northwestern Spain. *Forest Ecology and Management* **259**:5, 943-952. [[Crossref](#)]
150. Steven S. Montgomery, Chris T. Walsh, Malcolm Haddon, Caitlin L. Kesby, Daniel D. Johnson. 2010. Using length data in the Schnute Model to describe growth in a metapenaeid from waters off Australia. *Marine and Freshwater Research* **61**:12, 1435. [[Crossref](#)]
151. S. P. Griffiths, G. C. Fry, F. J. Manson, D. C. Lou. 2010. Age and growth of longtail tuna (*Thunnus tonggol*) in tropical and temperate waters of the central Indo-Pacific. *ICES Journal of Marine Science* **67**:1, 125-134. [[Crossref](#)]
152. Hsin-i Wu, Amit Chakraborty, Bai-Lian Li, Charles M. Kenerley. 2009. Formulating variable carrying capacity by exploring a resource dynamics-based feedback mechanism underlying the population growth models. *Ecological Complexity* **6**:4, 403-412. [[Crossref](#)]
153. A. García Vázquez, J.-C. Alonso, F. Carvajal, J. Moreau, J. Nuñez, J.-F. Renno, S. Tello, V. Montreuil, F. Duponchelle. 2009. Life-history characteristics of the large Amazonian migratory catfish *Brachyplatystoma rousseauxii* in the Iquitos region, Peru. *Journal of Fish Biology* **75**:10, 2527-2551. [[Crossref](#)]
154. Marcelo Francisco de Nóbrega, Rosangela Paula Lessa. 2009. Age and growth of the king mackerel (*Scomberomorus Cavalla*) off the northeastern coast of Brazil. *Brazilian Journal of Oceanography* **57**:4, 273-285. [[Crossref](#)]
155. LauretteCombeL. Combe, Abraham J.Escobar-GutiérrezA.J. Escobar-Gutiérrez. 2009. Sénescence d'un pied de maïs : évolution de la floraison à la récolte. *Botany* **87**:11, 1036-1053. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
156. D. Combes, Abraham J. Escobar-Gutiérrez. Morphogenetically Active Radiation within Tree Canopies: Estimations from PAR and Solar Broadband Irradiance Measurements 72-78. [[Crossref](#)]
157. Lisa K.SchwarzL.K. Schwarz, Michael C.RungeM.C. Runge. 2009. Hierarchical Bayesian analysis to incorporate age uncertainty in growth curve analysis and estimates of age from length: Florida manatee (*Trichechus manatus*) carcasses. *Canadian Journal of Fisheries and Aquatic Sciences* **66**:10, 1775-1789. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)] [[Supplemental Material](#)]
158. Abraham J. Escobar-Gutiérrez, Didier Combes, Miroslava Rakocevic, Christophe de Berranger, Annie Eprincharde-Ciesla, Hervé Sinoquet, Claude Varlet-Grancher. 2009. Functional relationships to estimate Morphogenetically Active Radiation (MAR) from PAR and solar broadband irradiance measurements: The case of a sorghum crop. *Agricultural and Forest Meteorology* **149**:8, 1244-1253. [[Crossref](#)]
159. L. Brossard, J.-Y. Dourmad, J. Rivest, J. van Milgen. 2009. Modelling the variation in performance of a population of growing pig as affected by lysine supply and feeding strategy. *animal* **3**:08, 1114. [[Crossref](#)]

160. Nathanael C. Overman, David A. Beauchamp, Hans B. Berge, Michael M. Mazur, Jenifer K. McIntyre. 2009. Differing Forage Fish Assemblages Influence Trophic Structure in Neighboring Urban Lakes. *Transactions of the American Fisheries Society* **138**:4, 741-755. [[Crossref](#)]
161. Rocío Bustos, Ángel Luque, José G. Pajuelo. 2009. Age estimation and growth pattern of the island grouper, *Mycteroperca fusca* (Serranidae) in an island population on the northwest coast of Africa. *Scientia Marina* **73**:2, 319-328. [[Crossref](#)]
162. James T. Thorson, Colin A. Simpfendorfer. 2009. Gear selectivity and sample size effects on growth curve selection in shark age and growth studies. *Fisheries Research* **98**:1-3, 75-84. [[Crossref](#)]
163. Dimitris Charalampopoulos, José Antonio Vázquez, Severino S. Pandiella. 2009. Modelling and validation of *Lactobacillus plantarum* fermentations in cereal-based media with different sugar concentrations and buffering capacities. *Biochemical Engineering Journal* **44**:2-3, 96-105. [[Crossref](#)]
164. Tong-Jiang Xu, Yen-Peng Ting. 2009. Fungal bioleaching of incineration fly ash: Metal extraction and modeling growth kinetics. *Enzyme and Microbial Technology* **44**:5, 323-328. [[Crossref](#)]
165. M. J. Kingsford. 2009. Contrasting patterns of reef utilization and recruitment of coral trout (*Plectropomus leopardus*) and snapper (*Lutjanus carponotatus*) at One Tree Island, southern Great Barrier Reef. *Coral Reefs* **28**:1, 251-264. [[Crossref](#)]
166. S.S. Montgomery, G.W. Liggins, J.R. Craig, J.R. McLeod. 2009. Growth of the spiny lobster *Jasus verreauxi* (Decapoda: Palinuridae) off the east coast of Australia. *New Zealand Journal of Marine and Freshwater Research* **43**:1, 113-123. [[Crossref](#)]
167. Vivian Haist, Paul A. Breen, Paul J. Starr. 2009. A multi-stock, length-based assessment model for New Zealand rock lobster (*Jasus Edwardsii*). *New Zealand Journal of Marine and Freshwater Research* **43**:1, 355-371. [[Crossref](#)]
168. Francisco Cerna, Roberto Licandeo. 2009. Age and growth of the shortfin mako ( *Isurus oxyrinchus* ) in the south-eastern Pacific off Chile. *Marine and Freshwater Research* **60**:5, 394. [[Crossref](#)]
169. André E. Punt, Rik C. Buckworth, Catherine M. Dichmont, Yimin Ye. 2009. Performance of methods for estimating size-transition matrices using tag-recapture data. *Marine and Freshwater Research* **60**:2, 168. [[Crossref](#)]
170. Joseph M. O'Malley. 2009. Spatial and Temporal Variability in Growth of Hawaiian Spiny Lobsters in the Northwestern Hawaiian Islands. *Marine and Coastal Fisheries* **1**:1, 325-342. [[Crossref](#)]
171. Marcelo Francisco de Nóbrega, Rosangela Paula Lessa. 2009. Age and growth of Spanish mackerel (*Scomberomorus brasiliensis*) off the northeastern coast of Brazil. *Neotropical Ichthyology* **7**:4, 667-676. [[Crossref](#)]
172. Ashley J. Williams, Leanne M. Currey, Gavin A. Begg, Cameron D. Murchie, Aaron C. Ballagh. 2008. Population biology of coral trout species in eastern Torres Strait: Implications for fishery management. *Continental Shelf Research* **28**:16, 2129-2142. [[Crossref](#)]

173. Patricia Adame, Miren del Río, Isabel Cañellas. 2008. A mixed nonlinear height–diameter model for pyrenean oak (*Quercus pyrenaica* Willd.). *Forest Ecology and Management* **256**:1–2, 88–98. [[Crossref](#)]
174. Maria Ana de Castelo Branco, Olga Moura, Sofia Gamito. 2008. A growth equation for sole postlarvae (*Solea senegalensis* Kaup, 1858) in earthen ponds in the region of Tagus and Sado Estuaries (Portugal). *Aquaculture Research* **39**:8, 795–800. [[Crossref](#)]
175. Stelios Katsanevakis, Christos D. Maravelias. 2008. Modelling fish growth: multi-model inference as a better alternative to a priori using von Bertalanffy equation. *Fish and Fisheries* **9**:2, 178–187. [[Crossref](#)]
176. Alex De Robertis, Kresimir Williams. 2008. Weight–Length Relationships in Fisheries Studies: The Standard Allometric Model Should Be Applied with Caution. *Transactions of the American Fisheries Society* **137**:3, 707–719. [[Crossref](#)]
177. A. Silva, P. Carrera, J. Massé, A. Uriarte, M.B. Santos, P.B. Oliveira, E. Soares, C. Porteiro, Y. Stratoudakis. 2008. Geographic variability of sardine growth across the northeastern Atlantic and the Mediterranean Sea. *Fisheries Research* **90**:1–3, 56–69. [[Crossref](#)]
178. Jon T. Schnute, Laura J. Richards. Surplus Production Models 105–126. [[Crossref](#)]
179. Lee-Anne Proudfoot, Sven Kaehler, Christopher D. McQuaid. 2008. Using growth band autofluorescence to investigate large-scale variation in growth of the abalone *Haliotis midae*. *Marine Biology* **153**:5, 789–796. [[Crossref](#)]
180. JOHN F. MOXNES, KJELL HAUSKEN. 2008. THE DYNAMIC BIOENERGY OF ANIMALS WITH A DIGESTIVE TRACT. *International Journal of Modern Physics C* **19**:02, 307–349. [[Crossref](#)]
181. . References 320–338. [[Crossref](#)]
182. Rodrigo Wiff, Rubén Roa-Ureta. 2008. Predicting the slope of the allometric scaling of consumption rates in fish using the physiology of growth. *Marine and Freshwater Research* **59**:10, 912. [[Crossref](#)]
183. F. Kozusko, M. Bourdeau. 2007. A unified model of sigmoid tumour growth based on cell proliferation and quiescence. *Cell Proliferation* **40**:6, 824–834. [[Crossref](#)]
184. Lisa J. Natanson, James A. Sulikowski, Jeff R. Kneebone, Paul C. Tsang. 2007. Age and growth estimates for the smooth skate, *Malacoraja senta*, in the Gulf of Maine. *Environmental Biology of Fishes* **80**:2–3, 293–308. [[Crossref](#)]
185. A. J. Williams, B. D. Mapstone, C. R. Davies. 2007. Spatial and interannual patterns in growth of an exploited coral-reef fish. *Journal of Fish Biology* **71**:4, 970–992. [[Crossref](#)]
186. D.E. Benth, B.M. Osei, C.D. Ellingwood, J.P. Hoffmann. 2007. Analysis of a Schnute postulate-based unified growth model for model selection in evolutionary computations. *Biosystems* **90**:2, 467–474. [[Crossref](#)]
187. Mahadev Sharma, John Parton. 2007. Height–diameter equations for boreal tree species in Ontario using a mixed-effects modeling approach. *Forest Ecology and Management* **249**:3, 187–198. [[Crossref](#)]
188. A. B. Dougherty, K. M. Bailey, K. L. Mier. 2007. Interannual differences in growth and hatch date distributions of age-0 year walleye pollock *Theragra chalcogramma* (Pallas) sampled from

- the Shumagin Islands region of the Gulf of Alaska, 1985 ? 2001. *Journal of Fish Biology* **71**:3, 763-780. [[Crossref](#)]
189. LAURA ROGERS-BENNETT, DONALD W. ROGERS, STEVEN A. SCHULTZ. 2007. MODELING GROWTH AND MORTALITY OF RED ABALONE (*HALIOTIS RUFESCENS*) IN NORTHERN CALIFORNIA. *Journal of Shellfish Research* **26**:3, 719-727. [[Crossref](#)]
190. J. Hoey, M. I. McCormick, A. S. Hoey. 2007. Influence of depth on sex-specific energy allocation patterns in a tropical reef fish. *Coral Reefs* **26**:3, 603-613. [[Crossref](#)]
191. Gerard Kuper, Elmer Sterken. Modelling the development of world records in running 7-31. [[Crossref](#)]
192. André Dumas, James France, Dominique P. Bureau. 2007. Evidence of three growth stanzas in rainbow trout (*Oncorhynchus mykiss*) across life stages and adaptation of the thermal-unit growth coefficient. *Aquaculture* **267**:1-4, 139-146. [[Crossref](#)]
193. Seiichi Sakanoue. 2007. Extended logistic model for growth of single-species populations. *Ecological Modelling* **205**:1-2, 159-168. [[Crossref](#)]
194. S. D. Newsome, M. A. Etnier, D. Gifford-Gonzalez, D. L. Phillips, M. van Tuinen, E. A. Hadly, D. P. Costa, D. J. Kennett, T. P. Guilderson, P. L. Koch. 2007. The shifting baseline of northern fur seal ecology in the northeast Pacific Ocean. *Proceedings of the National Academy of Sciences* **104**:23, 9709-9714. [[Crossref](#)]
195. A. Maia, N. Queiroz, H. N. Cabral, A. M. Santos, J. P. Correia. 2007. Reproductive biology and population dynamics of the shortfin mako, *Isurus oxyrinchus Rafinesque*, 1810, off the southwest Portuguese coast, eastern North Atlantic. *Journal of Applied Ichthyology* **23**:3, 246-251. [[Crossref](#)]
196. AKIRA NISHIMURA, TOMONORI HAMATSU, OSAMU SHIDA, IKUO MIHARA, TAKASHI MUTOH. 2007. Interannual variability in hatching period and early growth of juvenile walleye pollock, *Theragra chalcogramma*, in the Pacific coastal area of Hokkaido. *Fisheries Oceanography* **16**:3, 229-239. [[Crossref](#)]
197. Ahmad Mirfazaelian, Jeffrey W. Fisher. 2007. Organ Growth Functions in Maturing Male Sprague-Dawley Rats Based on a Collective Database. *Journal of Toxicology and Environmental Health, Part A* **70**:12, 1052-1063. [[Crossref](#)]
198. J Paige Eveson, Tom Polacheck, Geoff M Laslett. 2007. Consequences of assuming an incorrect error structure in von Bertalanffy growth models: a simulation study. *Canadian Journal of Fisheries and Aquatic Sciences* **64**:4, 602-617. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
199. Ahmad Mirfazaelian, Kyu-Bong Kim, Sookwang Lee, Hyo J. Kim, James V. Bruckner, Jeffrey W. Fisher. 2007. Organ Growth Functions in Maturing Male Sprague-Dawley Rats. *Journal of Toxicology and Environmental Health, Part A* **70**:5, 429-438. [[Crossref](#)]
200. Jason M Cope, André E Punt. 2007. Admitting ageing error when fitting growth curves: an example using the von Bertalanffy growth function with random effects. *Canadian Journal of Fisheries and Aquatic Sciences* **64**:2, 205-218. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
201. Kristina M Wasson, Stephen A Watts, Larry R McEdward, Benjamin G Miner, Thomas A Ebert. **37**, 55. [[Crossref](#)]

202. Thomas A Ebert. Chapter 6 Growth and survival of postsettlement sea urchins 95-134. [[Crossref](#)]
203. Graeme P. Ewing, Jeremy M. Lyle, Raymond J. Murphy, John M. Kalish, Philippe E. Ziegler. 2007. Validation of age and growth in a long-lived temperate reef fish using otolith structure, oxytetracycline and bomb radiocarbon methods. *Marine and Freshwater Research* **58**:10, 944. [[Crossref](#)]
204. S. Dimov, N. Kirilov, S. Peykov, I. Ivanova. 2007. Growth Characteristics of Twenty Lactobacillus Delbrueckii Strains Isolated from Bulgarian Home Made Yoghurts. *Biotechnology & Biotechnological Equipment* **21**:2, 172-176. [[Crossref](#)]
205. Rubén Roa-Ureta, Alexander I. Arkhipkin. 2007. Short-term stock assessment of *Loligo gahi* at the Falkland Islands: sequential use of stochastic biomass projection and stock depletion models. *ICES Journal of Marine Science* **64**:1, 3-17. [[Crossref](#)]
206. F. Köhn, A. R. Sharifi, H. Simianer. 2007. Modeling the growth of the Goettingen minipig1. *Journal of Animal Science* **85**:1, 84-92. [[Crossref](#)]
207. José G. Pajuelo, Iciar Martínez, José A. González, José M. Lorenzo, Antonio García-Mederos, Rosa Domínguez-Seoane, Carmen M. Hernández-Cruz. 2006. Growth pattern and age estimation of the coastal sparid fish *Pagrus auriga* on the Canary Islands shelf. *Fisheries Research* **82**:1-3, 7-13. [[Crossref](#)]
208. Gregor M. Cailliet, Wade D. Smith, Henry F. Mollet, Kenneth J. Goldman. 2006. Age and growth studies of chondrichthyan fishes: the need for consistency in terminology, verification, validation, and growth function fitting. *Environmental Biology of Fishes* **77**:3-4, 211-228. [[Crossref](#)]
209. Miguel Araya, Luis A. Cubillos. 2006. Evidence of two-phase growth in elasmobranchs. *Environmental Biology of Fishes* **77**:3-4, 293-300. [[Crossref](#)]
210. Lisa J. Natanson, Nancy E. Kohler, Daniele Ardizzone, Gregor M. Cailliet, Sabine P. Wintner, Henry F. Mollet. 2006. Validated age and growth estimates for the shortfin mako, *Isurus oxyrinchus*, in the North Atlantic Ocean. *Environmental Biology of Fishes* **77**:3-4, 367-383. [[Crossref](#)]
211. Rosa Domínguez-Seoane, José G. Pajuelo, José M. Lorenzo, Antonio G. Ramos. 2006. Age and growth of the sharpsnout seabream *Diplodus puntazzo* (Cetti, 1777) inhabiting the Canarian archipelago, estimated by reading otoliths and by backcalculation. *Fisheries Research* **81**:2-3, 142-148. [[Crossref](#)]
212. Stephen Haslett, R.B. Millar, K.B. Newman, Martyn Plummer, A.H. Welsh. 2006. INVITED DISCUSSION OF THE PAPER BY DOMIJAN, JORGENSEN & REID. *Australian <html\_ent glyph="&" ascii="&"> New Zealand Journal of Statistics* **48**:3, 393-413. [[Crossref](#)]
213. M. Malosetti, R. G. F. Visser, C. Celis-Gamboa, F. A. Eeuwijk. 2006. QTL methodology for response curves on the basis of non-linear mixed models, with an illustration to senescence in potato. *Theoretical and Applied Genetics* **113**:2, 288-300. [[Crossref](#)]
214. Luis Diaz-Balteiro, Luiz C.E. Rodriguez. 2006. Optimal rotations on Eucalyptus plantations including carbon sequestration—A comparison of results in Brazil and Spain. *Forest Ecology and Management* **229**:1-3, 247-258. [[Crossref](#)]

215. Fernando Castedo Dorado, Ulises Diéguez-Aranda, Marcos Barrio Anta, Marina Sánchez Rodríguez, Klaus von Gadow. 2006. A generalized height–diameter model including random components for radiata pine plantations in northwestern Spain. *Forest Ecology and Management* **229**:1–3, 202–213. [[Crossref](#)]
216. Andre S. Barreto, Fernando C. W. Rosas. 2006. COMPARATIVE GROWTH ANALYSIS OF TWO POPULATIONS OF PONTOPORIA BLAINVILLEI ON THE BRAZILIAN COAST. *Marine Mammal Science* **22**:3, 644–653. [[Crossref](#)]
217. G. Kruitwagen, T. Hecht, H. B. Pratap, S. E. Wendelaar Bonga. 2006. Changes in morphology and growth of the mudskipper (*Periophthalmus argentilineatus*) associated with coastal pollution. *Marine Biology* **149**:2, 201–211. [[Crossref](#)]
218. S. D. H. Bishop, M. P. Francis, C. Duffy, J. C. Montgomery. 2006. Age, growth, maturity, longevity and natural mortality of the shortfin mako shark (*Isurus oxyrinchus*) in New Zealand waters. *Marine and Freshwater Research* **57**:2, 143. [[Crossref](#)]
219. Gudmundur Gudmundsson. 2005. Stochastic growth. *Canadian Journal of Fisheries and Aquatic Sciences* **62**:8, 1746–1755. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
220. John Stewart, Julian M. Hughes. 2005. Longevity, growth, reproduction and a description of the fishery for silver sweep *Scorpius lineolatus* off New South Wales, Australia. *New Zealand Journal of Marine and Freshwater Research* **39**:4, 827–838. [[Crossref](#)]
221. J. X. He, L. G. Rudstam, J. L. Forney, A. J. VanDeValk, D. J. Stewart. 2005. Long-term patterns in growth of Oneida Lake walleye: a multivariate and stage-explicit approach for applying the von Bertalanffy growth function. *Journal of Fish Biology* **66**:5, 1459–1470. [[Crossref](#)]
222. Mariano García-Rodríguez, Pilar Pereda, Jorge Landa, Antonio Esteban. 2005. On the biology and growth of the anglerfish *Lophius budegassa* Spinola, 1807 in the Spanish Mediterranean: a preliminary approach. *Fisheries Research* **71**:2, 197–208. [[Crossref](#)]
223. A. T. KAMUKURU, T. HECHT, Y. D. MGAYA. 2005. Effects of exploitation on age, growth and mortality of the blackspot snapper, *Lutjanus fulvivlamma*, at Mafia Island, Tanzania. *Fisheries Management and Ecology* **12**:1, 45–55. [[Crossref](#)]
224. M. A. H Penna, M. A Villacorta-Corrêa, T. Walter, M. Petrere-JR. 2005. Growth of the tambaqui *Colossoma macropomum* (Cuvier) (Characiformes: Characidae): which is the best model?. *Brazilian Journal of Biology* **65**:1, 129–139. [[Crossref](#)]
225. Gavin A. Begg. Life History Parameters 119–150. [[Crossref](#)]
226. Alexander I. Arkhipkin, Rubén Roa-Ureta. 2005. Identification of ontogenetic growth models for squid. *Marine and Freshwater Research* **56**:4, 371. [[Crossref](#)]
227. Christian Lydersen, Kit M. Kovacs. 2005. Growth and population parameters of the world's northernmost harbour seals *Phoca vitulina* residing in Svalbard, Norway. *Polar Biology* **28**:2, 156–163. [[Crossref](#)]
228. Mahadev Sharma, Shu Yin Zhang. 2004. Height–Diameter Models Using Stand Characteristics for *Pinus banksiana* and *Picea mariana*. *Scandinavian Journal of Forest Research* **19**:5, 442–451. [[Crossref](#)]



229. S. UTHICKE, D. WELCH, J.A.H. BENZIE. 2004. Slow Growth and Lack of Recovery in Overfished Holothurians on the Great Barrier Reef: Evidence from DNA Fingerprints and Repeated Large-Scale Surveys. *Conservation Biology* **18**:5, 1395-1404. [[Crossref](#)]
230. Michael A Etnier. 2004. Reevaluating evidence of density-dependent growth in northern fur seals (*Callorhinus ursinus*) based on measurements of archived skeletal specimens. *Canadian Journal of Fisheries and Aquatic Sciences* **61**:9, 1616-1626. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
231. John D. Gage, Roslyn M. Anderson, Paul A. Tyler, Rachel Chapman, Emily Dolan. 2004. Growth, reproduction and possible recruitment variability in the abyssal brittle star *Ophiocten hastatum* (Ophiuroidea: Echinodermata) in the NE Atlantic. *Deep Sea Research Part I: Oceanographic Research Papers* **51**:6, 849-864. [[Crossref](#)]
232. J.J Colbert, Michael Schuckers, Desta Fekedulegn, James Rentch, Máirtín MacSiúrtáin, Kurt Gottschalk. 2004. Individual tree basal-area growth parameter estimates for four models. *Ecological Modelling* **174**:1-2, 115-126. [[Crossref](#)]
233. G.J. Partridge, G.A. Sarre, N.G. Hall, G.I. Jenkins, J. Chaplin, I.C. Potter. 2004. Comparisons between the growth of *Acanthopagrus butcheri* cultured from broodstock from two estuarine populations that are reproductively isolated and differ markedly in growth rate. *Aquaculture* **231**:1-4, 51-58. [[Crossref](#)]
234. Viktor Chrobok, Milan Meloun, Eva Šimáková. 2004. Descriptive growth model of the height of stapes in the fetus: a histopathological study of the temporal bone. *European Archives of Oto-Rhino-Laryngology* **261**:1, 25-29. [[Crossref](#)]
235. Rosângela Lessa, Francisco M Santana, Fábio H Hazin. 2004. Age and growth of the blue shark *Prionace glauca* (Linnaeus, 1758) off northeastern Brazil. *Fisheries Research* **66**:1, 19-30. [[Crossref](#)]
236. John Stewart, Douglas J. Ferrell, Bryan van der Walt. 2004. Sizes and ages in commercial landings with estimates of growth, mortality and yield per recruit of yellowtail kingfish (*Seriola lalandi*) from New South Wales, Australia. *Marine and Freshwater Research* **55**:5, 489. [[Crossref](#)]
237. Jens Floeter, Axel Temming. 2003. North sea fish and higher trophic levels: a review. *Senckenbergiana maritima* **33**:1-2, 269-442. [[Crossref](#)]
238. Jie Zheng. 2003. Uncertainties of natural mortality estimates for eastern Bering Sea snow crab, *Chionoecetes opilio*. *Fisheries Research* **65**:1-3, 411-425. [[Crossref](#)]
239. TERRANCE J. QUINN. 2003. RUMINATIONS ON THE DEVELOPMENT AND FUTURE OF POPULATION DYNAMICS MODELS IN FISHERIES. *Natural Resource Modeling* **16**:4, 341-392. [[Crossref](#)]
240. JON T. SCHNUTE. 2003. DESIGNING FISHERY MODELS: A PERSONAL ADVENTURE. *Natural Resource Modeling* **16**:4, 393-413. [[Crossref](#)]
241. S. Kayombo, T.S.A. Mbwette, J.H.Y. Katima, S.E Jorgensen. 2003. Effects of substrate concentrations on the growth of heterotrophic bacteria and algae in secondary facultative ponds. *Water Research* **37**:12, 2937-2943. [[Crossref](#)]
242. José G. Pajuelo, José M. Lorenzo, Rosa Domínguez-Seoane. 2003. Age estimation and growth of the zebra seabream *Diplodus cervinus cervinus* (Lowe, 1838) on the Canary Islands shelf (Central-east Atlantic). *Fisheries Research* **62**:1, 97-103. [[Crossref](#)]

243. Ph. Grosjean, Ch. Spirlet, M Jangoux. 2003. A functional growth model with intraspecific competition applied to a sea urchin, *Paracentrotus lividus*. *Canadian Journal of Fisheries and Aquatic Sciences* **60**:3, 237-246. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
244. Liu Zhao-gang, Li Feng-ri. 2003. The generalized Chapman-Richards function and applications to tree and stand growth. *Journal of Forestry Research* **14**:1, 19-26. [[Crossref](#)]
245. Emily B Szalai, Guy W Fleischer, James R Bence. 2003. Modeling time-varying growth using a generalized von Bertalanffy model with application to bloater (*Coregonus hoyi*) growth dynamics in Lake Michigan. *Canadian Journal of Fisheries and Aquatic Sciences* **60**:1, 55-66. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
246. P.V Radebe, B.Q Mann, L.E Beckley, A Govender. 2002. Age and growth of *Rhabdosargus sarba* (Pisces: Sparidae), from KwaZulu-Natal, South Africa. *Fisheries Research* **58**:2, 193-201. [[Crossref](#)]
247. A. Tsoularis, J. Wallace. 2002. Analysis of logistic growth models. *Mathematical Biosciences* **179**:1, 21-55. [[Crossref](#)]
248. Geoff M Laslett, J Paige Eveson, Tom Polacheck. 2002. A flexible maximum likelihood approach for fitting growth curves to tag#recapture data. *Canadian Journal of Fisheries and Aquatic Sciences* **59**:6, 976-986. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
249. Frédéric Lardeux, Jules Cheffort. 2002. Age-grading and growth of *Wuchereria bancrofti* (Filariidea: Onchocercidae) larvae by growth measurements and its use for estimating blood-meal intervals of its Polynesian vector *Aedes polynesiensis* (Diptera: Culicidae). *International Journal for Parasitology* **32**:6, 705-716. [[Crossref](#)]
250. A. Jaworski, E. Kamler. 2002. Development of a bioenergetics model for fish embryos and larvae during the yolk feeding period. *Journal of Fish Biology* **60**:4, 785-809. [[Crossref](#)]
251. Axel Temming, Ulrich Damm. 2002. Life cycle of *Crangon crangon* in the North Sea: a simulation of the timing of recruitment as a function of the seasonal temperature signal. *Fisheries Oceanography* **11**:1, 45-58. [[Crossref](#)]
252. John Stewart, Douglas J. Ferrell. 2001. Age, growth, and commercial landings of yellowtail scad (*Trachurus novaezelandiae*) and blue mackerel (*Scomber australasicus*) off the coast of New South Wales, Australia. *New Zealand Journal of Marine and Freshwater Research* **35**:3, 541-551. [[Crossref](#)]
253. LEI YUANCAI, CARLOS PACHECO MARQUES, JOÃO MANUEL BENTO. 2001. FEATURES AND APPLICATIONS OF BERTALANFFY-RICHARDS' AND SCHNUTE'S GROWTH EQUATIONS. *Natural Resource Modeling* **14**:3, 433-451. [[Crossref](#)]
254. C. Shane Reese, James A Calvin, John C George, Raymond J Tarpley. 2001. Estimation of Fetal Growth and Gestation in Bowhead Whales. *Journal of the American Statistical Association* **96**:455, 915-938. [[Crossref](#)]
255. J P Kritzer, C R Davies, B D Mapstone. 2001. Characterizing fish populations: effects of sample size and population structure on the precision of demographic parameter estimates. *Canadian Journal of Fisheries and Aquatic Sciences* **58**:8, 1557-1568. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]



256. J. Millstein, C.E. O'Clair. 2001. Comparison of age-length and growth-increment general growth models of the Schnute type in the Pacific Blue Mussel, *Mytilus trossulus* Gould. *Journal of Experimental Marine Biology and Ecology* **262**:2, 155-176. [[Crossref](#)]
257. J.R. Chale-Matsau, A. Govender, L.E. Beckley. 2001. Age, growth and retrospective stock assessment of an economically extinct sparid fish, *Polysteganus undulosus*, from South Africa. *Fisheries Research* **51**:1, 87-92. [[Crossref](#)]
258. Robin L. Walker, Ian G. Burns, Jeff Moorby. 2001. Responses of plant growth rate to nitrogen supply: a comparison of relative addition and N interruption treatments. *Journal of Experimental Botany* **52**:355, 309-317. [[Crossref](#)]
259. C. J. Cieszewski. 2001. Three methods of deriving advanced dynamic site equations demonstrated on inland Douglas-fir site curves. *Canadian Journal of Forest Research* **31**:1, 165-173. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
260. Thomas A. Ebert. Growth and survival of post-settlement sea urchins 79-102. [[Crossref](#)]
261. J.F. Van Impe, K. Bernaerts, A.H. Geeraerd, F. Poschet, K.J. Versyck. Modelling and prediction in an uncertain environment 156-179. [[Crossref](#)]
262. J. Fargo, A.R. Kronlund. 2000. Variation in growth for Hecate Strait English sole (*Parophrys vetulus*) with implications for stock assessment. *Journal of Sea Research* **44**:1-2, 3-15. [[Crossref](#)]
263. J. Fargo, T. Wilderbuer. 2000. Population dynamics of rock sole (*Lepidopsetta bilineata*) in the North Pacific. *Journal of Sea Research* **44**:1-2, 123-144. [[Crossref](#)]
264. You-Gan Wang, Christopher J. Jackson. 2000. Growth curves with time-dependent explanatory variables. *Environmetrics* **11**:5, 597-605. [[Crossref](#)]
265. J. van Milgen, N. Quiniou, J. Noblet. 2000. Modelling the relation between energy intake and protein and lipid deposition in growing pigs. *Animal Science* **71**:01, 119-130. [[Crossref](#)]
266. F. Devlieghere, A.H. Geeraerd, K.J. Versyck, H. Bernaert, J.F. Van Impe, J. Debevere. 2000. Shelf life of modified atmosphere packed cooked meat products: addition of Na-lactate as a fourth shelf life determinative factor in a model and product validation. *International Journal of Food Microbiology* **58**:1-2, 93-106. [[Crossref](#)]
267. Adolf Willem Schepers, Jules Thibault, Christophe Lacroix. 2000. Comparison of simple neural networks and nonlinear regression models for descriptive modeling of *Lactobacillus helveticus* growth in pH-controlled batch cultures. *Enzyme and Microbial Technology* **26**:5-6, 431-445. [[Crossref](#)]
268. J.R. Chale-Matsau, A. Govender, L.E. Beckley. 1999. Age and growth of the queen mackerel *Scomberomorus plurilineatus* from KwaZulu-Natal, South Africa. *Fisheries Research* **44**:2, 121-127. [[Crossref](#)]
269. Christopher N. Rooper, Lewis J. Haldorson, Terrance J. Quinn II. 1999. Habitat factors controlling Pacific herring (*Clupea pallasii*) egg loss in Prince William Sound, Alaska. *Canadian Journal of Fisheries and Aquatic Sciences* **56**:6, 1133-1142. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
270. W. M. Mooij, J. M. Van Rooij, S. Wijnhoven. 1999. Analysis and Comparison of Fish Growth from Small Samples of Length-at-Age Data: Detection of Sexual Dimorphism in Eurasian Perch as an Example. *Transactions of the American Fisheries Society* **128**:3, 483-490. [[Crossref](#)]
271. S. D. Berrow, N. Huin, R. Humpidge, A. W. A. Murray, P. A. Prince. 1999. Wing and Primary Growth of the Wandering Albatross. *The Condor* **101**:2, 360-368. [[Crossref](#)]

272. T. Brey. Growth Performance and Mortality in Aquatic Macrobenthic Invertebrates 153-223. [[Crossref](#)]
273. F. Devlieghere, B. Van Belle, J. Debevere. 1999. Shelf life of modified atmosphere packed cooked meat products: a predictive model. *International Journal of Food Microbiology* 46:1, 57-70. [[Crossref](#)]
274. Douglas L. Arnold, Fred Bryce, Doris Miller, Robert Stapley, Stephen Malcolm, Stephen Hayward. 1998. The Toxicological Effects Following the Ingestion of Chinook Salmon from the Great Lakes by Sprague-Dawley Rats during a Two-Generation Feeding-Reproduction Study. *Regulatory Toxicology and Pharmacology* 27:1, S18-S27. [[Crossref](#)]
275. Duncan G. Worthington, Neil L. Andrew. 1997. Does covariation between growth and reproduction compromise the use of an alternative size limit for the blacklip abalone, *Haliotis rubra*, in NSW, Australia?. *Fisheries Research* 32:3, 223-231. [[Crossref](#)]
276. A. J. Booth. 1997. On the life history of the lesser gurnard (Scorpaeniformes: Triglidae) inhabiting the Agulhas Bank, South Africa. *Journal of Fish Biology* 51:6, 1155-1173. [[Crossref](#)]
277. Li Fengri, Wang Yonghe, Hou Lijun. 1997. Comparison of the Chapman-Richards Function with the Schnute model in stand growth. *Journal of Forestry Research* 8:3, 137-143. [[Crossref](#)]
278. Lei Yuancai, C.Pacheco Marques, F.Wolfango Macedo. 1997. Comparison of Schnute's and Bertalanffy-Richards' growth functions. *Forest Ecology and Management* 96:3, 283-288. [[Crossref](#)]
279. B.A. van der Walt, L.E. Beckley. 1997. Age and growth of *Sarpa salpa* (Pisces: Sparidae) off the east coast of South Africa. *Fisheries Research* 31:3, 241-248. [[Crossref](#)]
280. Marc H. Griffiths. 1997. The application of per-recruit models to *Argyrosomus inodorus*, an important South African sciaenid fish. *Fisheries Research* 30:1-2, 103-115. [[Crossref](#)]
281. E. R. Falkenhagen. 1997. Growth modelling of *Pinus roxburghii* in South Africa. *The Southern African Forestry Journal* 178:1, 31-38. [[Crossref](#)]
282. Y Xiao. 1996. How does somatic growth rate affect otolith size in fishes?. *Canadian Journal of Fisheries and Aquatic Sciences* 53:7, 1675-1682. [[Citation](#)] [[PDF](#)] [[PDF Plus](#)]
283. KEVIN M. BAILEY, SUSAN J. PICQUELLE, STELLA M. SPRING. 1996. Mortality of larval walleye pollock *Theragra chalcogramma* in the western Gulf of Alaska, 1988-91. *Fisheries Oceanography* 5:s1, 124-136. [[Crossref](#)]
284. KEVIN M. BAILEY, ANNETTE L. BROWN, MARY M. YOKLAVICH, KATHRYN L. MIER. 1996. Interannual variability in growth of larval and juvenile walleye pollock *Theragra chalcogramma* in the western Gulf of Alaska, 1983-91. *Fisheries Oceanography* 5:s1, 137-147. [[Crossref](#)]
285. Nils T. Hagen. 1996. Tagging sea urchins: a new technique for individual identification. *Aquaculture* 139:3-4, 271-284. [[Crossref](#)]
286. Roland Billard, Jørgen (JOHN) O.T. Jensen. Gamete Removal, Fertilization and Incubation 291-364. [[Crossref](#)]
287. Roland Billard, Jørgen (JOHN) O.T. Jensen, William Pennell, William E. McLean, George K. Iwama. 29, 291. [[Crossref](#)]
288. M. H. Griffiths, T. Hecht. 1995. On the life-history of *Atractoscion aequidens*, a migratory sciaenid off the east coast of southern Africa. *Journal of Fish Biology* 47:6, 962-985. [[Crossref](#)]

289. T. Brey, L.S. Peck, J. Gutt, S. Hain, W.E. Arntz. 1995. Population Dynamics Of *Magellania Fragilis*, a Brachiopod Dominating a Mixed-Bottom Macrobenthic Assemblage on the Antarctic Shelf. *Journal of the Marine Biological Association of the United Kingdom* **75**:04, 857. [[Crossref](#)]
290. G. A. De Leo, M. Gatto. 1995. A size and age-structured model of the European eel (*Anguilla anguilla* L.). *Canadian Journal of Fisheries and Aquatic Sciences* **52**:7, 1351-1367. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
291. J. Zheng, M. C. Murphy, G. H. Kruse. 1995. A length-based population model and stock-recruitment relationships for red king crab, *Paralithodes camtschaticus*, in Bristol Bay, Alaska. *Canadian Journal of Fisheries and Aquatic Sciences* **52**:6, 1229-1246. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
292. R.I.C.C. Francis. 1995. An alternative mark-recapture analogue of Schnute's growth model. *Fisheries Research* **23**:1-2, 95-111. [[Crossref](#)]
293. Christopher Habicht, James E. Seeb, Richard B. Gates, Irvin R. Brock, Carmen A. Olito. 1994. Triploid Coho Salmon Outperform Diploid and Triploid Hybrids Between Coho Salmon and Chinook Salmon During Their First Year. *Canadian Journal of Fisheries and Aquatic Sciences* **51**:S1, 31-37. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
294. M. Maida, J.C. Coll, P.W. Sammarco. 1994. Shedding new light on scleractinian coral recruitment. *Journal of Experimental Marine Biology and Ecology* **180**:2, 189-202. [[Crossref](#)]
295. GUY E. SKINNER, JOHN W. LARKIN, E. JEFFERY RHODEHAMEL. 1994. MATHEMATICAL MODELING OF MICROBIAL GROWTH: A REVIEW. *Journal of Food Safety* **14**:3, 175-217. [[Crossref](#)]
296. Anesh Govender. 1994. Growth of the king mackerel (*Scomberomorus commerson*) off the coast of Natal, South Africa— from length and age data. *Fisheries Research* **20**:1, 63-79. [[Crossref](#)]
297. H. Lundqvist, S. McKinnell, H. Fångstam, I. Berglund. 1994. The effect of time, size and sex on recapture rates and yield after river releases of *Salmo salar* smolts. *Aquaculture* **121**:1-3, 245-257. [[Crossref](#)]
298. Yongshun Xiao. 1994. Growth Models with Corrections for the Retardative Effects of Tagging. *Canadian Journal of Fisheries and Aquatic Sciences* **51**:2, 263-267. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
299. Marc Mangel. 1994. Climate change and salmonid life history variation. *Deep Sea Research Part II: Topical Studies in Oceanography* **41**:1, 75-106. [[Crossref](#)]
300. D. A. Milton, S. J. M. Blaber, N. J. F. Rawlinson. 1993. Age and growth of three species of clupeids from Kiribati, tropical central south Pacific. *Journal of Fish Biology* **43**:1, 89-108. [[Crossref](#)]
301. PETER B. ADAMS, STEPHEN RALSTON, THOMAS E. LAIDIG. 1993. Occurrence of an exceptional catch of pelagic juvenile lingcod (*Ophiodon elongatus*) off Point Reyes, California. *Fisheries Oceanography* **2**:2, 97-100. [[Crossref](#)]
302. T. Wijtes, P.J. McClure, M.H. Zwietering, T.A. Roberts. 1993. Modelling bacterial growth of *Listeria monocytogenes* as a function of water activity, pH and temperature. *International Journal of Food Microbiology* **18**:2, 139-149. [[Crossref](#)]

303. IAN A. McLAREN. 1993. GROWTH IN PINNIPEDS. *Biological Reviews* **68**:1, 1-79. [[Crossref](#)]
304. Alan M. Johnson, Ida S. Umphers. 1993. A nonlinear regression model for describing averaged growth curves of mice. *Communications in Statistics - Theory and Methods* **22**:12, 3405-3418. [[Crossref](#)]
305. Laura J. Richards, Jon T. Schnute, A. R. Kronlund, Richard J. Beamish. 1992. Statistical Models for the Analysis of Ageing Error. *Canadian Journal of Fisheries and Aquatic Sciences* **49**:9, 1801-1815. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
306. E. Garcia-Berthou, R. Moreno-Amich. 1992. Age and growth of an Iberian cyprinodont, *Aphanius iberus* (Cuv. & Val.), in its most northerly population. *Journal of Fish Biology* **40**:6, 929-937. [[Crossref](#)]
307. T. J. Mulligan, B. M. Leaman. 1992. Length-at-Age Analysis: Can You Get What You See?. *Canadian Journal of Fisheries and Aquatic Sciences* **49**:4, 632-643. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
308. Timothy T. Baker, Robert Lafferty, Terrance J. Quinn. 1991. A general growth model for mark-recapture data. *Fisheries Research* **11**:3-4, 257-281. [[Crossref](#)]
309. M.R. Lipinski, M.A. Roeleveld. 1990. Minor extension of the von Bertalanffy growth theory. *Fisheries Research* **9**:4, 367-371. [[Crossref](#)]
310. Alexander J. P. Raat. 1990. Production, consumption and prey availability of northern pike (*Esox lucius*), pikeperch (*Stizostedion lucioperca*) and European catfish (*Silurus glanis*): a bioenergetics approach. *Hydrobiologia* **200-201**:1, 497-509. [[Crossref](#)]
311. David W. Welch, Gordon A. McFarlane. 1990. Quantifying the Growth of Female Pacific Hake (*Merluccius productus*): An Example of Measuring Uncertainty and Bias in Non-linear Parameter Estimation. *Canadian Journal of Fisheries and Aquatic Sciences* **47**:4, 672-681. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
312. J. D. Gage. 1990. Skeletal growth bands in brittle stars: microstructure and significance as age markers. *Journal of the Marine Biological Association of the United Kingdom* **70**:01, 209. [[Crossref](#)]
313. Jon T. Schnute, Laura J. Richards. 1990. A Unified Approach to the Analysis of Fish Growth, Maturity, and Survivorship Data. *Canadian Journal of Fisheries and Aquatic Sciences* **47**:1, 24-40. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
314. Annie Duncan. 1989. Food limitation and body size in the life cycles of planktonic rotifers and cladocerans. *Hydrobiologia* **186-187**:1, 11-28. [[Crossref](#)]
315. Grant G. Thompson. 1989. Analytic solution of a dynamic pool model incorporating constant marginal cost and discount rates. *Mathematical Biosciences* **95**:2, 139-159. [[Crossref](#)]
316. J.G.P.W. Clevers. 1989. Application of a weighted infrared-red vegetation index for estimating leaf Area Index by Correcting for Soil Moisture. *Remote Sensing of Environment* **29**:1, 25-37. [[Crossref](#)]
317. Jon T. Schnute, Laura J. Richards, Alan J. Cass. 1989. Fish Survival and Recruitment: Investigations Based on a Size-Structured Model. *Canadian Journal of Fisheries and Aquatic Sciences* **46**:5, 743-769. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]

318. A. J. P. Raat. 1989. Growth and production of O+ bream (*Abramis brama*), O+ roach (*Rutilus rutilus*) and O+ carp (*Cyprinus carpio*) in 10 drainable 0.1 Ha ponds. *Hydrobiological Bulletin* **23**:1, 67-72. [[Crossref](#)]
319. J. KRIETER, E. KALM. 1989. Growth, feed intake and mature size in Large White and Pietrain pigs. *Journal of Animal Breeding and Genetics* **106**:1-6, 300-311. [[Crossref](#)]
320. Robert L. Burr. 1988. Inferring the Distribution of the Parameters of the von Bertalanffy Growth Model from Length Moments. *Canadian Journal of Fisheries and Aquatic Sciences* **45**:10, 1779-1788. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
321. D.F. Alderdice, J.O.T. Jensen, F.P.J. Velsen. 1988. Incubation of sablefish (*Anoplopoma fimbria*) in a system designed for culture of fragile marine teleost eggs. *Aquaculture* **71**:3, 271-283. [[Crossref](#)]
322. Richard B. Deriso, Ana M. Parma. 1988. Dynamics of Age and Size for a Stochastic Population Model. *Canadian Journal of Fisheries and Aquatic Sciences* **45**:6, 1054-1068. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
323. D.F. Alderdice, J.O.T. Jensen, F.P.J. Velsen. 1988. Preliminary trials on incubation of sablefish eggs (*Anoplopoma fimbria*). *Aquaculture* **69**:3-4, 271-290. [[Crossref](#)]
324. H. Lundqvist, W. C. Clarke, H. Johansson. 1988. The influence of precocious sexual maturation on survival to adulthood of river stocked Baltic salmon, *Salmo salar*, smolts. *Ecography* **11**:1, 60-69. [[Crossref](#)]
325. D.F. Alderdice. 3 Osmotic and Ionic Regulation in Teleost Eggs and Larvae 163-251. [[Crossref](#)]
326. J. D. Gage. 1987. Growth of the deep-sea irregular sea urchins *Echinosigra phiale* and *Hemiaster expergitus* in the Rockall Trough (N.E. Atlantic Ocean). *Marine Biology* **96**:1, 19-30. [[Crossref](#)]
327. J.D. Gage, P.A. Tyler, D. Nichols. 1986. Reproduction and growth of *Echinus acutus* var. *norvegicus* Düben & Koren and *E. elegans* Düben & Koren on the continental slope off Scotland. *Journal of Experimental Marine Biology and Ecology* **101**:1-2, 61-83. [[Crossref](#)]
328. J. D. Gage, P. A. Tyler. 1985. Growth and recruitment of the deep-sea urchin *Echinus affinis*. *Marine Biology* **90**:1, 41-53. [[Crossref](#)]
329. L.C.A. CORSTEN. 1985. CURRENT STATISTICAL ISSUES IN AGRICULTURAL RESEARCH. *Statistica Neerlandica* **39**:2, 159-168. [[Crossref](#)]
330. Jon Schnute, Skip McKinnell. 1984. A Biologically Meaningful Approach to Response Surface Analysis. *Canadian Journal of Fisheries and Aquatic Sciences* **41**:6, 936-953. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
331. D.E. Bentil, O.M. Bonsu, C.D. Ellingwood, J.P. Hoffmann. Modeling uncertainty in population dynamics 274-278. [[Crossref](#)]