



## Pigeonhole Principle

To prove formally that there is no DFA that accepts  $L$  we need:

*The Pigeonhole Principle:* If  $A$  and  $B$  are finite sets and  $|A| > |B|$  then there is no 1-1 function from  $A$  to  $B$ , i.e., if we assign each element of  $A$  (the “pigeons”) to an element of  $B$  (the “pigeonholes”) eventually we must put more than one pigeon in the same hole.

## Proof that $L$ is not Regular

The proof is by contradiction. Suppose  $L$  is regular. Then there is a DFA  $M$  such that  $L = L(M)$ .

- Let  $n$  = number of states in  $M$ .
- Given  $a^\ell b^\ell$  for  $\ell > n$ ,  $M$  must be in some state  $p$  more than once while the  $a$ 's are scanned, by the pigeonhole principle.
- Partition  $a^\ell b^\ell$  into  $x, y$ , and  $z$ , where  $y$  is the string of  $a$ 's scanned between two times state  $p$  is entered. Let  $i = |y|$ .

Observe: We can leave out  $y$  or repeat  $y$  any number of times and end up in the same state. But then for any  $k \geq 0$ ,  $a^{\ell+(k-1)i}b^\ell \in L(M)$ ! E.g.,  $a^{\ell-i}b^\ell \in L(M)$ .

## The Pumping Lemma

**Theorem:** Let  $L$  be a regular language. Then there is a number  $n > 0$  (the “pumping length” of  $L$ ) such that for every string  $w$  in  $L$  such that  $|w| \geq n$ , we can break  $w$  into three strings,  $w = xyz$ , such that:

1.  $|xy| \leq n$
2.  $y \neq \epsilon$
3.  $xy^kz \in L$  for each  $k \geq 0$ .

## Proof of the Pumping Lemma

- Let  $n$  be the number of states in the finite automaton  $M = (Q, \Sigma, \delta, q_0, F)$  which accepts  $L$ . Let  $w = w_1w_2...w_\ell$  be a string of length  $\ell \geq n$ . Let  $r_1...r_{\ell+1}$  be the sequence of states  $M$  enters into while processing  $w$ .
- By the Pigeonhole Principle, two of the states among the first  $\ell + 1$  states are the same. Call the first  $r_s$  and the second  $r_t$ .
- – Let  $x = w_1...w_{s-1}$ ,  $y = w_s...w_{t-1}$ ,  $z = w_t...w_\ell$ .  
– We can easily verify each of the conditions of the lemma.

## Proving a Language $L$ is not Regular

The Pumping Lemma gives a condition that must be satisfied by every regular language. How can we use it to show a language is *not* regular?

*Contrapositive:* For any language  $L$ :

*IF* for every  $n > 0$ , there exists a string  $w \in L$ ,  $|w| \geq n$ , such that for *any* decomposition of  $w$  into  $xyz$  with  $|xy| \leq n$ , there is some  $k \geq 0$  such that  $xy^kz \notin L$

*THEN*  $L$  is *not* regular

**Example:**  $L = \{a^r b^s \mid r \geq s\}$

We are given  $n > 0$ .

We pick  $w = a^n b^n \in L$ .

We are given  $xyz$  with the following properties:

1.  $w = xyz$
2.  $|xy| \leq n$
3.  $y \neq \epsilon$ .

We pick  $k = 0$ .

Now, since  $|xy| \leq n$ , it *must* be the case that  $xy = a^j$  for some  $j \geq 0$ . Since  $y \neq \epsilon$ , it *must* be the case that  $y = a^i$  with  $i > 0$ . So  $xy^k z = xy^0 z = a^{n-i} b^n \notin L$  since there are more  $b$ 's than  $a$ 's. So  $L$  is *not* regular. (Pumping down)

## Using Closure Properties

**Theorem:** The class of languages accepted by finite automata is closed under

1. union;
2. concatenation;
3. star;
4. complementation;
5. intersection.
6. reversal



## Example 2

$L = \{w \in \{a, b\}^* \mid w \text{ has an equal number of } a's \text{ and } b's\}$

If  $L$  is regular then  $L \cap L(a^*b^*)$  is regular, since the regular languages are closed under intersection. But  $L \cap L(a^*b^*) = \{a^n b^n \mid n \geq 0\}$ . which we already showed is not regular, giving a contradiction.

## Or Using the Pumping Lemma

We are given  $n > 0$ .

We pick  $w = a^n b^n \in L$ .

We are given  $xyz$  with the following properties:

1.  $w = xyz$
2.  $|xy| \leq n$
3.  $y \neq \epsilon$ .

We pick  $k = 2$ .

Now, since  $|xy| \leq n$ , it *must* be the case that  $xy = a^j$  for some  $j \geq 0$ . Since  $y \neq \epsilon$ , it *must* be the case that  $y = a^i$  with  $i > 0$ . So  $xy^kz = xy^2z = a^{n+i}b^n \notin L$  since  $n + i \neq n$ . . So  $L$  is *not* regular.

## Example 3

$$L = \{ww \mid w \in \{0,1\}^*\}$$

## Example 3

We are given  $n > 0$ .

We pick  $w = 0^n 1^n 0^n 1^n \in L$ .

We are given  $xyz$  with the following properties:

1.  $w = xyz$
2.  $|xy| \leq n$
3.  $y \neq \epsilon$ .

We pick  $k = 0$ .

Now, since  $|xy| \leq n$ , it *must* be the case that  $xy = 0^j$  for some  $j \geq 0$ . Since  $y \neq \epsilon$ , it *must* be the case that  $y = 0^i$  with  $i > 0$ . So  $xy^k z = xy^0 z = 0^{n-i} 1^n 0^n 1^n \notin L$  since  $n - i \neq n$ . So  $L$  is *not* regular.

## Example 4

$$L = \{010^n1^n \mid n \geq 0\}$$

More than one case for the decomposition  $xyz$ . (What are the possible values of  $xy$  and  $y$ ?)