How to calculate the 95% confidence interval for the slope in a linear regression model in R

Ask Question

Here is an exercise from Introductory Statistics with R:

With the rmr data set, plot metabolic rate versus body weight. Fit a linear regression model to the relation. According to the fitted model, what is the predicted metabolic rate for a body weight of 70 kg? Give a 95% confidence interval for the slope of the line.

rmr data set is in the 'ISwR' package. It looks like this:

> rmr	
body.weigh	nt metabolic.rate
1 49.	9 1079
2 50.	8 1146
3 51.	8 1115
4 52.	6 1161
5 57.	6 1325
6 61.	4 1351
7 62.	3 1402
8 64.	9 1365
9 43.	1 870
10 48.	1 1372
11 52.	2 1132
12 53.	5 1172
13 55.	0 1034
14 55.	0 1155
15 56.	0 1392
16 57.	8 1090
17 59.	0 982
18 59.	
19 59.	2 1342
20 59.	5 1027
21 60.	0 1316
22 62.	1 1574
23 64.	9 1526
24 66.	0 1268
25 66.	4 1205
26 72.	8 1382
27 74.	8 1273
28 77.	1 1439
29 82.	0 1536
30 82.	0 1151

Join Stack Overflow to learn, share knowledge, and build your career.

By using our site, you acknowledge that you have read and understand our Cookie Policy, Privacy Policy, and our Terms of Service.

```
37
         103.0
                         1382
38
         104.5
                         1414
39
         107.7
                         1473
40
         110.2
                         2074
41
         122.0
                         1777
42
         123.1
                         1640
43
         125.2
                          1630
44
         143.3
                         1708
```

I know how to calculate the predicted y at a given x but how can I calculate the confidence interval for the slope?



```
edited Mar 2 '13 at 22:19

NPE
337k 57 721 859

asked Mar 2 '13 at 22:09

Yu Fu
555 1 5 13
```

5 have you tried ?confint ? – Arun Mar 2 '13 at 22:14

1 Answer

Let's fit the model:

```
> library(ISwR)
> fit <- lm(metabolic.rate ~ body.wei&</pre>
> summary(fit)
Call:
lm(formula = metabolic.rate ~ body.wei
Residuals:
    Min
             1Q Median
-245.74 -113.99 -32.05 104.96 484.8
Coefficients:
            Estimate Std. Error t valu
(Intercept) 811.2267 76.9755 10.53
body.weight 7.0595
                         0.9776 7.22
Signif. codes: 0 '***, 0.001 '**, 0.6
Residual standard error: 157.9 on 42 \ensuremath{\text{c}}
Multiple R-squared: 0.5539, Adjusted F
F-statistic: 52.15 on 1 and 42 DF, p-
```

to learn, share knowledge, and build your career.

By using our site, you acknowledge that you have read and understand our , , and our

This can be computed using

confint :

edited Mar 3 '13 at 8:40

answered Mar 2 '13 at 22:15



NPE 337k

57 721 859

13 This is equivalent to the following:
 coef=summary(fit)\$coefficients[2,1]
 err=summary(fit)\$coefficients[2,2]
 coef + c(-1,1)*err*qt(0.975,
 42) [1] 5.086656 9.032400 : it's the
 estimated coefficient +- qt(1-alpha/2,
 df) standard errors - ds440 Mar 2 '13
 at 22:59 *

- 1 Thank you, NPE! So estimated coefficient +/- two standard errors is an approximation and the latter method provides a accurate way to calculate the confidence interval, right? Yu Fu Mar 2 '13 at 23:09
- Yes, the two SE is a good ballpark: if the linear model assumptions are correct then it will follow a T distribution so as sample size increases it approaches ~1.96, for smaller samples it is higher. – ds440 Mar 2 '13 at 23:21
- 1 @NPE: Are you assuming that a Gaussian pdf for the slope values? If this hypothesis does not hold you could use bootstrap methods. – ipcgandre Sep 12 '14 at 11:53

to learn, share knowledge, and build your career.

By using our site, you acknowledge that you have read and understand our

, and our