

# PH 235 SPRING 2017

## ASSIGNMENT 6

**Assigned Date:** March 16, 2017

**Due Date:** 11:50pm, March 29, 2017 only via moodle (See syllabus for late policy)

**Total Points:** 100


All problems in this assignment are from the optional textbook for the course *Computational Physics* authored by Mark Newman.

### Instructions:

1. Create a folder titled YourLastName-HW6.
2. In this parent folder, create subfolders, one for each of the problems below and name each subfolder as YourLastName-ExerciseNumber for easier tracking.
3. Each file you submit should be of the format YourLastName-ExerciseNumber-Filename 4. Populate the folders with (a) the solution code for each problem, (b) screen shots of results when possible and (c) a text document with a short description of your approach, whether you successfully solved the problem. If you did not solve the problem, explain solution status (partial credit will be given for reasonable attempts).
5. Submit a zipped version of the parent folder to the moodle assignment link.

### Problem 1: Ex 8.10 Cometary Orbits (40 points)

Many comets travel in highly elongated orbits around the Sun. For much of their lives they are far out in the solar system, moving very slowly, but on rare occasions their orbit brings them close to the Sun for a fly-by and for a brief period of time they move very fast indeed:



This is a classic example of a system for which an adaptive step size method is useful, because for the large periods of time when the comet is moving slowly we can use long time-steps, so that the program runs quickly, but short time-steps are crucial in the brief but fast-moving period close to the Sun.

The differential equation obeyed by a comet is straightforward to derive. The force between the Sun, with mass  $M$  at the origin, and a comet of mass  $m$  with position vector  $\mathbf{r}$  is  $GMm/r^2$  in direction  $-\mathbf{r}/r$  (i.e., the direction towards the Sun), and hence Newton's second law tells us that

$$m \frac{d^2 \mathbf{r}}{dt^2} = - \left( \frac{GMm}{r^2} \right) \frac{\mathbf{r}}{r}.$$

Canceling the  $m$  and taking the  $x$  component we have

$$\frac{d^2 x}{dt^2} = -GM \frac{x}{r^3},$$

and similarly for the other two coordinates. We can, however, throw out one of the coordinates because the comet stays in a single plane as it orbits. If we orient our axes so that this plane is perpendicular to the  $z$ -axis, we can forget about the  $z$  coordinate and we are left with just two second-order equations to solve:

$$\frac{d^2 x}{dt^2} = -GM \frac{x}{r^3}, \quad \frac{d^2 y}{dt^2} = -GM \frac{y}{r^3},$$

where  $r = \sqrt{x^2 + y^2}$ .

1. Turn these two second-order equations into four first-order equations, using the methods you have learned.
2. Write a program to solve your equations using the fourth-order Runge–Kutta method with a *fixed* step size. You will need to look up the mass of the Sun and Newton's gravitational constant  $G$ . As an initial condition, take a comet at coordinates  $x = 4$  billion kilometers and  $y = 0$  (which is somewhere out around the orbit of Neptune) with initial velocity  $v_x = 0$  and  $v_y = 500 \text{ m s}^{-1}$ . Make a graph showing the trajectory of the comet (i.e., a plot of  $y$  against  $x$ ).


Choose a fixed step size  $h$  that allows you to accurately calculate at least two full orbits of the comet. Since orbits are periodic, a good indicator of an accurate calculation is that successive orbits of the comet lie on top of one another on your plot. If they do not then you need a smaller value of  $h$ . Give a short description of your findings. What value of  $h$  did you use? What did you observe in your simulation? How long did the calculation take?

3. Make a copy of your program and modify the copy to do the calculation using an adaptive step size. Set a target accuracy of  $\delta = 1$  kilometer per year in the position of the comet and again plot the trajectory. What do you see? How do the speed, accuracy, and step size of the calculation compare with those in part (b)?
4. Modify your program to place dots on your graph showing the position of the comet at each Runge–Kutta step around a single orbit. You should see the steps getting closer together when the comet is close to the Sun and further apart when it is far out in the solar system.

Calculations like this can be extended to cases where we have more than one orbiting body. We can include planets, moons, asteroids, and others. Analytic calculations are impossible for such complex systems, but with careful numerical solution of differential equations we can calculate the motions of objects throughout the entire solar system.

### Problem 2: Ex 8.15 The Lagrange point (30 points)

The nonlinear pendulum's movement discussed in Example 8.6 is non-linear and yet perfectly regular and periodic—there are no surprises. A *double pendulum*, on the other hand, is completely the opposite—chaotic and unpredictable. A double pendulum consists of a normal pendulum with another pendulum hanging from its end. For simplicity let us ignore friction, and assume that both pendulums have bobs of the same mass  $m$  and massless arms of the same length  $\ell$ . Thus the setup looks like this:



The position of the arms at any moment in time is uniquely specified by the two angles  $\theta_1$  and  $\theta_2$ . The equations of motion for the angles are most easily derived using the Lagrangian formalism, as follows.

The heights of the two bobs, measured from the level of the pivot are

$$h_1 = -\ell \cos \theta_1, \quad h_2 = -\ell(\cos \theta_1 + \cos \theta_2),$$

so the potential energy of the system is

$$V = mgh_1 + mgh_2 = -mg\ell(2 \cos \theta_1 + \cos \theta_2),$$

where  $g$  is the acceleration due to gravity. The (linear) velocities of the two bobs are given by

$$v_1 = \ell \dot{\theta}_1, \quad v_2^2 = \ell^2 [\dot{\theta}_1^2 + \dot{\theta}_2^2 + 2\dot{\theta}_1 \dot{\theta}_2 \cos(\theta_1 - \theta_2)],$$

where  $\dot{\theta}$  means the derivative of  $\theta$  with respect to time  $t$ . (If you don't see where the second velocity equation comes from, it's a good exercise to derive it for yourself from the geometry of the pendulum.) Now the total kinetic energy is

$$T = \frac{1}{2}mv_1^2 + \frac{1}{2}mv_2^2 = m\ell^2 [\dot{\theta}_1^2 + \frac{1}{2}\dot{\theta}_2^2 + \dot{\theta}_1 \dot{\theta}_2 \cos(\theta_1 - \theta_2)],$$

and the Lagrangian of the system is

$$\mathcal{L} = T - V = m\ell^2 [\dot{\theta}_1^2 + \frac{1}{2}\dot{\theta}_2^2 + \dot{\theta}_1 \dot{\theta}_2 \cos(\theta_1 - \theta_2)] + mg\ell(2 \cos \theta_1 + \cos \theta_2).$$

Then the equations of motion are given by the Euler–Lagrange equations

$$\frac{d}{dt} \left( \frac{\partial \mathcal{L}}{\partial \dot{\theta}_1} \right) = \frac{\partial \mathcal{L}}{\partial \theta_1}, \quad \frac{d}{dt} \left( \frac{\partial \mathcal{L}}{\partial \dot{\theta}_2} \right) = \frac{\partial \mathcal{L}}{\partial \theta_2},$$

which in this case give

$$\begin{aligned} 2\ddot{\theta}_1 + \ddot{\theta}_2 \cos(\theta_1 - \theta_2) + \dot{\theta}_2^2 \sin(\theta_1 - \theta_2) + 2\frac{g}{\ell} \sin \theta_1 &= 0, \\ \ddot{\theta}_2 + \ddot{\theta}_1 \cos(\theta_1 - \theta_2) - \dot{\theta}_1^2 \sin(\theta_1 - \theta_2) + \frac{g}{\ell} \sin \theta_2 &= 0, \end{aligned}$$

where the mass  $m$  has canceled out.

These are second-order equations, but we can convert them into first-order ones by the usual method, defining two new variables,  $\omega_1$  and  $\omega_2$ , thus:

$$\dot{\theta}_1 = \omega_1, \quad \dot{\theta}_2 = \omega_2.$$

In terms of these variables our equations of motion become

$$\begin{aligned} 2\dot{\omega}_1 + \dot{\omega}_2 \cos(\theta_1 - \theta_2) + \omega_2^2 \sin(\theta_1 - \theta_2) + 2\frac{g}{\ell} \sin \theta_1 &= 0, \\ \dot{\omega}_2 + \dot{\omega}_1 \cos(\theta_1 - \theta_2) - \omega_1^2 \sin(\theta_1 - \theta_2) + \frac{g}{\ell} \sin \theta_2 &= 0. \end{aligned}$$

Finally we have to rearrange these into the standard form of Eq. (8.29) with a single derivative on the left-hand side of each one, which gives

$$\begin{aligned} \dot{\omega}_1 &= -\frac{\omega_1^2 \sin(2\theta_1 - 2\theta_2) + 2\omega_2^2 \sin(\theta_1 - \theta_2) + (g/\ell) [\sin(\theta_1 - 2\theta_2) + 3 \sin \theta_1]}{3 - \cos(2\theta_1 - 2\theta_2)}, \\ \dot{\omega}_2 &= \frac{4\omega_1^2 \sin(\theta_1 - \theta_2) + \omega_2^2 \sin(2\theta_1 - 2\theta_2) + 2(g/\ell) [\sin(2\theta_1 - \theta_2) - \sin \theta_2]}{3 - \cos(2\theta_1 - 2\theta_2)}. \end{aligned}$$

(This last step is quite tricky and involves some trigonometric identities. If you're not certain of how the calculation goes you may find it useful to go through the derivation for yourself.)

These two equations, along with the equations  $\dot{\theta}_1 = \omega_1$  and  $\dot{\theta}_2 = \omega_2$ , give us four first-order equations which between them define the motion of the double pendulum.

1. Derive an expression for the total energy  $E = T + V$  of the system in terms of the variables  $\theta_1$ ,  $\theta_2$ ,  $\omega_1$ , and  $\omega_2$ , plus the constants  $g$ ,  $\ell$ , and  $m$ .
2. Write a program using the fourth-order Runge–Kutta method to solve the equations of motion for the case where  $\ell = 40$  cm, with the initial conditions  $\theta_1 = \theta_2 = 90^\circ$  and  $\omega_1 = \omega_2 = 0$ . Use your program to calculate the total energy of the system assuming that the mass of the bobs is 1 kg each, and make a graph of energy as a function of time from  $t = 0$  to  $t = 100$  seconds.

Because of energy conservation, the total energy should be constant over time (actually it should be zero for this particular set of initial conditions), but you will find that it is not perfectly constant because of the approximate nature of the solution of the differential equation. Choose a suitable value of the step size  $h$  to ensure that the variation in energy is less than  $10^{-5}$  Joules over the course of the calculation.

3. Make a copy of your program and modify the copy to create a second program that does not produce a graph, but instead makes an animation of the motion of the double pendulum over time. At a minimum, the animation should show the two arms and the two bobs.

Hint: You will probably find the function rate useful in order to make your program run at a steady speed. You will probably also find that the value of  $h$  needed to get the required accuracy in your solution gives a frame-rate much faster than any that can reasonably be displayed in your animation, so you won't be able to display every time-step of the calculation in the animation. Instead you will have to arrange the program so that it updates the animation only once every several Runge–Kutta steps.

### **Problem 3: Ex 7.4 Fourier Filtering and Smoothing (20 points)**

In the on-line resources you'll find a file called `dow.txt`. It contains the daily closing value for each business day from late 2006 until the end of 2010 of the Dow Jones Industrial Average, which is a measure of average prices on the US stock market.

Write a program to do the following:

1. Read in the data from `dow.txt` and plot them on a graph.
2. Calculate the coefficients of the discrete Fourier transform of the data using the function `rfft` from `numpy.fft`, which produces an array of  $\frac{1}{2}N + 1$  complex numbers.
3. Now set all but the first 10% of the elements of this array to zero (i.e., set the last 90% to zero but keep the values of the first 10%).
4. Calculate the inverse Fourier transform of the resulting array, zeros and all, using the function `irfft`, and plot it on the same graph as the original data. You may need to vary the colors of the two curves to make sure they both show up on the graph. Comment on what you see. What is happening when you set the Fourier coefficients to zero?
5. Modify your program so that it sets all but the first 2% of the coefficients to zero and run it again.