

①

Sec 10. Scattering Amplitudes & Feynman Rules.

With $Z(J) = e^{iW(J)}$

we can calculate time-ordered amplitudes through functional derivatives.

$$\frac{1}{i} \frac{\delta}{\delta J} i \Delta(x_1 - x_2) \equiv \langle 0 | T(\varphi(x_1) \varphi(x_2)) | 0 \rangle, \text{ with } \delta_j = \frac{1}{i} \frac{\delta}{\delta J(x_j)}$$

$$= \delta_1 \delta_2 Z(J)|_{J=0} = \delta_1 \delta_2 i W(J)|_{J=0} - \delta_1 i W(J)|_{J=0} \delta_2 i W(J)|_{J=0}$$

$$= \delta_1 \delta_2 i W(J)|_{J=0}$$

$$\text{as } \delta_j W(J)|_{J=0} = \langle 0 | (\varphi(x_j)) | 0 \rangle = 0$$

δ_i removes a source and gives a label x_i

$\frac{1}{i} \Delta(x_1 - x_2)$ is given by all diagrams with two sources.

at lowest order



Thus we get

$$\frac{1}{i} \Delta(x_1 - x_2) = \frac{1}{i} \Delta(x_1 - x_2) + O(g^2)$$

Feynman Propagator

$$\Delta(x - x') = \frac{i k^4 \delta(x - x')}{(2\pi)^4 k^2 + m^2 - i\epsilon}$$

$$\langle 0 | T(\varphi(x_1) \varphi(x_2) \varphi(x_3) \varphi(x_4)) | 0 \rangle = \delta_1 \delta_2 \delta_3 \delta_4 Z(J)$$

$$= [\delta_1 \delta_2 \delta_3 \delta_4 i W(J) + (\delta_1 \delta_2 i W(J))(\delta_3 \delta_4 i W(J))$$

$$+ (\delta_1 \delta_3 i W(J))(\delta_2 \delta_4 i W(J))$$

$$+ (\delta_1 \delta_4 i W(J))(\delta_2 \delta_3 i W(J))]|_{J=0}$$

$$= \delta_1 \delta_2 \delta_3 \delta_4 i W(J)|_{J=0} + \frac{1}{i} \Delta(x_1 - x_2) \Delta(x_3 - x_4) + \dots$$

JED

(2)

We can insert these terms to the LSZ relation
for $2 \rightarrow 2$ process

$$\langle f | i \rangle = i^4 \int d^4 x_1 d^4 x_2 d^4 x_3 d^4 x_4 e^{i(k_1 x_1 + k_2 x_2 - k'_1 x'_1 - k'_2 x'_2)}$$

$$(-\partial_1^2 + m^2)(-\partial_2^2 + m^2)(-\partial_3^2 + m^2)(-\partial_4^2 + m^2)$$

$$\langle 0 | T(\varphi(x_1) \varphi(x_2) \varphi(x'_1) \varphi(x'_2)) | 0 \rangle.$$

There are several products of Green's functions contributing,

let's check $\frac{1}{i} \Delta(x_1 - x'_1) \frac{1}{i} \Delta(x_2 - x'_2)$

$$\int d^4 x_1 d^4 x_2 d^4 x'_1 d^4 x'_2 e^{i(k_1 x_1 + k_2 x_2 - k'_1 x'_1 - k'_2 x'_2)} F(x_{11}) F(x_{22})$$

where we defined $F(x_{ij}) = (-\partial_i^2 + m^2)(-\partial_j^2 + m^2) \Delta(x_{ij})$.

$$x_{ij} = x_i - x_j, \bar{k}_{ij} = (k_i + k_j)/2.$$

$$= (2\pi)^4 \delta^4(k_1 - k'_1) (2\pi)^4 \delta^4(k_2 - k'_2) F(\bar{k}_{11}) F(\bar{k}_{22})$$

$F(k)$ is ~~not~~ $F(x_{ij})$'s Fourier Transform. check

but clearly this represents no scattering as
 $k_1 = k'_1, k_2 = k'_2$

In general, we are only interested in diagrams where something happened (scattering),

which is "sufficiently" represented by

Fully Connected diagrams.

what if some connected,
so not? They can be
factorized.

(3)


From here on, we restrict ourselves to those diagrams alone and define Connected correlation functions via.

$$\langle 0 | T \varphi(x_1) \dots \varphi(x_E) | 0 \rangle_C = \delta_1 \dots \delta_E i W(J) |_{J=0}.$$

Back to $2 \rightarrow 2$ process.

$$\langle 0 | T (\varphi(x_1) \varphi(x_2) \varphi(x'_1) \varphi(x'_2)) | 0 \rangle_C = \delta_1 \delta_2 \delta_1 \delta_2 i W(J) |_{J=0}$$

at the lowest order in g , the contributing process is




with the 4 (comes)

gluons/ gluon away by δ .

There are $4!$ ways to $\delta_1 \delta_2 \delta_3 \delta_4$

which are 3 groups (A2B pairing, C&D pairing, and each with 8 identical exchange of the 4 vertices) reflection diagrams.



Note ~~here~~ that the factor 8 nicely cancels the symmetry factor $S=8$ of the original source diagram.

This is a general result for TREE-level diagrams (those without loops), that the symmetry factor get cancelled with explicit labelling. This is the first non-zero contribution to the amplitude.

(4)

Now. \rightarrow each propagator is $\frac{1}{i}$ as $\frac{1}{i} \frac{\delta}{\delta j} : \Delta \cdot \frac{1}{i} \frac{\delta}{\delta j}$.

$$\begin{aligned} & \langle 0 | T(\phi(x_1)\phi(x_2)\phi(x'_1)\phi(x'_2)) | 0 \rangle_C \\ &= (ig)^2 \left(\frac{1}{i} \right)^5 \int d^4y d^4z \Delta(y-z) \times \\ & [\Delta(x_1-y) \Delta(x_2-y) \Delta(x'_1-z) \Delta(x'_2-z) \\ & + \Delta(x_1-y) \Delta(x_2-z) \Delta(x'_1-z) \Delta(x'_2-y) \\ & + \Delta(x_1-z) \Delta(x_2-y) \Delta(x'_1-z) \Delta(x'_2-y)] + O(g^4). \end{aligned}$$

Put in LSZ. Each Klein-Gordon wave operator acts on a propagator to give.

$$(-\partial_i^2 + m^2) \Delta(x_i - y) = \delta^4(x_i - y) \leftarrow \text{free-particle propagator,}$$

(note that Δ is as it is from Z(i)).

Then we get.

$$\begin{aligned} \langle f | i \rangle &= (ig)^2 \frac{1}{i} \int d^4y d^4z \Delta(y-z) \\ & [e^{-i(k_1 y + k_2 z - k'_1 z - k'_2 y)} \\ & + e^{-i(k_1 y + k_2 z - k'_1 z - k'_2 y)} \\ & + e^{-i(k_1 z + k_2 y - k'_1 z - k'_2 y)}] + O(g^4). \end{aligned}$$

knowing that $\Delta(y-z) = \int \frac{d^4k}{(2\pi)^4} \frac{e^{-ik(y-z)}}{k^2 + m^2 - ie}$.

we can get

(5)

$$\langle f | i \rangle = -ig^2 \int d^4 k \quad e$$

$$\langle f | i \rangle = -ig^2 \int \frac{d^4 k}{(2\pi)^4} \frac{1}{k^2 + m^2 - ie}$$

$$x [(2\pi)^4 \delta^4(k_1 + k_2 + k) + (2\pi)^4 \delta^4(k'_1 + k'_2 + k)]$$

$$+ (2\pi)^4 \delta^4(k_1 - k'_1 + k) (2\pi)^4 \delta^4(k_2 - k'_2 + k)$$

$$+ (2\pi)^4 \delta^4(k_1 - k'_2 + k) (2\pi)^4 \delta^4(k_2 - k'_1 + k)] + O(g^4).$$

$$= ig^2 \cdot (2\pi)^4 \cdot \delta^4(k_1 + k_2 - k'_1 - k'_2)$$

$$\left[\frac{1}{(k_1 + k_2)^2 + m^2 - ie} + \frac{1}{(k_1 - k'_1)^2 + m^2 - ie} + \frac{1}{(k_1 - k'_2)^2 + m^2 - ie} \right] + O(g^4).$$

The overall Dirac delta shows the energy momentum conservation. So

$$\langle f | i \rangle = (2\pi)^4 \delta^4(k_{in} - k_{out}) \underbrace{-ie}_{\text{The scattering Matrix Element}}$$

Bring QFT to anomalies. Feynman Rules

Now we can finally arrive at the Feynman Rules that we learned from the lecture process for calculating $\langle f | i \rangle$.

The Feynman Rules are:

(6)

① Draw lines (external lines) for each incoming or outgoing particles

② leave one end of each external line free and connect the other end to a vertex

where exactly three lines meet.

for ϕ^3 theory.

Include extra internal internal lines to do this if necessary.

Draw all diagrams that are Topologically Inequivalent.

③ On each incoming line, draw arrow towards the vertex

On each outgoing line, draw arrow away from the vertex

On each internal line, draw it arbitrarily.

④ Assign each line its own momentum. The four momentum of external lines are their own momentum.

⑤ Thinking of the four-momenta as following along the arrows, and conserve four-momenta at each vertex.

(For a tree diagram, this fix all ~~momenta~~ momenta)

⑥ The value of each diagram contains the following factors
each external line has, 1;

each internal line with momentum k , $\frac{-i}{k^2 + m^2 - i\epsilon}$

each vertex, $i \gamma_5 g$.

(7)

7 (loop). A diagram with 1 ~~top~~^{internal} closed loops will have 1 internal momenta surfaced by rule #5.

Integrate over these 4-momenta with measure

$$\frac{d^4 p_i}{(2\pi)^4}$$

8. ~~1~~ (loop) A loop diagram may have left over symmetry factors if there are exchange of internal

propagators and vertices that leave the diagram unchanged.

Divide the value by such ~~sym~~ leftover symmetry factors.

9. (counter terms) Include diagrams with the

counterterm vertex that connects two propagators,

each with the same four momentum k . The value of

this vertex is $-i(Ak^2 + Bm^2)$, where


$A = z_\varphi - 1$ and $B = z_m - 1$, each are expected

to be $\mathcal{O}(g^2)$

10. The value of Π is given by a sum over the values

of all these diagrams.

(8)



At tree level, we have.

$$iT = (igg)^2 \left(+ \frac{-i}{(k_1+k_2)^2+m^2-i\epsilon} + \frac{-i}{(k_1-k_1')^2+m^2-i\epsilon} \right. \\ \left. + \frac{-i}{(k_1-k_2)^2+m^2-i\epsilon} \right) + O(g^4)$$

Now we can proceed to get observable cross section from the scattering amplitudes from iT .