

CHAPTER 2: BRIEF QFT

2.1 BASIC GROUP THEORY

Here is a quick refresher on the group theory, which one could have learned from the mathematical physics class and the QFT class. The logic is the same: I hope you have a better understanding of where the “rules” in particle physics come from.

Group theory is used in many places in particle physics, in understanding global symmetries, (local) gauge symmetries, particle representations, etc. Here, we recap the basics.

A Group G is defined by a set of elements $g_i \in G$ and its multiplication rule, $g_1 \cdot g_2$, which satisfy the following,

- **Closure:** if $g_1 \in G$ and $g_2 \in G$, then $g_1 \cdot g_2 \in G$;
- **Associativity:** if $g_1, g_2, g_3 \in G$, then $g_1 \cdot (g_2 \cdot g_3) = (g_1 \cdot g_2) \cdot g_3$ (note the order is unchanged);
- **Identity:** there exists $I \in G$ such that $I \cdot g_i = g_i \cdot I = g_i$;
- **Inverse:** if $g_1 \in G$, then there exists $g_1^{-1} \in G$ such that $g_1^{-1} \cdot g_1 = g_1 \cdot g_1^{-1} = I$.

Here are a few simple examples:

- $\mathbb{Z}_2 = \{-1, 1\}$, with the multiplication rule defined as multiplication.
- $\{\mathbb{Z}\}$, the set of integers, with the multiplication rule defined as addition. One can see that if $n, m \in \mathbb{Z}$, $m \cdot n = m + n \in \mathbb{Z}$ and the identity is 0 and the inverse of each element m is $-m$.

Q & As

(Chimeln) Can $\{x \in \mathbb{R} | x \neq 0\}$, the set of all real numbers with 0 removed, be a group?

(Chimeln) What is the identity?

(Chimeln) What is the multiplication?

(Chimeln) Can $\{\mathbb{R}\}$, the set of all real numbers, be a group?

The groups we will use in this class can be categorized differently. For instance, the groups can be **discrete** or **continuous**, which describes the group member’s continuity behavior. Importantly, we also deal with the following types of groups,

- **Abelian group:** for all $g_1, g_2 \in G$, $g_1 \cdot g_2 = g_2 \cdot g_1$;
- **Non-abelian group:** there exists $g_1, g_2 \in G$, $g_1 \cdot g_2 \neq g_2 \cdot g_1$.

In particle physics, we often use groups (with n degrees) made up of elements of n -by- n matrices. Particularly,

- **Unitary Group**: with all group elements $g_i^\dagger = g_i$; e.g., U(1), U(3)
- **Special Unitary Group**: additionally $\det[g_i] = 1$; e.g., SU(2), SU(3), SU(5)
- **Orthogonal Group** group: with all group elements $g_i^T = g_i$; e.g., O(3)
- **Special Orthogonal Group**: additionally $\det[g_i] = 1$ e.g., SO(3)

Another critically important concept is representation. **A representation¹** of group G is a mapping, D of elements of G onto a set of linear operators with the following properties

- $D(I) = 1$, where 1 is the identity operator in the space on which the linear operators act.
- $D(g_1)D(g_2) = D(g_1 \cdot g_2)$, in other words, the group multiplication law is mapped onto the natural multiplication in the linear space on which the linear operators act.

One can view the group multiplication rule as a mapping table where group elements (column) group-multiplied by group elements (row) gives another group element. For a group of n elements (this is to say the group G is of **order** n), this forms an $n \times n$ generalized matrix D whose matrix elements are group members g_i . Now define $D(g_i)$ as a $n \times n$ matrix where group element g_i in D is replaced by 1, and all others are set to zero. Then, we get a set (total numbers is n) of $n \times n$ matrices, which automatically satisfies the group multiplication rules with its natural matrix multiplication. This is called the **regular representation** of a group, which can be **reducible**. The regular representation $D(g_i)$ satisfy the following, treating the group elements g_i to form an orthonormal basis for vector space $|g_i\rangle$:

$$D(g_i) |g_j\rangle = |g_i \cdot g_j\rangle. \quad (2.1.1)$$

Clearly, one can obtain the **matrix elements** of representation for arbitrary group element g , $D(g)$ via,

$$D(g)_{ij} = \langle g_i | D(g) | g_j \rangle \quad (2.1.2)$$

One can see such a n-by-n matrix representation $D(g)$ constructed this way satisfies the group multiplication rule by the linear matrix natural multiplication,

$$[D(g_a)D(g_b)]_{ik} = \sum_j D(g_a)_{ij} D(g_b)_{jk} = \sum_j \langle g_i | D(g_a) | g_j \rangle \langle g_j | D(g_b) | g_k \rangle \quad (2.1.3)$$

$$\begin{aligned} &= \langle g_i | D(g_a) D(g_b) | g_k \rangle = \langle g_i | D(g_a) | g_b \cdot g_k \rangle = \langle g_i | g_a \cdot g_b \cdot g_k \rangle \\ &= \langle g_i | (g_a \cdot g_b) \cdot g_k \rangle = \langle g_i | D(g_a \cdot g_b) | g_k \rangle = D(g_a \cdot g_b)_{ik}. \end{aligned} \quad (2.1.4)$$

¹Here I directly quote the wording in Ref. [Geo99].

In the second line, I used the fact that $\sum_k |g_k\rangle \langle g_k|$ is complete and in the third line the associativity of group G . Unitary representations ($D(g)^\dagger = D(g)$) are particularly important, and all representations of finite groups are equivalent to unitary representations.

This regular representation can form a new **equivalent representation**, by a similarity transformation S ,

$$D'(g) = S^{-1}D(g)S, \quad (2.1.5)$$

and this new representation satisfies all group transformation rules.

An important concept in representation is its reducibility. A representation is reducible if it has an **invariant subspace**, that the multiplication of $D(g)$ on any of the elements in the subspace remains in the subspace. This will allow us to decompose a representation into a “sum” of irreducible representations. As the representations of the SM elementary fermions are found to live in irreducible representations called fundamental representations, and the gauge boson lives in the adjoint representations. We do not delve much further here into the representation theory. But if you work on model building, I highly recommend you to read more of Ref. [Geo99].

Q & As

Q: There is a jump between the group multiplication rule, $g_1 \cdot g_2$, to the claim that I can represent the group by matrices following natural multiplication in the linear space. Do representations exist for all groups?

A: Yes. See the discussion about the construction of a regular representation.

Q: How are we sure that we are able to align the group elements to form an orthonormal basis (also using the bracket notation)?

A: One can declare rules of the basis by treating each group element as a label and define a new inner product rule that only the group element squared in the bracket notation yields one; otherwise, it yields zero. In other words, Kronecker Delta. This shall be separated from the group multiplication rules.

Back to one of the simplest group \mathbb{Z}_2 . The group shall be more generally described as two elements g_1 and g_2 , which satisfy $g_1 = I$ and $g_1 \cdot g_2 = g_2$, $g_2 \cdot g_2 = g_1$. One can then decide how to represent this group by various representations. In our first encounter of \mathbb{Z}_2 , we’ve already chosen a one-dimensional representation $D(g_1) = 1$, $D(g_2) = -1$. One can also have a trivial representation of $D(g_1) = D(g_2) = 1$, or an identity matrix of arbitrary dimension. We can also construct the regular representation using the procedure mentioned above; we have a table of multiplication mapping

$$\begin{pmatrix} g_1 \cdot g_1 & g_1 \cdot g_2 \\ g_2 \cdot g_1 & g_2 \cdot g_2 \end{pmatrix} = \begin{pmatrix} g_1 & g_2 \\ g_2 & g_1 \end{pmatrix}. \quad (2.1.6)$$

Then, the regular representation is

$$D(g_1) = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, D(g_2) = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad (2.1.7)$$

363 and it satisfies

$$D(g_1)D(g_1) = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = D(g_1 \cdot g_1) = D(g_1), \quad (2.1.8)$$

$$D(g_2)D(g_1) = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} = D(g_2 \cdot g_1) = D(g_2), \quad (2.1.9)$$

$$D(g_2)D(g_2) = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = D(g_2 \cdot g_2) = D(g_1) \dots \quad (2.1.10)$$

364 We can take a peek at how the simplest discrete symmetry works in the
 365 SM. For instance, in the SM, we have a positively charged pion and a
 366 negatively charged pion, represented by $|\pi^\pm\rangle$. Defining a charged
 367 conjugation operator \hat{C} such that $\hat{C}|\pi^\pm\rangle = |\pi^\mp\rangle$. In the basis of the two
 368 states $|\pi^+\rangle, |\pi^-\rangle$, \hat{C} is simply $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$. Together with the identity matrix I ,
 369 $\{I, \hat{C}\}$ is a representation of the \mathbb{Z}_2 group. If the commutator $[\hat{C}, \hat{H}] = 0$,
 370 then the Hamiltonian is invariant under this group, which means the
 371 system respects this symmetry. This means many of the properties of π^+
 372 are the same as that of π^- . One can also construct a linear combination of
 373 pions to be the eigenstate of \hat{C} with eigenvalues ± 1 for $1/\sqrt{2}(|\pi^+\rangle \pm |\pi^-\rangle)$. As
 374 $\hat{C}^2 = I$, the eigenvalue can only be ± 1 . Since the only non-trivial
 375 “operation” of \mathbb{Z}_2 is \hat{C} (or g_2 that has $g_2 \cdot g_2 = g_1 = I$), we can call it a parity.
 376 One can claim these two eigenstate charge conjugations are even/odd.
 377 There are many other parity/conjugation symmetries in the SM.

378 2.2 LIE ALGEBRA

379 Now, we turn to a somewhat more complex class of group, which is
 380 extremely powerful in helping us understand quantum field theory and
 381 particle physics, the **continuous group**. One can imagine a successful
 382 reapplication of all the previous tricks in the discrete group in the previous
 383 section, but you can immediately see some inconvenience that the linear
 384 vector space spanned by the group elements where I want to construct
 385 regular representation is infinite, and hence, the representations are
 386 infinite-dimensional. The continuity comes to the rescue.

387 Given that the group is continuous, we can claim it smoothly depends
 388 on some continuous parameter α , which itself can be multi-dimensional
 389 (we call it **degrees**). This implies the group elements can span in a number
 390 of independent directions.

$$g \rightarrow g(\alpha). \quad (2.2.1)$$

391 Because α now labeled the group elements, we can also use it to label the
 392 elements in a representation $D(\alpha)$. As a group, identity exist; we demand
 393 that identity group in a desirable (which we are constructing, or assuming

the existence for now) representation is an identity (matrix) and the smooth parameters α taking value zero is the identity,

$$g(0) = I, \quad (2.2.2)$$

and hence for a representation $D(\alpha)$,

$$D(\alpha)|_{\alpha=0} = 1. \quad (2.2.3)$$

Here 1 is a n-by-n (the dimension of this representation $D(\alpha)$) identity matrix. Now, due to smoothness and continuity, we can reach group elements near the identity by a small “distance” $d\alpha$ via the expansion,

$$D(d\alpha) = 1 + id\alpha_a X_a + \dots \quad (2.2.4)$$

$$X_a \equiv -i \frac{\partial}{\partial \alpha_a} D(\alpha)|_{\alpha=0} \quad (2.2.5)$$

Here (and hereafter, unless otherwise specified), we use the “Einstein Summation” that all repeated indices are summed over. You can see a spans over all the degrees where identity is continuously connected, and X_a determines “how” the group elements span away from identity in each degree. X_a s are called **generators** of the group. For the α parametrization that is sufficiently unique (i.e., being parsimonious, that one needs to specify all α_a s to specify a group element), X_a will be independent. Again, you can already get a sense that a represents independent directions, the group element can span near the identity element, and X_a specifies how it spans. We include i here for general convenience; if the representation $D(\alpha)$ is unitary, the generator X_a will be Hermitian and vice versa.

How about when we are somewhat far away from identity? And how about very far away from identity? We hope to arrive at some representation that is consistent and nice, which, hopefully, will also have some nice smoothness in the expansion as well. A natural choice is to assume the representation follows the exponential parametrization of the continuous group. One can reach group elements further away from identity by repeated small excursions,

$$D(d\alpha) = 1 + i d\alpha_a X_a, \text{ such that} \quad (2.2.6)$$

$$D(\alpha) = \lim_{k \rightarrow \infty} \left(1 + i \frac{\alpha_a}{k} X_a \right)^k = \exp [i\alpha_a X_a]. \quad (2.2.7)$$

One can see that the generators here obey natural addition and multiplications, which are in the linear vector space.

We can immediately find usage of this representation in our familiar physics case. For simple free parameter wavefunction (let’s not worry about normalization as we can either wave-packet it or normalize over infinity with proper counting), we have,

$$\langle x|p \rangle = e^{ipx} \quad (2.2.8)$$

Under the translation of $x \rightarrow x - a$, the coordinate system is shifted by $+a$. One can define a unitary operator (which is a group element of the translation group) in this representation,

$$\hat{U}(a) = e^{-ia\hat{P}} \quad (2.2.9)$$

and we have

$$\langle x|\hat{U}(a)|p\rangle = \langle x|e^{-ia\hat{P}}|p\rangle = \langle x|e^{-iap}|p\rangle = e^{ip(x-a)}. \quad (2.2.10)$$

Clearly, the translation a is the continuous parameter that the group elements smoothly (exponentially) depend on, and the momentum operator \hat{P} is the generator of the spatial translation group.

In fact, for three spatial dimensions, there are three independent translation directions. Their generators commute with each other.

$$[\hat{P}_i, \hat{P}_j] = 0, \quad (2.2.11)$$

implying that

$$\hat{U}(a)\hat{U}(b) = \hat{U}(a+b), \quad (2.2.12)$$

satisfying simple addition rules on consecutive translations. The momentum operator \hat{P} is Hermitian. Hence, the translation operator is indeed unitary. In the relativistic case, the four-dimensional energy momentum operator is similar.

How can we use it? Suppose a system with Hamiltonian \hat{H} , satisfying

$$[\hat{P}, \hat{H}] = 0, \quad (2.2.13)$$

we know the linear momentum is conserved. On the other hand, it is completely equivalent to say, in this case,

$$[\hat{U}(a), \hat{H}] = 0, \quad (2.2.14)$$

and the system is translational invariant. In fact, the commutation of the generator with Hamiltonian directly means a corresponding symmetry of the system exists. The system shall be invariant under the symmetry transformation. The above fact is closely tied to the Noether theorem, which we will revisit later in this note.

But life is not always as simple as this above case. As one is fully entitled to, when spanning further away from identity, higher order terms should enter, and the parameter dependence might not be as (conveniently) simple as $D(\alpha)D(\beta) = D(\alpha + \beta)$ for two consecutive operations. Let's examine it. In general, without loss of generality, we can say

$$D(\alpha)D(\beta) = D(\gamma), \quad (2.2.15)$$

452 that consecutive group multiplication gives me another group member.
 453 Then, we work out small operations first,

$$e^{i\alpha_a X_a} e^{i\beta_b X_b} = e^{i\gamma_c X_c}, \quad (2.2.16)$$

$$i\gamma_c X_c = \ln [1 + e^{i\alpha_a X_a} e^{i\beta_b X_b} - 1], \quad (2.2.17)$$

$$(2.2.18)$$

454 Denoting the $K \equiv e^{i\alpha_a X_a} e^{i\beta_b X_b} - 1$, we have,

$$K = (1 + i\alpha_a X_a - \frac{1}{2}(\alpha_a X_a)^2 + \dots)(1 + i\beta_b X_b - \frac{1}{2}(\beta_b X_b)^2 + \dots) - 1 \quad (2.2.19)$$

$$= i\alpha_a X_a + i\beta_b X_b - \frac{1}{2}(\alpha_a X_a)^2 - \frac{1}{2}(\beta_b X_b)^2 - \alpha_a X_a \beta_b X_b + \dots \quad (2.2.20)$$

455 On the other hand,

$$\ln [1 + K] = K - \frac{1}{2}K^2 + \frac{1}{3}K^3 \dots \quad (2.2.21)$$

456 For infinitesimal operation, $K \simeq 0$, we can keep track of the leading terms.
 457 In particular, counting α and β in the same order and tracking up to the
 458 second order involves keeping track of the leading terms of $-\frac{1}{2}K^2$ term.

$$\begin{aligned} i\gamma_c X_c &= i\alpha_a X_a + i\beta_b X_b - \frac{1}{2}(\alpha_a X_a)^2 - \frac{1}{2}(\beta_b X_b)^2 - \alpha_a X_a \beta_b X_b - \frac{1}{2}(i\alpha_a X_a + i\beta_b X_b)^2 + \dots \\ &= i(\alpha + \beta)_a X_a - \frac{1}{2}[\alpha_a X_a, \beta_b X_b] + \dots \end{aligned} \quad (2.2.22)$$

$$= i(\alpha + \beta)_a X_a - \frac{1}{2}\alpha_a \beta_b [X_a, X_b] + \dots \quad (2.2.23)$$

459 These somewhat non-surprising results (we already learned the
 460 exponential operator identities in quantum mechanics) have very nice
 461 features. First, one can see that if the generators commute with each other
 462 $[X_a, X_b] = 0$, then $\gamma = \alpha + \beta$, we get back to the simplest product rule.
 463 Second, since we are discussing the general arbitrary excursions from
 464 identities labeled by α and β , the key features of the group structure are
 465 encoded independently of them. They are embedded in the commutator of
 466 generators,

$$[X_a, X_b] = if_{abc} X_c. \quad (2.2.24)$$

467 This is the **Lie Algebra** of the group. f_{abc} encodes the full information of
 468 the groups and is called **structure constant**. Clearly, $f_{abc} = -f_{bac}$. It
 469 captures the full information in the following sense: when we do
 470 excursions from the identity via the exponential parameters, it tells us
 471 what additional modifications we need to include to be consistent with the
 472 group multiplication. One might worry about higher order terms, but they
 473 are explicitly constructible as higher order commutators (non-commutator

part cancels) are just repeated usage of f_{abc} . Once the group is specified, we know the structure constant. The rest is actually to study different representations (all in the exponential parameterization as done here), equivalently different sets of X_a , that are consistent with the Lie algebra.

Before I give the first example. I want to mention two other useful results. One can show or encounter Jacobi Identity in various forms, e.g.,

$$[X_a, [X_b, X_c]] + \text{cyclic permutation} = 0. \quad (2.2.25)$$

Implementing the structure constant, Jacobi Identity means,

$$f_{ade}f_{dbc} + f_{bde}f_{dca} + f_{cde}f_{dab} = 0. \quad (2.2.26)$$

One can further show that the structure constants themselves, as generators, form a representation of the group,

$$(A_a)_{bc} = -if_{abc} \quad (2.2.27)$$

automatically satisfies

$$[A_a, A_b] = if_{abc}A_c. \quad (2.2.28)$$

This representation is called **adjoint representation**. It is an n -by- n dimensional representation of the group, and n is the degree of the group.

Let's begin with one simple example: for the structure constant f_{abc} of ϵ_{ijk} , where $i, j, k \in 1, 2, 3$, what generators can satisfy this and what physical system can we use it for? They are the structure constants for the rotation! ϵ_{ijk} is the fully antisymmetric tensor, with $\epsilon_{123} = 1$.

One can construct an adjoint representation directly (let me call it J_i , and $(J_i)_{jk} = -i\epsilon_{ijk}$, such that),

$$J_1 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -i \\ 0 & i & 0 \end{pmatrix}, J_2 = \begin{pmatrix} 0 & 0 & i \\ 0 & 0 & 0 \\ -i & 0 & 0 \end{pmatrix}, J_3 = \begin{pmatrix} 0 & -i & 0 \\ i & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}. \quad (2.2.29)$$

These are generators for rotations around three axes in the linear vector space by $(\hat{x}, \hat{y}, \hat{k})$, and the smooth parameters of these three rotations are the three Euler angles. These are the generators of the fundamental representation of the $SO(3)$ group, a three-degree special orthogonal group.

Certainly, we can ask if a two-dimensional irreducible representation exists. The answer is yes, and we know that well, the Pauli matrices. So, a two-dimensional representation of the rotational group can be written as,

$$J_1 = \frac{\sigma_1}{2} = \frac{1}{2} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, J_2 = \frac{\sigma_2}{2} = \frac{1}{2} \begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix}, J_3 = \frac{\sigma_3}{2} = \frac{1}{2} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (2.2.30)$$

They also satisfy the Lie algebra with the structure constant ϵ_{ijk} . Once again, we are already familiar with this in quantum mechanics. These are

the spin operators for spin-1/2 fermions, which obey the same commutation relations of the angular momentum operators and are part of the total angular momentum. From the group theory point of view, these are the generators in the fundamental representation of $SU(2)$, the two-degree special unitary group. One can see these generators in different representations are closely related. In fact, $SO(3)$ is isomorphic to $SU(2)/\{I, -I\}$ ², which means there is one-to-one mapping between these two.

What about one-dimensional representation? One can find the only way to satisfy the Lie algebra is the trivial representation, where all group elements are represented by identity (here in 1-D, just the number 1). The generators are nothing but

$$J_1 = J_2 = J_3 = 0. \quad (2.2.31)$$

What's so useful about this knowledge? In fact, most directly, we understand how to write down the spin projection operators for different systems. For a spin-0 particle, it is a spin singlet, and it transforms in the 1-d representation under rotation. For a spin-1/2 particle, it is a spin doublet (two-level system of s_i eigenvalue $\pm 1/2$), and it transforms in the 2-d representation under rotation. For a spin-1 particle, we call it a vector, and it is a spin triplet (three-level system of s_i eigenvalue of 0, ± 1). We can understand and show the symmetries and conserved currents when expressing the system in such basis and representations. Without any surprise, one can develop a method to write down a $(2s + 1)$ by $(2s + 1)$ representation for a spin- s system. Note that this is applying the representation to one of the quantum properties of a state, which we call spin. We can also *separately* apply these generators and exponential representations to other properties of a state, such as its orbital angular momentum. While the generators can be of the same form, they act on different subspaces of the Hilbert space of the wave function.

A curious reader may ask what happens to a possible generator of the identity matrix. In the exponential representation, the trial generator is no longer the identity matrix but zero. The exponential of zero is identity, which is a trivial representation. The inclusion of the identity matrix in the generator will actually change the group from special to non-special, so $SU(2)$ becomes $U(2)$ and $SO(3)$ becomes $O(3)$.

While one can see that $SU(2)$ and $SO(3)$ have generators follow the angular momentum operator rule, $SU(N)$ and $SO(N)$ have other usages. For instance, $SO(N)$ group is a set of continuous (real) linear transformations that leaves the inner product of N real numbers invariant. The group elements in the fundamental, irreducible representation would

²a quotient group of $SU(2)$

540 be a real N by N matrix D ,

$$\begin{pmatrix} x_1 \\ x_2 \\ \dots \\ x_N \end{pmatrix} \rightarrow \begin{pmatrix} x'_1 \\ x'_2 \\ \dots \\ x'_N \end{pmatrix} = D \begin{pmatrix} x_1 \\ x_2 \\ \dots \\ x_N \end{pmatrix} \quad (2.2.32)$$

541 The invariance of the inner product means

$$\vec{x} \cdot \vec{x} = \vec{x}' \cdot \vec{x}', \quad (2.2.33)$$

542 which requires that

$$D^T \cdot D = \mathbb{1}. \quad (2.2.34)$$

543 Here, you can see why they are called orthogonal groups. The new vectors
 544 x'_i are orthogonal to each other if the original basis is orthonormal. For
 545 $SO(N)$, there are N^2 real parameters in the matrices, and the above
 546 condition removes N parameters (from equating the diagonal elements
 547 r.h.s.) and $N(N-1)/2$ parameters (from equating the off-diagonal elements
 548 of the r.h.s., and it is symmetric). Hence, the total number of free
 549 parameters is $N^2 - N - N(N-1)/2 = N(N-1)/2$. This is the degree of this
 550 group that a linear transformation leaves the inner product invariant, and
 551 hence the number of generators. If we apply this to $SO(3)$, we know there
 552 should be $3 \times (3-1)/2 = 3$ generators. If we, instead, only require the inner
 553 product to multiply proportional to the original inner product for arbitrary
 554 vectors, one boundary condition is removed, and hence, there will be one
 555 more generator, and that would be the identity matrix, which changes the
 556 length of the inner product.

557 Similarly, for $SU(N)$, the linear mapping keeps the inner product of $\vec{x}^* \cdot$
 558 \vec{x} invariant. It would be $2N^2$ degree of freedom, and after subtracting the
 559 conditions from $D^\dagger D = \mathbb{1}$ ³ one gets $N(N+1)/2$ degrees. For the case of
 560 $SU(2)$, one gets $2 \times (2+1)/2$ generators.

561 2.3 POINCARÉ GROUP AND 562 LORENTZ GROUP

563 Poincaré group is a semi-direct product of the translation group and the
 564 Lorentz group. Its importance in relativistic physics is self-evident.
 565 However, it is not a simple group, and we will have to deal with some
 566 richer structures and features than the abovementioned groups. The
 567 Lorentz group is a transformation that preserves distance with a metric

³ $D^\dagger D$ being real removes $(N^2 - N)$ d.o.f., as the diagonal entries are automatically diagonal. Then, the rest of the counting is the same as the $SO(N)$ case.

568 containing opposite signs between the distance in time and the distance in
569 space.

$$x^\mu \rightarrow x'^\mu = \Lambda^\mu_\nu x^\nu, \text{ such that } x^2 \equiv x \cdot x \equiv g_{\mu\nu} x^\mu x^\nu = x'^2 \quad (2.3.1)$$

570 Here $g_{\mu\nu}$ is the metric (in particle physics, most of us take the most-minus
571 metric)

$$g_{\mu\nu} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}, \quad (2.3.2)$$

572 and the covariant form $g^{\mu\nu}$ is a matrix of exactly the same form. We can
573 use them to raise and lower Lorentz indices.

574 The Poincaré group further is translational invariance, which preserves
575 separations between points.

$$x^\mu \rightarrow x'^\mu = \Lambda^\mu_\nu x^\nu + a^\mu, \quad (2.3.3)$$

$$y^\mu \rightarrow y'^\mu = \Lambda^\mu_\nu y^\nu + a^\mu, \text{ such that} \quad (2.3.4)$$

$$(x - y)^2 = (x' - y')^2 \quad (2.3.5)$$

576 One can perform two Poincaré transformations consecutively, and it
577 should be equal to one single Poincaré transformation with some
578 parametric correspondence. We have

$$x^\mu \rightarrow x'^\mu \rightarrow x''^\mu \quad (2.3.6)$$

$$\begin{aligned} x''^\mu &= \Lambda''^\mu_\nu x'^\nu + a''^\mu \\ &= \Lambda'^\mu_\rho (\Lambda^\rho_\nu x^\nu + a'^\nu) + a''^\mu \\ &= (\Lambda' \Lambda)^\mu_\rho x^\rho + (\Lambda'^\mu_\nu a'^\nu + a''^\mu), \end{aligned} \quad (2.3.7)$$

579 implying

$$x''^\mu = \Lambda''^\mu_\nu x^\nu + a''^\mu, \quad (2.3.8)$$

580 with

$$\Lambda'' = \Lambda' \Lambda, \quad a''^\mu = \Lambda'^\mu_\nu a'^\nu + a''^\mu. \quad (2.3.9)$$

581 You can see here why we say the Poincaré group is a semi-direct product
582 group of the translation group and Lorentz group.

583 We are already somewhat familiar with the translation group and its
584 generator from the previous section. Let's focus on the Lorentz part and
585 then put them together.

586 The Lorentz part could be an $SO(4)$ group if the metric would be an
587 identity matrix. However, due to the opposite sign in the space and time
588 part in the metric, it becomes a non-compact group. For instance, we can
589 elongate the spatial part arbitrarily so long that we enlarge the time part
590 correspondingly; the separation is unchanged between points, hence

satisfying the Poincaré invariance. It demands some more complex treatment that we discuss in this section.

Note that we are working in a representation of Lorentz and Poincaré group that we are most familiar with; that is how the 3+1-D coordinates transform under these symmetries. This representation is called the defining representation of these groups. What we want to do in this section is to understand the important properties of these symmetries in this representation and then derive the structure constants of these groups. This will allow us to start understanding different representations of these groups, and they play important roles in defining fundamental particles.

The Lorentz symmetry require $g_{\mu\nu}x^\mu x^\nu = g_{\mu\nu}x'^\mu x'^\nu$ and hence that

$$g_{\mu\nu}\Lambda^\mu_\alpha\Lambda^\nu_\beta = g_{\alpha\beta}, \quad (2.3.10)$$

and the identity group element is

$$\delta^\mu_\nu = 1_{4\times 4}. \quad (2.3.11)$$

Now multiply Eq. (2.3.10) by $g^{\alpha\rho}$ and contract the indices that get raised and lowered, we get,

$$\Lambda^\rho_\nu\Lambda^\nu_\beta = \delta^\rho_\beta. \quad (2.3.12)$$

Noting that⁴,

$$\Lambda^\rho_\nu = (\Lambda^T)^\rho_\nu = (\Lambda^{-1})^\rho_\nu, \quad (2.3.13)$$

and hence

$$\Lambda^T\Lambda = 1_{4\times 4}, \quad (2.3.14)$$

implying

$$\det[\Lambda^T\Lambda] = \det[\Lambda^T]\det[\Lambda] = \det[\Lambda]^2 = 1, \quad \det[\Lambda] = \pm 1. \quad (2.3.15)$$

Clearly, there is an identity group element in this representation, δ^μ_ν , and it has a determinant of +1. We call the subset of Lorentz transformation with determinant +1 *proper* and those with determinant -1 *improper*.

Furthermore, a subset of Lorentz transformation has Λ^0_0 component greater or equal to unity⁵, $\Lambda^0_0 \geq 1$, which preserves the time direction⁶, are called *orthochronous*.

The Proper Orthochronous (P.O.) subset of Lorentz transformation forms a subgroup of Lorentz transformation; it satisfies all the group properties

⁴here the notation is a bit confusing, but I don't have a better way. $\Lambda^\rho_\nu = (\Lambda^\rho_\nu)^T$ (Here I am transposing the matrix, not only the elements). One transposes Λ to get Λ^T , exchanging the corresponding labels.

⁵From Eq. (2.3.10), we have $(\Lambda^0_0)^2 - (\Lambda^i_0)^2 = 1$

⁶For Lorentz four vectors with time-like separation $x^2 \geq 0$, the positive time vectors will remain positive time for arbitrary Lorentz transformation satisfying $\Lambda^0_0 \geq 1$.

and, in addition, can reach every group element by successive infinitesimal transformations. The P.O. subgroup of the Lorentz group is very powerful and convenient for us since we can use Lie Algebra to fully describe it. It is denoted as $SO^+(1,3)$, and for most of the time of this lecture, when we refer to Lorentz symmetry and invariance, in particular when talking about generators, we meant this subgroup. This subgroup is also “central” in the sense that it is continuously connected to the identity group element δ^μ_ν , as the identity is P.O. as well.

Before we dive into the representations. It is useful to comment on the other parts of the Lorentz group. They are connected to the P.O. subgroup by two meaningful operators (and their product), which are the time-reversal operator \mathcal{T} and the parity operator \mathcal{P} ,

$$\mathcal{T}^\mu_\nu = \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad \mathcal{P}^\mu_\nu = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}, \quad (2.3.16)$$

be mindful of the Lorentz indices. Clearly, one can reach all four domains of the Lorentz transformation by applying $\mathbb{1}$, \mathcal{T} , \mathcal{P} and \mathcal{PT} on the P.O. subgroup.

Now, let’s talk about the exponential representation of the P.O. subgroup of the Lorentz group in its defining representation. For short, I will call this a representation of the Lorentz group, instead of calling it a representation of the P.O. subgroup of the Lorentz group.

We can play the game of infinitesimal transformations, and let’s express them (later on, I will add translation),

$$\Lambda^\mu_\nu = \delta^\mu_\nu + \alpha^\mu_\nu. \quad (2.3.17)$$

Then, given Eq. (2.3.12), we have

$$(\delta^\alpha_\nu + \alpha^\alpha_\nu + \dots)(\delta^\nu_\beta + \alpha^\nu_\beta + \dots) = \delta^\alpha_\beta, \quad (2.3.18)$$

and tracking to order α (the continuous parameter that a representation of the Lorentz group smoothly depends on), we get

$$\alpha^\alpha_\beta + \alpha^\alpha_\beta = 0. \quad (2.3.19)$$

Hence, this excursion parameter is antisymmetric,

$$\alpha_{\alpha\beta} = -\alpha_{\beta\alpha}. \quad (2.3.20)$$

Q & As

Q: (Chimeln) How many generators does the Lorentz group have?

Q: (Chimeln) How many generators does the Poincaré group have?

Q: (Chimeln) Is this representation unitary? Are all the generators in this representation hermitian?

Q: [Eq. \(2.3.17\)](#) is a Taylor expansion near the identity; how do I ensure there are no new directions (d.o.f.) when I have a large expansion?

A: By assuming the existence of exponential representation and the continuity as well as smoothness of the dependence, I can cover all the group elements smoothly connected to identity by repeated infinitesimal transformations. In this sense, for this fully connected group space, there are no new directions. In other words, the emergence of a new d.o.f., which is linearly independent from the original ones, means discontinuity in the group element. However, for the full Lorentz group, there are large expansions, discontinuously connected to the P.O. subgroup, by the group elements of \mathcal{T} , \mathcal{P} , and \mathcal{PT} .

Since it is real-valued, as an anti-symmetry 4-by-4 number, there are six d.o.f, and hence six generators $M^{\mu\nu}$, who are also anti-symmetric. We already know what these six d.o.f. are: three rotations and three boosts. Of course, one can also combine these directions linearly into new directions, but it is merely a linear transformation of the generators and expansion parameters.

We can denote the Lorentz generators as $\hat{M}^{\mu\nu}$, such that,

$$\hat{U}(\alpha) = e^{\frac{i}{2}\alpha_{\mu\nu}\hat{M}^{\mu\nu}}, \quad (2.3.21)$$

For an appropriate choice of generators, one can identify the three rotations and three boosts as

$$\hat{K}_i = \hat{M}_{i0} \quad (2.3.22)$$

$$\hat{J}_i = -\frac{1}{2}\epsilon_{ijk}\hat{M}^{jk}. \quad (2.3.23)$$

Correspondingly, the three boost parameters α_{0i} are the rapidity in each direction, and the three rotation parameters $n_i \equiv \epsilon_{ijk}\alpha_{jk}$ are the three Euler angles.

We already know the translation and its generator from [Eq. \(2.2.9\)](#). Hence, we can write down a representation (the defining representation in 3+1D, in the exponential form, for the P.O. subgroup) for the Poincaré transformation,

$$\hat{U}(a, \Lambda) = \hat{U}(a)\hat{U}(\Lambda). \quad (2.3.24)$$

Note *the order matters*; as usual, we apply the transformations from the rightmost operator to the left. There is no ambiguity in this notation that,

$$\begin{aligned} \hat{U}(a) &= \hat{U}(a, 1), \\ \hat{U}(\Lambda) &= \hat{U}(0, \Lambda). \end{aligned} \quad (2.3.25)$$

In the above equations, we did a small “trick” that we replaced the expansion parameter from α^μ_ν to Λ^μ_ν , and there shouldn’t be any ambiguities as there is one (set of parameters) to one correspondence between α and Λ for P.O. subset of Lorentz transformations. We are simply too used to the Lorentz transformations in their defining representation, and the space-time transformation is quite a rudimentary building block in particle physics.

We know exactly how Poincaré transformation should be from Eqs. (2.3.6) to (2.3.9), and hence we can derive the commutation relations, the Lie Algebra, between the generators.

$$\hat{U}(a)\hat{U}(a') = \hat{U}(a + a') \quad (2.3.26)$$

$$\hat{U}^{-1}(\Lambda)\hat{U}(\Lambda')\hat{U}(\Lambda) = \hat{U}(\Lambda^{-1}\Lambda'\Lambda) \quad (2.3.27)$$

$$\hat{U}(\Lambda)\hat{U}(a) = \hat{U}(\Lambda a)\hat{U}(\Lambda) \quad (2.3.28)$$

Expanding the above with infinitesimal transformations labeled by a and $\alpha_{\mu\nu}$, with their corresponding generators \hat{P} and $\hat{M}^{\mu\nu}$, at linear order on the expansion parameters, allows us to find the consistent commutation relations.

We can use Eq. (2.3.26) to find that,

$$[\hat{P}_\mu, \hat{P}_\nu] = 0. \quad (2.3.29)$$

We have already done this from a different angle in the earlier section in Eqs. (2.2.11) and (2.2.12).

We can use Eq. (2.3.27) to find that⁷,

$$[\hat{M}^{\mu\nu}, \hat{M}^{\rho\sigma}] = i \left(\hat{M}^{\mu\rho} g^{\nu\sigma} + \hat{M}^{\nu\sigma} g^{\mu\rho} - \hat{M}^{\mu\sigma} g^{\nu\rho} - \hat{M}^{\nu\rho} g^{\mu\sigma} \right). \quad (2.3.30)$$

So far, the above contains the commutator between generators within translation and Lorentz. What about the commutator between the translation generators and Lorentz generators? We can use Eq. (2.3.28). Here, I will take a slower approach and show it step by step to set an example. In fact, I will derive Eq. (2.3.28) first from the defining property in Eq. (2.3.7) and then derive the commutation relations between \hat{P} and \hat{M} .

⁷Note that the argument of the right-hand side $\Lambda^{-1}\Lambda'\Lambda$ is still a ground member of the Lorentz Group in the defining representation. Hence, it can be written as $\Lambda'' \equiv \Lambda^{-1}\Lambda'\Lambda = \delta + \alpha''$ and hence the exponential form in $\hat{U}(\Lambda'')$.

$$\hat{U}(a', \Lambda') \hat{U}(a, \Lambda) = \hat{U}(\Lambda' a + a', \Lambda' \Lambda) \quad (2.3.31)$$

$$\begin{aligned} \hat{U}(a') \hat{U}(\Lambda') \hat{U}(a) \hat{U}(\Lambda) &= \hat{U}(\Lambda' a + a') \hat{U}(\Lambda' \Lambda) \\ &= \hat{U}(a') \hat{U}(\Lambda' a) \hat{U}(\Lambda') \hat{U}(\Lambda) \end{aligned} \quad (2.3.32)$$

$$\hat{U}(\Lambda') \hat{U}(a) = \hat{U}(\Lambda' a) \hat{U}(\Lambda') \quad (2.3.33)$$

$$\hat{U}(\Lambda) \hat{U}(a) \hat{U}^{-1}(\Lambda) = \hat{U}(\Lambda a), \quad (2.3.34)$$

686 We derive Eq. (2.3.28) here in Eq. (2.3.33) after left-multiplying $\hat{U}^{-1}(a')$ and
 687 right-multiplying $\hat{U}^{-1}(\Lambda)$ on both side of the equation. Since Λ no longer
 688 appears in Eq. (2.3.33) and below (arrived by right-multiplying it with
 689 $\hat{U}^{-1}(\Lambda')$), we can unprime Λ' and rewrite it as Λ .

690 Now, we can expand according to the plan and the expansion
 691 parameters⁸,

$$\hat{U}(\Lambda) e^{i\hat{P}^\alpha a_\alpha} \hat{U}(\Lambda^{-1}) = e^{i\hat{P}^\mu \Lambda_\mu^\alpha a_\alpha} \quad (2.3.35)$$

$$\begin{aligned} (1 + \frac{i}{2} \alpha_{\mu\nu} \hat{M}^{\mu\nu} + \dots) \hat{P}^\alpha (1 - \frac{i}{2} \alpha_{\mu\nu} \hat{M}^{\mu\nu} + \dots) &= \hat{P}^\mu \Lambda_\mu^\alpha = \hat{P}^\mu (\delta_\mu^\alpha + \alpha_\mu^\alpha) \\ \frac{i}{2} \alpha_{\mu\nu} [\hat{M}^{\mu\nu}, \hat{P}^\alpha] &= \frac{1}{2} (\hat{P}^\mu \alpha_{\mu\nu} g^{\nu\alpha} - \hat{P}^\nu \alpha_{\mu\nu} g^{\mu\alpha}) \\ [\hat{M}^{\mu\nu}, \hat{P}^\alpha] &= -i (\hat{P}^\mu g^{\nu\alpha} - \hat{P}^\nu g^{\mu\alpha}). \end{aligned} \quad (2.3.36)$$

692 From the first line to the second line, we match the leading non-trivial
 693 term in a_α (it is a_α to the first order). From the second line to the third line,
 694 we match the leading non-trivial term in α (it is α to the first order). We
 695 use $g^{\nu\alpha}$ to lower the α index to match $\alpha_{\mu\nu}$ on both sides and get rid of it in
 696 the last line. We relabel the dummy indices and split the r.h.s. into two
 697 terms to make it an anti-symmetric form, which is merely an aesthetic
 698 choice.

699 The commutation relation between the generator of translation and
 700 Lorentz transformation, shown in Eq. (2.3.36), shows that these two
 701 transformations are “coupled”, which are not simple ones that commute
 702 and can be exchanged. One sees again the semi-direct product nature of
 703 the Poincaré group from the Translation group and the Lorentz group.

704 Having understood the above, the first direct application of this
 705 space-time symmetry is how we define a one-particle state. One-particle
 706 state is the building block for particle phenomenology, and there are a lot
 707 to discuss in terms of interaction picture, canonical normalization, cluster
 708 decomposition, etc. For now, we simply want to have a clear definition of
 709 isolated pristine one-particle state that we can prepare and later

⁸Note that I can always relabel dummy indices that are contacted over, so I choose them “wisely”

experiment with. As already done in quantum mechanics, we want to find a maximally compatible set of operators such that we can define the wavefunction uniquely as a set of eigenstates of these operators that spans over the whole Hilbert space. A natural choice is to use the momentum operator, which in most cases is observable, and further, in high energy limit, the particle-wave duality does tell us the particle description provides a simpler calculation basis. Clearly, the Lorentz invariant operator $\hat{P}_\mu \hat{P}^\mu$ also commutes with \hat{P}^μ , and this is the mass squared operator \hat{M}^2 . How about Lorentz generators $M^{\mu\nu}$? Unfortunately, they do not commute with \hat{P}^μ , as shown in Eq. (2.3.36). We will show in the next section that a subset of $\hat{M}^{\mu\nu}$ commutes with \hat{P}^μ and \hat{M}^2 , and they are identified as the spin quantum numbers for a point-like one-particle state. So, the operator set is

$$\{\hat{P}^\mu, \hat{M}^2, \text{subset of } \hat{M}^{\mu\nu} \text{ that commutes}\} \quad (2.3.37)$$

and we are able to specify the one-particle state by the eigenvalues,

$$|p, m, \text{some additional space-time quantum numbers}\rangle. \quad (2.3.38)$$

In the next section, we see how we can specify these additional space-time quantum numbers.

For now, the one-particle state have some degeneracy but we can still understand their behaviors, e.g.,

$$\begin{aligned} \hat{P}^\mu |p, m\rangle &= p^\mu |p, m\rangle \\ \hat{U}(a) |p, m\rangle &= e^{i\hat{P}^\mu a_\mu} |p, m\rangle \\ \hat{M}^2 |p, m\rangle &= m^2 |p, m\rangle \end{aligned} \quad (2.3.39)$$

What about $\hat{U}(\Lambda) |p, m\rangle$? From Eq. (2.3.35), write down the term linear in \hat{P} , and replace Λ^{-1} by Λ , we find

$$\begin{aligned} \hat{U}(\Lambda^{-1}) \hat{P}^\mu \hat{U}(\Lambda) &= \Lambda^\mu{}_\nu \hat{P}^\nu \\ \hat{P}^\mu \hat{U}(\Lambda) &= \hat{U}(\Lambda) \Lambda^\mu{}_\nu \hat{P}^\nu \\ \hat{P}^\mu \left(\hat{U}(\Lambda) |p, m\rangle \right) &= \hat{U}(\Lambda) \left(\Lambda^\mu{}_\nu \hat{P}^\nu |p, m\rangle \right) \\ &= \Lambda^\mu{}_\nu p^\nu \left(\hat{U}(\Lambda) |p, m\rangle \right) \end{aligned} \quad (2.3.40)$$

Hence $\left(\hat{U}(\Lambda) |p, m\rangle \right)$ is an eigenstate of \hat{P}^μ with eigenvalue of $\Lambda^\mu{}_\nu p^\nu$.

Consequently, we conclude,

$$\hat{U}(\Lambda) |p, m\rangle = |\Lambda p, m\rangle. \quad (2.3.41)$$

However, as I declared that there are degeneracies (more non-trivial compatible operators with \hat{P}^μ exist from the Poincaré group, the resulting one-particle state not merely changed its momentum. We will get the fuller result in the next section.

2.4 LITTLE GROUP, WIGNER ROTATION, SPIN, AND HELICITY

Instead of directly informing the reader of the representations and corresponding constructions of the Poincaré group for particle physics, here I show how to get general properties of the representation. In particular, we introduce the idea of Wigner's Little Group transformation and show how they can be utilized to specify the one-particle states. Another reason for me to introduce it here is that Little Group does buy us additional handles in particle physics, in particular the scattering processes, and in the modern helicity amplitude program.⁹

From the previous section, we know the key is to identify a subset of Lorentz generators $\hat{M}^{\mu\nu}$ that commutes with the translation generator \hat{P}^μ (and hence it automatically commutes with \hat{M}^2). We know Lorentz boost would change the momentum of a state, and hence, it would not commute with the momentum operator, which is the generator for translation. Would the answer be simply a rotation? The answer is no. Physically, the result of translating and then rotating would be different if one rotates and then translates (unless the translation direction is the same as the spatial rotation vector direction or the translation is in the time direction). Specifically,

$$[\hat{J}_k, \hat{P}^\mu] = -\frac{1}{2}\epsilon_{ijk}[\hat{M}_{ij}, \hat{P}^\mu] = i\frac{1}{2}\epsilon_{ijk}(\hat{P}^i g^{j\mu} - \hat{P}^j g^{i\mu}) = \begin{cases} 0 & \text{if } \mu = 0, k \\ i\epsilon_{ijk}\hat{P}^i g^{jj} & \text{if } \mu = j, j! = 0, k \end{cases} \quad (2.4.1)$$

We can see that two rotation generators commute with the translation generator. They will reappear later in a slightly subtle context.

We will need to construct some other generators of the Poincaré group, which depends on the original generators. We understand that the additional operators we can use to specify the one-particle state cannot be boost, which does not commute with \hat{P} , and it cannot be rotation, which does not commute with \hat{P} in general, as shown in Eq. (2.4.1).

We introduce the concept of the little group [Wig39; Kim16], which is a subgroup of the Lorentz group that leaves the momentum four-vector invariant. (Isn't this what we are asking for in Eq. (2.3.37)?) An easy way to understand and construct the generators of this subgroup would be to see their effects on momentum in their so-called standard form, which is $\tilde{p} = (m, 0, 0, 0)$ for massive particles and $\tilde{p} = (p, 0, 0, p)$ for massless particles. The resulting Lie algebra for these two cases will be related but somewhat

⁹For reviews of the modern helicity amplitude program and the usage of helicity weights, see, e.g., [Che18; EH13].

different.¹⁰

2.4.1 MASSIVE CASE

Let's denote the little group, a subset of Lorentz transformations Λ , as $\tilde{\Lambda}$. We then have,

$$\tilde{\Lambda} p^\mu = p^\mu. \quad (2.4.2)$$

In the rest frame, $\tilde{p} = (m, 0, 0, 0)$. We then have,

$$\tilde{\Lambda}^\mu{}_\nu \tilde{p}^\nu = (\delta^\mu{}_\nu + \tilde{\alpha}^\mu{}_\nu) \tilde{p}^\nu = \tilde{p}^\nu, \quad (2.4.3)$$

implying

$$\begin{aligned} \tilde{\alpha}^\mu{}_\nu \tilde{p}^\nu &= \tilde{\alpha}_{\mu\nu} \tilde{p}^\nu = 0 \\ \tilde{\alpha}^{i0} &= -\tilde{\alpha}^{0i} = 0. \end{aligned} \quad (2.4.4)$$

From the above, we can see that three degrees are possible in the little group. We are their generators, and how do we write them down with only Lorentz covariant quantities? Clearly, the index of $\mu = 0$ is special in this case, and the only covariant way to pick it would be multiplying the \hat{P}^μ operator. We can define,

$$\tilde{\alpha}_{\mu\nu} \hat{M}^{\mu\nu} = \hat{M}^{\mu\nu} \tilde{\alpha}_{\mu\nu} = -\frac{1}{2} \hat{M}^{\mu\nu} \epsilon_{\mu\nu\rho\sigma} \hat{P}^\rho n^\sigma, \quad (2.4.5)$$

so there is a one-to-one mapping between the possible non-zero little group expansion directions $\tilde{\alpha}_{\mu\nu}$ and the parameter n^σ for particle at rest. When Lorentz transformation act on a massive particle at rest (rest frame, r.f.),

$$-\frac{1}{2} \hat{M}^{\mu\nu} \epsilon_{\mu\nu\rho\sigma} \hat{P}^\rho n^\sigma \stackrel{\text{r.f.}}{=} -\frac{1}{2} \hat{M}^{\mu\nu} \epsilon_{\mu\nu 0\sigma} n^\sigma m = -\frac{1}{2} \hat{M}^{ij} \epsilon_{ijk} n^k m = m \hat{J}_k n^k. \quad (2.4.6)$$

Interestingly, in the rest frame of a massive particle, the little group transformation is equivalent to rotation (the rotation generator is defined as in Eq. (2.3.23)). From the above, one can define a Lorentz covariant generalized rotation using a new operator constructed from Eq. (2.4.5), called the Pauli-Lubanski operator, which is the generator of this generalized rotation,

$$\hat{W}_\sigma \equiv -\frac{1}{2} \epsilon_{\mu\nu\rho\sigma} \hat{M}^{\mu\nu} \hat{P}^\rho, \quad (2.4.7)$$

$$\hat{U}(\tilde{\Lambda}) |\tilde{p}, m\rangle \equiv \left(1 + i n^\sigma \hat{W}_\sigma + \dots\right) |\tilde{p}, m\rangle. \quad (2.4.8)$$

¹⁰For a more systematic discussion, see Refs. [Kim16; Wei05].

792 We have

$$\hat{U}(\tilde{\Lambda}) |\tilde{p}, m\rangle = |\tilde{\Lambda}\tilde{p}, m\rangle = |\tilde{p}, m\rangle, \quad (2.4.9)$$

$$\hat{W}_\sigma |\tilde{p}, m\rangle = m\hat{J}_i |\tilde{p}, m\rangle = ms_i |\tilde{p}, m\rangle, \quad (2.4.10)$$

$$\hat{W}_\sigma \hat{W}^\sigma |\tilde{p}, m\rangle = -m^2 \hat{J}^2 |\tilde{p}, m\rangle = m^2 s(s+1) |\tilde{p}, m\rangle, \quad (2.4.11)$$

793 where s and s_i are the eigenvalues of the \hat{J}^2 and \hat{J}_i operators, from
 794 **Eq. (2.4.6)** and the fact $[\hat{J}_i, \hat{J}_j] = i\epsilon_{ijk}\hat{J}_k$ (thus all the nice relations with
 795 raising and lowering operators and relations to \hat{J}^2).

Q & As

Q: (Chimeln) What is $\hat{W}_\sigma \hat{P}^\sigma$?

Q: (Chimeln) What is $[\hat{W}_\sigma, \hat{P}^\mu]$?

797 We can work out its commutation relations,

$$\hat{W}_\sigma \hat{P}^\sigma = 0, \quad (2.4.12)$$

$$[\hat{W}_\sigma, \hat{P}^\mu] = 0, \quad (2.4.13)$$

$$[\hat{M}_{\mu\nu}, \hat{W}_\sigma] = -i \left(\hat{W}_\mu g_{\nu\sigma} - \hat{W}_\nu g_{\mu\sigma} \right). \quad (2.4.14)$$

798 The last line above is the same commutation relation as momentum
 799 operators \hat{P} , which is a Lorentz four-vector. It implies that \hat{W}_σ transforms
 800 as a Lorentz four-vector. Now, we feel more confident to say \hat{W}^σ is a
 801 generalization of the three rotation generators in a Lorentz covariant
 802 four-vector form. Its commutation relation is,

$$[\hat{W}_\mu, \hat{W}_\nu] = i\epsilon_{\mu\nu\rho\sigma} \hat{W}^\rho \hat{P}^\sigma. \quad (2.4.15)$$

803 Knowing \hat{W}_σ is a Lorentz four-vector with three effective angles n^σ (one
 804 degree is mapped out by p^0), we can now declare and complete
 805 **Eq. (2.3.37)**,

$$\{\hat{P}^\mu, \hat{M}^2, \frac{\hat{W}_\sigma \hat{W}^\sigma}{m^2}, \frac{\hat{W}_\sigma}{m}\}. \quad (2.4.16)$$

806 Equivalently, we can choose to the set to be

$$\{\hat{P}^\mu, \hat{M}^2, \hat{J}^2, \hat{J}_i\}. \quad (2.4.17)$$

807 Note that $\hat{W}^\sigma \hat{W}_\sigma$ is a Lorentz invariant quantity, m^2 is a Lorentz invariant
 808 quantity. One might worry about the compatibility between \hat{J}_i with \hat{P}^μ , we
 809 can declare that we want to specify the quantum state in the rest frame of
 810 the particle, or we declare that we want to specify the eigenvalue of \hat{W}_σ/m .
 811 In fact, an even better definition, motivated by our observation in

Eq. (2.4.1), is to define the J_k along the momentum direction, and we have proven that they commute. Nevertheless, for massive particles, we can fully specify the state (according to its space-time symmetries) via

$$|p, m, s, s_i\rangle. \quad (2.4.18)$$

Now that we have fully specified the state (that there is no degeneracy in space-time symmetries), we can understand how such a state transforms in a general Lorentz transformation beyond those specified in Eq. (2.3.39). Before that, let's define a pure boost operator,

$$\begin{aligned} \hat{L}(p)^\mu \tilde{p}^\nu &= p^\mu, \\ \hat{U}(L(p)) |\tilde{p}, m\rangle &= |p, m\rangle, \end{aligned} \quad (2.4.19)$$

which transforms a massive state (momentum as a Lorentz four-vector) from its rest frame to a frame with momentum of p^μ .

We want to complete the results in Eq. (2.3.39),

$$\begin{aligned} \hat{U}(\Lambda) |p, m, s, s_i\rangle &= (\text{unknown coefficient}) |\Lambda p, m, s, s'_i\rangle \\ &= \hat{U}(\Lambda) \hat{U}(L(p)) |\tilde{p}, m, s, s_i\rangle \end{aligned} \quad (2.4.20)$$

$$\begin{aligned} &= \hat{U}(L(\Lambda p)) \hat{U}^{-1}(L(\Lambda p)) \hat{U}(\Lambda) \hat{U}(L(p)) |\tilde{p}, m, s, s_i\rangle \\ &= \hat{U}(L(\Lambda p)) \hat{U}(L^{-1}(\Lambda p)) \hat{U}(\Lambda) \hat{U}(L(p)) |\tilde{p}, m, s, s_i\rangle \end{aligned} \quad (2.4.21)$$

We understand the first (from right to left) three operators do not change the four-momentum at rest¹¹,

$$\begin{aligned} L^{-1}(\Lambda p) \Lambda L(p) \tilde{p} &= L^{-1}(\Lambda p) \Lambda p \\ &= L^{-1}(\Lambda p) (\Lambda p) \\ &= \tilde{p}. \end{aligned} \quad (2.4.22)$$

Hence, $L^{-1}(\Lambda p) \Lambda L(p)$ is a member of the little group, and we can define, in short, a **Wigner Rotation**,

$$R(\Lambda p, p) \equiv L^{-1}(\Lambda p) \Lambda L(p). \quad (2.4.23)$$

We can work its impact on a state by inserting a complete set of states,

$$\begin{aligned} \hat{U}(R(\Lambda p, p)) |\tilde{p}, m, s, s_i\rangle &= \sum_{s'_i} |\tilde{p}, m, s, s'_i\rangle \langle \tilde{p}, m, s, s'_i| \hat{U}(R(\Lambda p, p)) |\tilde{p}, m, s, s_i\rangle \\ &= \sum_{s'_i} D_{s'_i, s_i}^{(s)}(R(\Lambda p, p)) |\tilde{p}, m, s, s'_i\rangle. \end{aligned} \quad (2.4.24)$$

¹¹Here, we can work with the parameters of the operators directly in the defining representation.

827 Here $D_{s'_i, s_i}^{(s)}(R(\Lambda p, p))$ are Wigner rotation matrices in the spin s
 828 representation, with¹²

$$D_{s'_i, s_i}^{(s)}(R(\Lambda p, p)) \equiv (e^{-i\theta \hat{n} \cdot \vec{J}^{(s)}})_{s'_i, s_i}. \quad (2.4.25)$$

829 The matrix elements are called Wigner d-functions.

830 Note that \hat{W}_σ is the generator for such class of rotations with three
 831 d.o.f.s, and they are the rotation generator in the defining representation
 832 in the rest frame, multiplying mass m . Hence, for a general representation,
 833 the Wigner rotation in the rest frame would rotate the spin components
 834 without changing the total spin quantum number as an eigenstate of the
 835 \hat{J}^2 operator with eigenvalue $s(s+1)$. Consequently, the Wigner rotation
 836 matrices correspond to a general rotation with a compound direction
 837 specified by \hat{n} (a normalized three vector) and an angle θ in the spin- s
 838 representation. We somewhat anticipate this result for a few reasons. The
 839 boost and rotation generators do not commute with each other; hence,
 840 also the Wigner rotation does not change momentum; it is a nest operation
 841 of boost, general Lorentz transformation (including boost and rotation),
 842 and then boost the new result momentum back to the rest four-vector. It
 843 cannot correspond to another boost as it will change the momentum, so it
 844 can only change the spin quantum number via rotation. Furthermore,
 845 we've also seen that the generator of Wigner rotation, the Pauli-Lubanski
 846 operator, is a three-rotation, and hence we expect it to appear here. Last
 847 but not least, since this Wigner rotation is acting on a particle at rest, it is
 848 identical to the rotation operator, and consequently, we get this result.

849 With the Wigner rotation matrices (elements) in Eq. (2.4.24), we can
 850 continue the evaluation in Eq. (2.4.21),

$$\begin{aligned} \hat{U}(\Lambda) |p, m, s, s_i\rangle &= \hat{U}(L(\Lambda p)) \hat{U}(L^{-1}(\Lambda p)) \hat{U}(\Lambda) \hat{U}(L(p)) |\tilde{p}, m, s, s_i\rangle \\ &= \hat{U}(L(\Lambda p)) \hat{U}(R(\Lambda p, p)) |\tilde{p}, m, s, s_i\rangle \\ &= \sum_{s'_i} D_{s'_i, s_i}^{(s)}(R(\Lambda p, p)) \hat{U}(L(\Lambda p)) |\tilde{p}, m, s, s'_i\rangle \\ &= \sum_{s'_i} D_{s'_i, s_i}^{(s)}(R(\Lambda p, p)) |\Lambda p, m, s, s'_i\rangle. \end{aligned} \quad (2.4.26)$$

851 We can see that general Lorentz transformations change the momentum of
 852 a particle and rotate its spin. Now, let's move to the massless case and try
 853 to unify the results in a more commonly used version.

854 2.4.2 MASSLESS CASE

855 For the massless case, we can go through a similar procedure, but we will
 856 find the commuting rotation corresponds to two degrees of rotation,

¹²The minus sign in the exponent is simply a result of the Lorentz inner product on the space components.

857 motivating us to define a new quantity called helicity. For simplicity of the
858 discussion, we work with the standard form of the momentum¹³

$$\tilde{p}^\mu \equiv (p, 0, 0, p). \quad (2.4.27)$$

859 We first check how Pauli-Lubanski operator acts on a massless state with
860 momentum \tilde{p} . From $\hat{W}_\sigma \hat{P}^\sigma = 0$ (see Eq. (2.4.13)),¹⁴

$$\hat{W}_\sigma \hat{P}^\sigma |\tilde{p}, 0\rangle = (\hat{W}_0 + \hat{W}_3)p |\tilde{p}, 0\rangle = 0. \quad (2.4.28)$$

861 On the other hand, $\hat{W}^\sigma \hat{W}_\sigma$ is a Lorentz Invariant (L.I.), mass dimension two
862 quantity. The only scale in $|\tilde{p}, 0\rangle$ is p , which is not L.I. but covariant. One
863 can Lorentz-boost along or against the z-direction to get arbitrary
864 momentum (including approaching zero not zero). Hence, the only possible
865 outcome of this operator on the state is zero,¹⁵

$$\hat{W}_\sigma \hat{W}^\sigma |\tilde{p}, 0\rangle = 0, \quad (2.4.29)$$

$$\begin{aligned} (\hat{W}_0)^2 - (\hat{W}_1)^2 - (\hat{W}_2)^2 - (\hat{W}_3)^2 |\tilde{p}, 0\rangle &= 0, \\ (\hat{W}_1)^2 + (\hat{W}_2)^2 |\tilde{p}, 0\rangle &= 0. \end{aligned} \quad (2.4.30)$$

866 To obtain the last line, we use that (with Eq. (2.4.28)),

$$(\hat{W}_0)^2 - (\hat{W}_3)^2 |\tilde{p}, 0\rangle = (\hat{W}_0 - \hat{W}_3)(\hat{W}_0 + \hat{W}_3) |\tilde{p}, 0\rangle = 0. \quad (2.4.31)$$

867 and (from Eq. (2.4.15))

$$[\hat{W}_0, \hat{W}_3] |\tilde{p}, 0\rangle = i\epsilon_{03ij} \hat{W}^i \hat{P}^j |\tilde{p}, 0\rangle = 0. \quad (2.4.32)$$

868 Furthermore, we can get, from Eq. (2.4.15),

$$\begin{aligned} [\hat{W}_1, \hat{W}_2] |\tilde{p}, 0\rangle &= i(\epsilon_{1203} \hat{W}^0 \hat{P}^3 + \epsilon_{1230} \hat{W}^3 \hat{P}^0) |\tilde{p}, 0\rangle \\ &= ip\epsilon_{1203} (\hat{W}^0 - \hat{W}^3) |\tilde{p}, 0\rangle \\ &= ip\epsilon_{12}^{03} (\hat{W}_0 + \hat{W}_3) |\tilde{p}, 0\rangle \\ &= 0. \end{aligned} \quad (2.4.33)$$

869 Similarly, one can get that,¹⁶

$$[\hat{W}_0, \hat{W}_1] |\tilde{p}, 0\rangle = i\epsilon_{0123} \hat{W}^2 \hat{P}^3 |\tilde{p}, 0\rangle = -ip\hat{W}_2 |\tilde{p}, 0\rangle \quad (2.4.34)$$

$$[\hat{W}_3, \hat{W}_1] |\tilde{p}, 0\rangle = i\epsilon_{3120} \hat{W}^2 \hat{P}^0 |\tilde{p}, 0\rangle = ip\hat{W}_2 |\tilde{p}, 0\rangle \quad (2.4.35)$$

$$[\hat{W}_0, \hat{W}_2] |\tilde{p}, 0\rangle = i\epsilon_{0213} \hat{W}^1 \hat{P}^3 |\tilde{p}, 0\rangle = ip\hat{W}_1 |\tilde{p}, 0\rangle \quad (2.4.36)$$

$$[\hat{W}_3, \hat{W}_2] |\tilde{p}, 0\rangle = i\epsilon_{3210} \hat{W}^1 \hat{P}^0 |\tilde{p}, 0\rangle = -ip\hat{W}_1 |\tilde{p}, 0\rangle. \quad (2.4.37)$$

¹³There is no magic here; you can choose general momentum, and we need to adjust the statements from z-direction to a linear combination proportional to the momentum direction.

¹⁴Note here the Lorentz indices are not contracted yet so no need to implement the metric.

¹⁵For a more systematic discussion, see Weinberg Vol.I [Wei05].

¹⁶Note that I flipped the sign when I lower the indices in the spatial part, consistent with the metric choice.

870 Hence we have,

$$[\hat{W}_3, \hat{W}_1 \pm i\hat{W}_2] |\tilde{p}, 0\rangle = \pm p(\hat{W}_1 \pm i\hat{W}_2) |\tilde{p}, 0\rangle \quad (2.4.38)$$

871 This is our familiar raising and lowering operator. Let's label a state with
872 its eigenvalue on \hat{W}_3/p , $|\tilde{p}, 0, s_3\rangle$. Then, one have

$$\hat{W}_3/p(\hat{W}_1 \pm i\hat{W}_2) |\tilde{p}, 0, s_3\rangle = (s_3 \pm 1)(\hat{W}_1 \pm i\hat{W}_2) |\tilde{p}, 0, s_3\rangle \quad (2.4.39)$$

873 and expect $(\hat{W}_1 \pm i\hat{W}_2) |\tilde{p}, 0, s_3\rangle \propto |\tilde{p}, 0, s_3 \pm 1\rangle$, i.e., the raising and lowering
874 operators change the eigenstate of s_3 to $s_3 \pm 1$. However, there is a different
875 possibility for Eq. (2.4.39) to hold, that is,

$$(\hat{W}_1 \pm i\hat{W}_2) |\tilde{p}, 0, s_3\rangle = 0. \quad (2.4.40)$$

876 It has to be the case that if I can raise or lower my state without getting a
877 null state, at least one of the following be nonzero,

$$(\hat{W}_1 \pm i\hat{W}_2)(\hat{W}_1 \mp i\hat{W}_2) |\tilde{p}, 0, s_3\rangle \neq 0, \quad (2.4.41)$$

878 which contradicts Eq. (2.4.30). Consequently, we have

$$\hat{W}_1 |\tilde{p}, 0\rangle = \hat{W}_2 |\tilde{p}, 0\rangle = 0. \quad (2.4.42)$$

879 One can already get a hint of the difference between the massive case
880 and the massless case in the sense that,

$$-\hat{W}_i \hat{W}^i / m^2 |\tilde{p}, m, s, s_i\rangle = \hat{J}^2 |\tilde{p}, m, s, s_i\rangle = s(s+1) |\tilde{p}, m, s, s_i\rangle, \quad (2.4.43)$$

$$-\hat{W}_i \hat{W}^i / p^2 |\tilde{p}, 0, s_3\rangle = \hat{W}_3^2 / p^2 |\tilde{p}, 0, s_3\rangle = s_3^2 |\tilde{p}, 0, s_3\rangle. \quad (2.4.44)$$

881 The total spin is just the spin in the z -direction for massless particles with
882 standard momentum; the action of raising and lowering operators gets
883 zero. So, I do not need another total spin quantum number for the
884 massless case.

885 Furthermore, from the definition,

$$\begin{aligned} \hat{W}_1 |\tilde{p}, 0\rangle &= -\frac{1}{2}\epsilon_{\mu\nu 01}\hat{M}^{\mu\nu}\hat{P}^0 - \frac{1}{2}\epsilon_{\mu\nu 31}\hat{M}^{\mu\nu}\hat{P}^3 |\tilde{p}, 0\rangle \\ &= p(\hat{J}_1 + \hat{K}_2) |\tilde{p}, 0\rangle \end{aligned} \quad (2.4.45)$$

$$\begin{aligned} \hat{W}_2 |\tilde{p}, 0\rangle &= -\frac{1}{2}\epsilon_{\mu\nu 02}\hat{M}^{\mu\nu}\hat{P}^0 - \frac{1}{2}\epsilon_{\mu\nu 32}\hat{M}^{\mu\nu}\hat{P}^3 |\tilde{p}, 0\rangle \\ &= -p(\hat{J}_2 + \hat{K}_1) |\tilde{p}, 0\rangle \end{aligned} \quad (2.4.46)$$

$$\begin{aligned} \hat{W}_0 |\tilde{p}, 0\rangle &= -\frac{1}{2}\epsilon_{\mu\nu 30}\hat{M}^{\mu\nu}\hat{P}^3 |\tilde{p}, 0\rangle \\ &= p\hat{J}_3 |\tilde{p}, 0\rangle \end{aligned} \quad (2.4.47)$$

$$\begin{aligned} \hat{W}_3 |\tilde{p}, 0\rangle &= -\frac{1}{2}\epsilon_{\mu\nu 03}\hat{M}^{\mu\nu}\hat{P}^0 |\tilde{p}, 0\rangle \\ &= -p\hat{J}_3 |\tilde{p}, 0\rangle, \end{aligned} \quad (2.4.48)$$

Combining with the commutation relations between rotation and boost generators, We can see clearly many equations in this section, e.g., $\hat{W}_0 + \hat{W}_3 |\tilde{p}, 0\rangle = 0$, confirming Eq. (2.4.28), and $[\hat{W}_1, \hat{W}_2] |\tilde{p}, 0\rangle = 0$, confirming Eq. (2.4.33).

The above motivates us to define a quantum number as an eigenstate of the Pauli-Lubanski operator in the non-trivial direction of \hat{W}_0 , \hat{W}_3 or a linear combination that is not $\hat{W}_0 + \hat{W}_3$. Furthermore, we know \hat{W}_σ transforms as a Lorentz 4-vector, from the commutation relations with Lorentz generator in Eq. (2.4.14). There are no other Lorentz covariant symbols in momentum space that are intrinsic to the state (e.g., not a reference four-vector); we do foresee that $\hat{W}_\sigma \propto \hat{P}_\sigma$. We can specify a new (mass-)dimensionless quantity, **helicity**,

$$\lambda \equiv -\frac{\hat{W}_\mu n^\mu}{\hat{P}_\mu n^\mu}. \quad (2.4.49)$$

Here n^μ is a reference vector, and we typically choose it as $n^\mu = (1, 0, 0, 0)$. This corresponds to defining the one-particle eigenstate according to its eigenvalue with respect to the rotation generator \hat{J}_3 . Compared to the case of massive particles, we lose the freedom to define the one-particle state with its eigenvalue on an arbitrary spatial axis. On the other hand, this quantum number choice is still consistent with what we find in Eq. (2.4.1), that rotation along the momentum direction is compatible with the momentum operator. In other words, we already foresee that we want to project the spin quantum number along the momentum direction from the very beginning of this section! This choice is independent of whether the underlying state is massive or massless. A more *commonly used* **helicity** definition, which is equivalent to Eq. (2.4.49) with the n^μ choice, is

$$\lambda \equiv -\frac{\hat{W}_0 n^0}{\hat{P}_0 n^0} = -\frac{\hat{W}_0 P^0}{\hat{P}_0 P^0} = \frac{\hat{W}_i \cdot \hat{P}^i}{|\vec{P}|^2} = \frac{\vec{J} \cdot \vec{p}}{|\vec{p}|}, \quad (2.4.50)$$

where we have used Eq. (2.4.12) Both definitions are consistent with each other, and they are *invariant* under rotation. one can see that this rightmost definition can be extendable to massive particles, which we will explore later.

Unlike the case for massive particles, one cannot define raising and lowering operators for helicity. Hence, a particle does not need to fill the full $2s + 1$ spin d.o.f.s. A one-particle state with specific helicity lives on its own. Why do we often see opposite helicities for massless particles? It is because we require some more symmetries of the theory.

Helicity is rotation invariant but boosts variant quantity, but the spin projection is a rotation variant quantity. It is even convenient many times to define the last quantum number for massive particles' helicity instead of spin projection.

923 For massless particles, the total spin quantum number is its spin
 924 projection along the momentum direction. The algebra is closed, so there
 925 are no $2s + 1$ states. Note that, under parity transformation,

$$\begin{aligned}\vec{p} &\xrightarrow{\mathcal{P}} -\vec{p} \\ s_i &\xrightarrow{\mathcal{P}} s_i \\ \lambda &\xrightarrow{\mathcal{P}} -\lambda.\end{aligned}\tag{2.4.51}$$

926 Spin projection being invariant as its generator \hat{M}^{ij} gets flipped twice.
 927 Helicity, as the spin projection along the momentum direction, gets flipped
 928 under parity. If we require a theory (with its corresponding one-particle
 929 state) to be an eigenstate of the parity transformation, we would need the
 930 existence of the opposite helicity state. Hence, we often see massless states
 931 come in helicity pairs.

932 One can unify the notation for one-particle state for massive and
 933 massless particles, both expressed as

$$|p, m, s, \lambda\rangle, \tag{2.4.52}$$

934 and for massless state $m = 0$, s is redundant with $s = |\lambda|$ (again, note that
 935 $-\hat{W}_i \hat{W}^i = W_3^2$, differ from the massive case), but it doesn't hurt to keep it.

936 Finally, we can also unify the discussion regarding a general Lorentz
 937 transformation's effect on a one-particle state. With helicity, we are
 938 motivated to exploit rotation, as both S and λ are invariant under rotation.
 939 We can define a general standard form momentum,

$$\tilde{p}' \equiv (E, 0, 0, p), \tag{2.4.53}$$

940 as we put all the momentum in the positive z-direction. For massless case,
 941 $E = p$ and massive case has $E = \sqrt{m^2 + p^2}$. For the massless case, the
 942 helicity is already defined to be projected in this direction. For the massive
 943 case, with a general momentum $p \equiv \Lambda \tilde{p}$, we can write

$$\begin{aligned}\hat{U}(\Lambda) |\tilde{p}, m, s, \lambda\rangle &= \hat{U}(R(p, \tilde{p}')) \hat{U}(L(\tilde{p}')) |\tilde{p}, m, s, \lambda\rangle \\ &= \hat{U}(R(p, \tilde{p}')) |\tilde{p}', m, s, \lambda\rangle.\end{aligned}\tag{2.4.54}$$

944 Basically, for a general momentum direction, one can define the helicity for
 945 a massive particle in its rest frame as the spin projection along the positive
 946 z-direction. The general momentum p is rewritten as a rotation of the
 947 general standard form momentum \tilde{p}' . The first step of boost along the
 948 positive z-direction will not change the spin projection along the positive
 949 z-direction; it transforms the state momentum from rest \tilde{p} to a new one
 950 that has the same three-momentum magnitude as that of the final p . Then
 951 one just performs a rotation, which again does not change λ . If we do this

in another order, it often involves other helicity values (for massive cases). For a three momentum with Euler angle (ϕ, θ) , the rotation $R(p, \tilde{p}')$ can be expressed as a series of operation,

$$\hat{R}(p, \tilde{p}') = \hat{R}_3(-\phi)R_2(-\theta)R_3(\phi) = e^{-i\phi\hat{J}_3}e^{-i\theta\hat{J}_2}e^{+i\phi\hat{J}_3}. \quad (2.4.55)$$

For massless case, one again can boost the standard form momentum further along the z-direction to match the magnitude of the resulting three-momentum $p = \Lambda\tilde{p} = R(\Lambda\tilde{p}, \tilde{p}')L(\tilde{p}', p)\tilde{p}$. The resulting effect is analogous to the above.

REFERENCES

- [Che18] Clifford Cheung. “TASI Lectures on Scattering Amplitudes”. In: *Proceedings, Theoretical Advanced Study Institute in Elementary Particle Physics : Anticipating the Next Discoveries in Particle Physics (TASI 2016): Boulder, CO, USA, June 6-July 1, 2016*. Ed. by Rouven Essig and Ian Low. 2018, pp. 571–623. DOI: [10.1142/9789813233348_0008](https://doi.org/10.1142/9789813233348_0008). arXiv: [1708.03872](https://arxiv.org/abs/1708.03872) [hep-ph].
- [EH13] Henriette Elvang and Yu-tin Huang. “Scattering Amplitudes”. In: (Aug. 2013). arXiv: [1308.1697](https://arxiv.org/abs/1308.1697) [hep-th].
- [Geo99] Howard Georgi. *Lie algebras in particle physics*. 2nd ed. Vol. 54. Reading, MA: Perseus Books, 1999.
- [Kim16] Y.S. Kim. *Wigner’s little groups*. <https://ysfine.com/wigner/wiglittle.pdf>. 2016.
- [Wei05] Steven Weinberg. *The Quantum theory of fields. Vol. 1: Foundations*. Cambridge University Press, June 2005. ISBN: 978-0-521-67053-1, 978-0-511-25204-4. DOI: [10.1017/CBO9781139644167](https://doi.org/10.1017/CBO9781139644167).
- [Wig39] Eugene P. Wigner. “On Unitary Representations of the Inhomogeneous Lorentz Group”. In: *Annals Math.* 40 (1939). Ed. by Y. S. Kim and W. W. Zachary, pp. 149–204. DOI: [10.2307/1968551](https://doi.org/10.2307/1968551).