

10.2 CLASSICAL FIELDS: THE LEGENDRE TRANSFORMATION AND SURFACE TERMS

Two pieces of classical machinery are used from the first lectures on: the Legendre transformation, which takes us from a Lagrangian to a Hamiltonian, and the rule that surface terms may be dropped when we vary an action.

THE LEGENDRE TRANSFORMATION

DEFINITION AND MATHEMATICAL FORMULATION

- **Single Variable:** For a convex function $f(x)$, the Legendre transform $g(p)$ is defined as:

$$g(p) = px(p) - f(x(p)), \quad \text{where} \quad p = \frac{df}{dx}.$$

Here, $x(p)$ is obtained by inverting $p = df/dx$.

- **Multiple Variables:** For $f(x_1, \dots, x_n)$, the transform for the first m variables is:

$$g(p_1, \dots, p_m, x_{m+1}, \dots, x_n) = \sum_{i=1}^m p_i x_i - f, \quad p_i = \frac{\partial f}{\partial x_i}.$$

The untransformed variables $x_{m+1}, \dots, x_n$ ride along as parameters.

KEY PROPERTIES

- **Involution:** Applying the transform twice returns f if f is convex.
- **Convexity Requirement:** Ensures uniqueness of $x(p)$. Non-convex functions require the Legendre–Fenchel transform.
- **Derivatives swap roles:** $dg/dp = x$, the inverse of $df/dx = p$. The slope of one function is the variable of the other.

APPLICATIONS IN PHYSICS

Classical mechanics. The transition from the **Lagrangian** $L(q, \dot{q}, t)$ to the **Hamiltonian** $H(q, p, t)$:

$$p_i = \frac{\partial L}{\partial \dot{q}_i}, \quad H = \sum_i p_i \dot{q}_i - L.$$

This yields Hamilton's first-order equations.

666 **Thermodynamics.** Thermodynamic potentials are related via Legendre
667 transforms:

$$\begin{aligned}H(S, p) &= U + pV, \\F(T, V) &= U - TS, \\G(T, p) &= H - TS.\end{aligned}$$

668 Each potential shifts control variables (e.g., $T \leftrightarrow S$). With the definition
669 above, F and G are *minus* the Legendre transforms; the overall sign is a
670 convention.

671 **Field theory.** From the **Lagrangian density** $\mathcal{L}(\phi, \partial_\mu \phi)$ to the **Hamiltonian**
672 **density**:

$$\pi(x) = \frac{\partial \mathcal{L}}{\partial \dot{\phi}}, \quad \mathcal{H} = \pi \dot{\phi} - \mathcal{L}.$$

673 GEOMETRIC INTERPRETATION

674 The Legendre transform constructs $g(p)$ as the maximum value of $px - f(x)$
675 for each p . This encodes $f(x)$ through its tangent lines (slopes p ,
676 intercepts $-g(p)$).

677 EXAMPLES

- 678 • Quadratic function: $f(x) = \frac{1}{2}kx^2 \rightarrow g(p) = \frac{p^2}{2k}$.
- 679 • Exponential function: $f(x) = e^x \rightarrow g(p) = p(\ln p - 1)$.

680 ADVANCED TOPICS

- 681 • **Symplectic Geometry:** Framework for Hamiltonian mechanics.
- 682 • **Contact Geometry:** Describes thermodynamic equilibrium spaces.
- 683 • **Hamilton–Jacobi Theory:** The generating function of the
684 canonical transformation to constant variables obeys a PDE, the
685 Hamilton–Jacobi equation.

686 In short, the Legendre transformation switches between conjugate
687 variables, enabling reformulations like Hamiltonian mechanics and
688 thermodynamic potentials. It preserves the information in the function
689 under convexity, adapting the description to the natural variables of the
690 problem.

691 DISCARDING SURFACE TERMS IN THE VARIATIONAL 692 PRINCIPLE

693 In quantum field theory, when deriving equations of motion such as the
694 Klein–Gordon equation via the principle of least action, boundary terms
695 arising from integration by parts are discarded under two justified
696 assumptions:

1. Vanishing Field Variations at Boundaries: The variational principle considers small perturbations $\delta\phi(x)$ around the classical field configuration. These variations are required to have *compact support*, meaning $\delta\phi(x) \rightarrow 0$ at spacetime infinity.¹ For example, when varying the action $S = \int d^4x \mathcal{L}(\phi, \partial_\mu\phi)$, integration by parts generates boundary terms of the form

$$\int d^4x \partial_\mu \left(\frac{\partial \mathcal{L}}{\partial(\partial_\mu\phi)} \delta\phi \right) = \oint_{\text{boundary}} d\Sigma_\mu \frac{\partial \mathcal{L}}{\partial(\partial_\mu\phi)} \delta\phi,$$

which vanish because $\delta\phi = 0$ on the boundary. This ensures surface terms do not contribute to the equations of motion, regardless of the original field $\phi(x)$'s behavior. The vanishing of $\delta\phi$ at boundaries is a formal requirement of the variational principle, not a physical assumption about $\phi(x)$.

2. Physical Boundary Conditions for Fields: While the field $\phi(x)$ itself might oscillate rapidly at infinity, physical consistency demands fields behave “nicely” at spacetime boundaries. For quantum fields, this is often enforced by:

- *Square-integrability:* $\phi \in L^2$. This controls the field on average; it does not by itself make $\phi(x)$ vanish pointwise at infinity (§10.5).
- *Tempered distributions:* Fields are assumed to not blow up uncontrollably (standard in axiomatic QFT).
- *Schwartz functions:* Fields and derivatives decay rapidly, ensuring boundary terms involving $\phi(x)$ are well-controlled.

Heuristically, rapid oscillations of $\phi(x)$ are further suppressed in the path integral formalism. Configurations with wildly oscillating phases e^{iS} destructively interfere (the *stationary phase approximation*), leaving only contributions near classical paths, where the action is stationary.

The Klein–Gordon equation arises as the condition for the action to be stationary under these variations. Crucially, boundary terms depend on $\delta\phi$ (not ϕ), and $\delta\phi$ vanishes at infinity by construction. Thus, discarding surface terms is valid in the variational framework, even if $\phi(x)$'s behavior at infinity appears ambiguous.

¹The physics argument: we want the amplitude between given initial and final states, which are asymptotic states. With those specified, there are no functional variations at infinity.