

10.15 SPINOR REPRESENTATIONS OF THE LORENTZ GROUP

This note connects the proper orthochronous Lorentz group $SO^+(1, 3)$, its Lie algebra $\mathfrak{so}(1, 3)$, its universal cover $SL(2, \mathbb{C})$, and the classification of its finite-dimensional representations. Conventions and the algebra

$$[J_i, J_j] = i\epsilon_{ijk}J_k, \quad [J_i, K_j] = i\epsilon_{ijk}K_k, \quad [K_i, K_j] = -i\epsilon_{ijk}J_k$$

are those of §10.3; a group element near the identity is $\exp(i\theta_i J_i + i\eta_i K_i)$, with rotation angles θ_i and rapidities η_i . $SO^+(1, 3)$ is the group of real 4×4 matrices Λ that preserve $g = \text{diag}(-1, +1, +1, +1)$ with $\det \Lambda = 1$ and $\Lambda^0_0 \geq 1$.

THE TEMPLATE: $SO(3)$ AND $SU(2)$

- **Algebras (local).** The Lie algebras are isomorphic, $\mathfrak{su}(2) \cong \mathfrak{so}(3)$: the same three generators J_i with $[J_i, J_j] = i\epsilon_{ijk}J_k$.
- **Groups (global).** The groups are not isomorphic. $SO(3)$ (real orthogonal 3×3 matrices) acts on vectors; $SU(2)$ (complex unitary 2×2 matrices of unit determinant) acts on 2-component spinors. $SU(2)$ is the **universal double cover** of $SO(3)$: the map is 2-to-1, with U and $-U$ giving the same rotation.

The double cover is visible in the action on a vector $\mathbf{x}$ written as the matrix $X = \mathbf{x} \cdot \boldsymbol{\sigma}$:

$$X \rightarrow X' = UXU^\dagger, \quad (-U)X(-U)^\dagger = UXU^\dagger.$$

Physically, a 2π rotation returns a vector to itself but multiplies a spinor by -1 ; a 4π rotation is needed to return a spinor to itself.

THE LORENTZ CASE: $SO^+(1, 3)$ AND $SL(2, \mathbb{C})$

The same pattern holds with six generators (three rotations J_i , three boosts K_i):

- **Algebras (local).** As 6-dimensional *real* Lie algebras, $\mathfrak{so}(1, 3) \cong \mathfrak{sl}(2, \mathbb{C})$, where $\mathfrak{sl}(2, \mathbb{C})$ is regarded as a real algebra with basis $\{\sigma_i/2, i\sigma_i/2\}$.
- **Groups (global).** $SL(2, \mathbb{C})$, the complex 2×2 matrices of unit determinant, is the universal double cover of $SO^+(1, 3)$: M and $-M$ map to the same Λ . Its defining representation and the complex conjugate of it are the 2-component Weyl spinors.

2267 A COMMON ERROR: $\mathfrak{so}(1, 3) \not\cong \mathfrak{su}(2) \oplus \mathfrak{su}(2)$

2268 It is tempting to write the Lorentz algebra as two copies of $\mathfrak{su}(2)$. As real
 2269 algebras this is wrong: $\mathfrak{su}(2) \oplus \mathfrak{su}(2) \cong \mathfrak{so}(4)$, the algebra of *Euclidean*
 2270 rotations in four dimensions. The difference is one sign. Take two
 2271 commuting $\mathfrak{su}(2)$'s with generators A_i , B_i , and set $J_i = A_i + B_i$, $K_i = A_i - B_i$.
 2272 Then

$$[K_i, K_j] = [A_i, A_j] + [B_i, B_j] = i\epsilon_{ijk}(A_k + B_k) = +i\epsilon_{ijk}J_k,$$

2273 whereas the Lorentz algebra has $-i\epsilon_{ijk}J_k$. So $\mathfrak{so}(1, 3) \not\cong \mathfrak{so}(4) \cong \mathfrak{su}(2) \oplus \mathfrak{su}(2)$
 2274 as real algebras, and because $[J_i, K_j] \neq 0$ the real algebra does not split
 2275 into commuting pieces.

2276 THE COMPLEXIFICATION TRICK

2277 Since the real algebra does not decompose, we complexify it: we allow
 2278 complex linear combinations of the generators, passing to $\mathfrak{so}(1, 3)_{\mathbb{C}}$, and
 2279 define

$$N_i = \frac{1}{2}(J_i + iK_i), \quad \tilde{N}_i = \frac{1}{2}(J_i - iK_i).$$

2280 The minus sign in $[K_i, K_j]$ is exactly what makes the two sets commute:

$$[N_i, N_j] = i\epsilon_{ijk}N_k, \quad [\tilde{N}_i, \tilde{N}_j] = i\epsilon_{ijk}\tilde{N}_k, \quad [N_i, \tilde{N}_j] = 0.$$

2281 The complexified algebra therefore does decompose:

$$\mathfrak{so}(1, 3)_{\mathbb{C}} \cong \mathfrak{sl}(2, \mathbb{C}) \oplus \mathfrak{sl}(2, \mathbb{C}) \cong \mathfrak{su}(2)_{\mathbb{C}} \oplus \mathfrak{su}(2)_{\mathbb{C}}.$$

2282 **Why this is allowed.** The finite-dimensional complex representations of a
 2283 real Lie algebra $\mathfrak{g}$ are in one-to-one correspondence with the
 2284 complex-linear representations of its complexification $\mathfrak{g}_{\mathbb{C}}$, and
 2285 irreducibility is preserved. So classifying the representations of the easy
 2286 algebra $\mathfrak{su}(2)_{\mathbb{C}} \oplus \mathfrak{su}(2)_{\mathbb{C}}$, which are pairs of ordinary angular-momentum
 2287 representations, classifies those of the physical algebra.

2288 **Counting dimensions.** $\mathfrak{su}(2)$ is a 3-dimensional real algebra (basis σ_i). Its
 2289 complexification $\mathfrak{su}(2)_{\mathbb{C}}$ is 3-dimensional over $\mathbb{C}$ and 6-dimensional over $\mathbb{R}$
 2290 (basis $\{\sigma_i, i\sigma_i\}$). So $\mathfrak{so}(1, 3)_{\mathbb{C}}$ is 6-dimensional over $\mathbb{C}$, i.e. a 12-dimensional
 2291 real space, with basis $\{\sigma_i^N, i\sigma_i^N\} \cup \{\sigma_i^{\tilde{N}}, i\sigma_i^{\tilde{N}}\}$.

2292 **The physical slice.** The physical algebra $\mathfrak{so}(1, 3)$ is a 6-dimensional real
 2293 slice of this 12-dimensional space. Inverting the definitions,

$$J_i = N_i + \tilde{N}_i, \quad K_i = -i(N_i - \tilde{N}_i),$$

2294 with *real* rotation angles and rapidities multiplying them. In a
 2295 finite-dimensional representation with N_i and $\tilde{N}_i$ Hermitian:

- 2296 • the rotation generators J_i are Hermitian, so rotations $\exp(i\theta_j J_j)$ are
 2297 unitary;

• the boost generators K_i are anti-Hermitian, so boosts $\exp(i\eta_j K_j) = \exp(\eta_j(N_j - \tilde{N}_j))$ are not unitary. This is not a choice: a finite-dimensional representation of the non-compact Lorentz group cannot be unitary (§10.3).

The six physical generators are this specific combination, not “two sets of Pauli matrices”.

A note on notation. Srednicki writes our $\tilde{N}_i$ as $N_i^\dagger$. On the Hilbert space of states, where the representation is unitary and J_i, K_i are Hermitian, this is a true adjoint. In the finite-dimensional representations below, both N_i and $\tilde{N}_i$ are represented by Hermitian matrices and $\tilde{N}_i \neq N_i^\dagger$; there the dagger is only a label. What survives is a relation between the group elements: the generators of the $(1, 2)$ representation are minus the Hermitian conjugates of those of the $(2, 1)$, so the matrices are related by $R = (L^\dagger)^{-1}$.

CLASSIFYING THE REPRESENTATIONS

Irreducible representations are labeled by a pair $(2n + 1, 2n' + 1)$, the dimensions under the N and $\tilde{N}$ algebras, with $n, n' = 0, \frac{1}{2}, 1, \dots$. Since $J_i = N_i + \tilde{N}_i$, the rotation content is spins $|n - n'|, \dots, n + n'$.

Representations with $n + n'$ an integer are representations of $SO^+(1, 3)$ itself; those with $n + n'$ half-integer are representations of $SL(2, \mathbb{C})$ only, and change sign under a 2π rotation.

$(2, 1)$: the left-handed Weyl spinor. $J_i = \frac{\sigma_i}{2}$, $K_i = -\frac{i\sigma_i}{2}$. Then $N_i = \frac{\sigma_i}{2}$ (dimension 2) and $\tilde{N}_i = 0$ (dimension 1). A left-handed spinor transforms as $\psi_L \rightarrow L\psi_L$, with $L = \exp(\frac{i}{2}\theta_i\sigma_i + \frac{1}{2}\eta_i\sigma_i) \in SL(2, \mathbb{C})$.

$(1, 2)$: the right-handed Weyl spinor. $J_i = \frac{\sigma_i}{2}$, $K_i = +\frac{i\sigma_i}{2}$. Then $N_i = 0$ and $\tilde{N}_i = \frac{\sigma_i}{2}$. A right-handed spinor transforms as $\psi_R \rightarrow R\psi_R$, with $R = \exp(\frac{i}{2}\theta_i\sigma_i - \frac{1}{2}\eta_i\sigma_i) = (L^\dagger)^{-1}$.

$(2, 1) \oplus (1, 2)$: the Dirac spinor. In the Weyl basis $\Psi = (\psi_L, \psi_R)^T$ transforms with $\text{diag}(L, R)$, generated by the blocks $J_i = \frac{1}{2}\text{diag}(\sigma_i, \sigma_i)$, $K_i = \frac{i}{2}\text{diag}(-\sigma_i, \sigma_i)$ of §10.3.

$(2, 2)$: the four-vector. A four-vector carries one index of each type.

Following Srednicki's $\sigma^\mu = (I, \boldsymbol{\sigma})$ contracted with the lower-index x_μ , map it to a 2×2 Hermitian matrix:

$$X = x_\mu \sigma^\mu = \begin{pmatrix} -x^0 + x^3 & x^1 - ix^2 \\ x^1 + ix^2 & -x^0 - x^3 \end{pmatrix}, \quad \det X = (x^0)^2 - |\mathbf{x}|^2 = -x_\mu x^\mu.$$

It transforms as

$$X \rightarrow X' = LX L^\dagger,$$

the $(2, 1)$ rule acting on the left index and its complex conjugate

(equivalent to the $(1, 2)$) on the right. Because $\det L = 1$, $\det X' = \det X$, and

2334 X' is again Hermitian, so this is a Lorentz transformation $x \rightarrow \Lambda(L)x$ of
 2335 the components; L and $-L$ give the same Λ . Its rotation content is
 2336 $\frac{1}{2} \otimes \frac{1}{2} = 0 \oplus 1$: the time component and a spatial vector.

Representation	$(d_N, d_{\tilde{N}})$	(n, n')	Transformation
Left-handed Weyl	$(2, 1)$	$(\frac{1}{2}, 0)$	$\psi_L \rightarrow L\psi_L$
Right-handed Weyl	$(1, 2)$	$(0, \frac{1}{2})$	$\psi_R \rightarrow (L^\dagger)^{-1}\psi_R$
Four-vector	$(2, 2)$	$(\frac{1}{2}, \frac{1}{2})$	$X \rightarrow LXL^\dagger, X = x_\mu\sigma^\mu$
Dirac spinor	$(2, 1) \oplus (1, 2)$	$(\frac{1}{2}, 0) \oplus (0, \frac{1}{2})$	$\Psi \rightarrow \text{diag}(L, (L^\dagger)^{-1})\Psi$

THE BASIC FINITE-DIMENSIONAL REPRESENTATIONS OF THE LORENTZ GROUP,
 WITH $L \in SL(2, \mathbb{C})$

2337 KEY DEFINITIONS

- 2338 • **Isomorphic** ($\cong$): the same structure, e.g. the same commutation
 2339 relations in a suitable basis.
- 2340 • **Universal double cover**, e.g. $SL(2, \mathbb{C}) \rightarrow SO^+(1, 3)$:
 - 2341 – *Cover*: a continuous, locally one-to-one projection from the
 2342 covering group onto the base group that respects
 2343 multiplication.
 - 2344 – *Double*: the map is exactly 2-to-1.
 - 2345 – *Universal*: the covering group is simply connected (every loop
 2346 can be shrunk to a point). $SL(2, \mathbb{C})$ is; $SO^+(1, 3)$ is not, since a
 2347 2π rotation is a loop that cannot be contracted, while a 4π
 2348 rotation can.