



DSC214

Topological Data Analysis

Topic 1: Basics

Instructor: Zhengchao Wan

Goal



- ▶ Fundamental Questions
 - ▶ What is a topological space?
 - ▶ What is a “continuous” way of turning one space to another?
 - ▶ When can we say two spaces are the “same”?

Overview

- ▶ **Fundamental concepts**
 - ▶ Topological space
 - ▶ Continuous maps
 - ▶ Homeomorphisms and homotopies
 - ▶ Manifolds


Overview

▶ Fundamental concepts

- ▶ Topological space
- ▶ Continuous maps
- ▶ Homeomorphisms and homotopies
- ▶ Manifolds


How we mathematically talk about space of interest

Set theory and beyond




Set theory and beyond

- ▶ Given a disk D (without boundary)




Set theory and beyond

- ▶ Given a disk D (without boundary)
- ▶ $x \in D$




Set theory and beyond

- ▶ Given a disk D (without boundary)
- ▶ $x \in D$
- ▶ $y \notin D$




Set theory and beyond

- ▶ Given a disk D (without boundary)
- ▶ $x \in D$
- ▶ $y \notin D$
- ▶ $z \notin D$




Set theory and beyond

- ▶ Given a disk D (without boundary)
- ▶ $x \in D$
- ▶ $y \notin D$
- ▶ $z \notin D$
- ▶ **D contacts both x and z**




Set theory and beyond

- ▶ Given a disk D (without boundary)
- ▶ $x \in D$
- ▶ $y \notin D$
- ▶ $z \notin D$
- ▶ **D contacts both x and z**
- ▶ x and z are in the “**closure**” of D



Why do we care?

- ▶ We want to rigorously define “continuous transformation”
 - ▶ A continuous map shouldn’t tear things apart
 - ▶ If S “contacts” x , under a continuous transformation, we want that $f(S)$ “contacts” $f(x)$




From <https://wgyory.wixsite.com/toolatetopologize/post/post-1>

Why do we care?

- ▶ We want to rigorously define “continuous transformation”
 - ▶ A continuous map shouldn’t tear things apart
 - ▶ If S “contacts” x , under a continuous transformation, we want that $f(S)$ “contacts” $f(x)$
- ▶ We keep track of **ALL** the relations “ S contacts x ” to make the above intuition rigorous!

Topology

- ▶ It turns out that we just need to specify all “open” subsets to keep track of all relations “ S contacts x ” on a given set



Topological space

Definition 1.1 (Topological space) A topological space is a set X endowed with a topological structure (a topology) \mathcal{T} such that the following conditions are satisfied:

1. Both the empty set and X are elements of \mathcal{T} .
2. Any union of arbitrarily many elements of \mathcal{T} is an element of \mathcal{T} .
3. Any intersection of finitely many elements of \mathcal{T} is an element of \mathcal{T} .

$$\begin{array}{ll} 1) & \emptyset, X \in \mathcal{T} \\ 2) & \bigcup_{i \in I} U_i \in \mathcal{T} \\ 3) & U \cap V \in \mathcal{T} \end{array}$$

Topological space

Definition 1.1 (Topological space) A topological space is a set X endowed with a topological structure (a topology) \mathcal{T} such that the following conditions are satisfied:

1. Both the empty set and X are elements of \mathcal{T} .
2. Any union of arbitrarily many elements of \mathcal{T} is an element of \mathcal{T} .
3. Any intersection of finitely many elements of \mathcal{T} is an element of \mathcal{T} .

1)	$\emptyset, X \in \mathcal{T}$
2)	$\bigcup_{i \in I} U_i \in \mathcal{T}$
3)	$U \cap V \in \mathcal{T}$

- ▶ \mathcal{T} is a system of subsets of X . It is called a **topology** on X .

Topological space

Definition 1.1 (Topological space) A topological space is a set X endowed with a topological structure (a topology) \mathcal{T} such that the following conditions are satisfied:

1. Both the empty set and X are elements of \mathcal{T} .
2. Any union of arbitrarily many elements of \mathcal{T} is an element of \mathcal{T} .
3. Any intersection of finitely many elements of \mathcal{T} is an element of \mathcal{T} .

1)	$\emptyset, X \in \mathcal{T}$
2)	$\bigcup_{i \in I} U_i \in \mathcal{T}$
3)	$U \cap V \in \mathcal{T}$

- ▶ \mathcal{T} is a system of subsets of X . It is called a **topology** on X .
- ▶ Examples:

Topological space

Definition 1.1 (Topological space) A topological space is a set X endowed with a topological structure (a topology) \mathcal{T} such that the following conditions are satisfied:

1. Both the empty set and X are elements of \mathcal{T} .
2. Any union of arbitrarily many elements of \mathcal{T} is an element of \mathcal{T} .
3. Any intersection of finitely many elements of \mathcal{T} is an element of \mathcal{T} .

1)	$\emptyset, X \in \mathcal{T}$
2)	$\bigcup_{i \in I} U_i \in \mathcal{T}$
3)	$U \cap V \in \mathcal{T}$

- ▶ \mathcal{T} is a system of subsets of X . It is called a **topology** on X .
- ▶ Examples:
 - ▶ Trivial topology $\{\emptyset, X\}$

Topological space

Definition 1.1 (Topological space) A topological space is a set X endowed with a topological structure (a topology) \mathcal{T} such that the following conditions are satisfied:

1. Both the empty set and X are elements of \mathcal{T} .
2. Any union of arbitrarily many elements of \mathcal{T} is an element of \mathcal{T} .
3. Any intersection of finitely many elements of \mathcal{T} is an element of \mathcal{T} .

1)	$\emptyset, X \in \mathcal{T}$
2)	$\bigcup_{i \in I} U_i \in \mathcal{T}$
3)	$U \cap V \in \mathcal{T}$

- ▶ \mathcal{T} is a system of subsets of X . It is called a **topology** on X .
- ▶ Examples:
 - ▶ Trivial topology $\{\emptyset, X\}$
 - ▶ Discrete topology $2^X = \text{all subsets of } X$

Topological space


Definition 1.1 (Topological space) A topological space is a set X endowed with a topological structure (a topology) \mathcal{T} such that the following conditions are satisfied:

1. Both the empty set and X are elements of \mathcal{T} .
2. Any union of arbitrarily many elements of \mathcal{T} is an element of \mathcal{T} .
3. Any intersection of finitely many elements of \mathcal{T} is an element of \mathcal{T} .

1)	$\emptyset, X \in \mathcal{T}$
2)	$\bigcup_{i \in I} U_i \in \mathcal{T}$
3)	$U \cap V \in \mathcal{T}$

- ▶ \mathcal{T} is a system of subsets of X . It is called a **topology** on X .
- ▶ Examples:
 - ▶ Trivial topology $\{\emptyset, X\}$
 - ▶ Discrete topology $2^X = \text{all subsets of } X$
 - ▶ **Metric space topology**

From <https://wgyory.wixsite.com/toolatetopologize/post/post-1>





Topology provides the formal language to keep track of all the Nearness relation: S contacts x . Hence, we can talk about limits, continuity, etc of functions as well as connectedness, compactness, etc of spaces.

Open / Closed sets

Definition 1.1 (Topological space) A topological space is a set X endowed with a topological structure (a topology) \mathcal{T} such that the following conditions are satisfied:

1. Both the empty set and X are elements of \mathcal{T} .
2. Any union of arbitrarily many elements of \mathcal{T} is an element of \mathcal{T} .
3. Any intersection of finitely many elements of \mathcal{T} is an element of \mathcal{T} .

- ▶ \mathcal{T} is a system of subsets of X . It is called a *topology* on X .
- ▶ Each set $A \in \mathcal{T}$ is called an *open set*

Open / Closed sets

Definition 1.1 (Topological space) A topological space is a set X endowed with a topological structure (a topology) \mathcal{T} such that the following conditions are satisfied:

1. Both the empty set and X are elements of \mathcal{T} .
2. Any union of arbitrarily many elements of \mathcal{T} is an element of \mathcal{T} .
3. Any intersection of finitely many elements of \mathcal{T} is an element of \mathcal{T} .

- ▶ \mathcal{T} is a system of subsets of X . It is called a *topology* on X .
- ▶ Each set $A \in \mathcal{T}$ is called an *open set*
- ▶ A set B is *closed* if its complement is open
 - ▶ i.e., there exists A such that $B = X \setminus A$


Closure, interior, boundary

Closure, interior, boundary

- ▶ Recall a set is closed if its complement is open
- ▶ Given a topological space (X, \mathcal{T}) and a subset $A \subseteq X$:
 - ▶ the *closure* of A , denoted by \bar{A} , is the smallest closed set containing A .
 - ▶ $\bar{A} = \bigcap_{\text{closed } C \supseteq A} C$
 - ▶ its *interior* A^o is the union of all open subsets of A .
 - ▶ the *boundary* of A is $\partial A = \bar{A} \setminus A^o$


Closure, interior, boundary

- ▶ Recall a set is closed if its complement is open
- ▶ Given a topological space (X, \mathcal{T}) and a subset $A \subseteq X$:
 - ▶ the *closure* of A , denoted by \bar{A} , is the smallest closed set containing A .
 - ▶ $\bar{A} = \bigcap_{\text{closed } C \supseteq A} C$
 - ▶ its *interior* A^o is the union of all open subsets of A .
 - ▶ the *boundary* of A is $\partial A = \bar{A} \setminus A^o$



Closure, interior, boundary

- ▶ Recall a set is closed if its complement is open
- ▶ Given a topological space (X, \mathcal{T}) and a subset $A \subseteq X$:
 - ▶ the *closure* of A , denoted by \bar{A} , is the smallest closed set containing A
 - ▶ its *exterior* is \bar{A}^c
 - ▶ “ S contacts x ” can be formally defined as $x \in \bar{S}$
 - ▶ the *boundary* of A is $\bar{A} \setminus A$



Examples in \mathbb{R}

- ▶ Let $A = [1,2)$
 - ▶ $\bar{A} = [1,2]$
 - ▶ $A^o = (1,2)$
 - ▶ $\partial A = \{1,2\}$

- ▶ For any given set X , one can define different topologies on top of that. Some of them can be bizarre, such as the trivial topology
- ▶ The most useful topology in this class is the **metric space topology**

Metric space

Definition 2 (Metric space). A metric space is a pair (X, d) where X is a set and d is a distance function $d : X \times X \rightarrow \mathbb{R}$ satisfying the following properties:

- $d(p, q) = 0$ if and only if $p = q$
- $d(p, q) = d(q, p), \forall p, q \in X;$
- $d(p, q) \leq d(p, r) + d(r, q), \forall p, q, r \in X.$

Metric space

Definition 2 (Metric space). A metric space is a pair (X, d) where X is a set and d is a distance function $d : X \times X \rightarrow \mathbb{R}$ satisfying the following properties:

- $d(p, q) = 0$ if and only if $p = q$
- $d(p, q) = d(q, p), \forall p, q \in X;$
- $d(p, q) \leq d(p, r) + d(r, q), \forall p, q, r \in X.$

► Examples:

- $(\mathbb{R}^k, \|\cdot\|_2)$ k-dimensional Euclidean space, equipped with the standard Euclidean distance
$$d(p, q) = \|p - q\|_2$$
- “Curved” space (manifolds), equipped with geodesic distance
 - e.g, the surface of earth.
- Space can also be discrete, as very often in data analysis
 - (P, d) : a set of points with pairwise distance (or similarity) given.
 - or graphs, equipped with shortest path metric.

Metric space topology

- ▶ Open ball:

- ▶ $B_o(c, r) = \{x \in X \mid d(c, x) < r\}$

Definition 3 (Metric space topology). *Given a metric space X , all metric balls $\{B_o(c, r) \mid c \in \mathbb{T} \text{ and } 0 < r \leq \infty\}$ and their union constituting the open sets define a topology on X .*

- ▶ Exercise: prove that this is a topology on X

Metric space topology

- ▶ Open ball:

- ▶ $B_o(c, r) = \{x \in X \mid d(c, x) < r\}$

Definition 3 (Metric space topology). *Given a metric space X , all metric balls $\{B_o(c, r) \mid c \in \mathbb{T} \text{ and } 0 < r \leq \infty\}$ and their union constituting the open sets define a topology on X .*

- ▶ Exercise: prove that this is a topology on X
- ▶ The set of metric balls is called a *basis* for this topology on X
 - ▶ it generates all open sets in this topology

Metric space topology

- ▶ Open ball:

- ▶ $B_o(c, r) = \{x \in X \mid d(c, x) < r\}$

Definition 3 (Metric space topology). *Given a metric space X , all metric balls $\{B_o(c, r) \mid c \in \mathbb{T} \text{ and } 0 < r \leq \infty\}$ and their union constituting the open sets define a topology on X .*

- ▶ Exercise: prove that this is a topology on X
- ▶ The set of metric balls is called a *basis* for this topology on X
 - ▶ it generates all open sets in this topology
- ▶ In general, when we refer to a common metric space, say Euclidean space, we refer to this metric space topology induced by standard metric.

Metric space topology on \mathbb{R}

Metric space topology on \mathbb{R}

- ▶ Each open ball is an open interval $(c - r, c + r)$

Metric space topology on \mathbb{R}

- ▶ Each open ball is an open interval $(c - r, c + r)$
- ▶ Each open set is a union of arbitrarily many open intervals (by definition)

Metric space topology on \mathbb{R}

- ▶ Each open ball is an open interval $(c - r, c + r)$
- ▶ Each open set is a union of arbitrarily many open intervals (by definition)
- ▶ Each open set is a countable union of open intervals

Metric space topology on \mathbb{R}

- ▶ Each open ball is an open interval $(c - r, c + r)$
- ▶ Each open set is a union of arbitrarily many open intervals (by definition)
- ▶ Each open set is a **countable** union of open intervals

Metric space topology on \mathbb{R}

- ▶ Each open ball is an open interval $(c - r, c + r)$
- ▶ Each open set is a union of arbitrarily many open intervals (by definition)
- ▶ Each open set is a **countable** union of open intervals

Exercise: why?

Subspace topology

- ▶ A topological space (X, \mathcal{T}) , say the Euclidean space
- ▶ Given a subset $Y \subseteq X$, the subspace topology (Y, \mathcal{T}_Y) , (inherited from (X, \mathcal{T})), is such that \mathcal{T}_Y consists of intersection between open sets in \mathcal{T} and Y .
- ▶ Common subspaces of Euclidean space
 - ▶ Euclidean d-ball: $\mathbb{B}^d = \{x \in \mathbb{R}^d \mid \|x\| \leq 1\}$
 - ▶ Open Euclidean d-ball: $\mathbb{B}_o^d = \{x \in \mathbb{R}^d \mid \|x\| < 1\}$
 - ▶ Euclidean d-sphere: $\mathbb{S}^d = \{x \in \mathbb{R}^{d+1} \mid \|x\| = 1\}$
 - ▶ Euclidean half-space: $\mathbb{H}^d = \{x \in \mathbb{R}^d \mid x_d \geq 0\}$

Subspace topology

- ▶ A topological space (X, \mathcal{T}) , say the Euclidean space
- ▶ Given a subset $Y \subseteq X$, the subspace topology (Y, \mathcal{T}_Y) , (inherited from (X, \mathcal{T})), is such that \mathcal{T}_Y consists of intersection between open sets in \mathcal{T} and Y .
- ▶ Example: $X = \mathbb{R}$ and $Y = [1,2]$. Then, $(1.5,2] = (1.5,3) \cap [1,2]$ is an open set in subspace topology.



Subspace topology

- ▶ A topological space (X, \mathcal{T}) , say the Euclidean space
- ▶ Given a subset $Y \subseteq X$, the subspace topology (Y, \mathcal{T}_Y) , (inherited from (X, \mathcal{T})), is such that \mathcal{T}_Y consists of intersection between open sets in \mathcal{T} and Y .
- ▶ Example: $X = \mathbb{R}$ and $Y = [1,2]$. Then, $(1.5,2] = (1.5,3) \cap [1,2]$ is an open set in subspace topology.



Connectivity

- ▶ Open sets determine connectivity.

Connectivity

- ▶ Open sets determine connectivity.
- ▶ A topological space (X, \mathcal{T}) is *disconnected* if there are two disjoint non-empty open sets $A, B \in \mathcal{T}$ so that $X = A \cup B$.

Connectivity


- ▶ Open sets determine connectivity.
- ▶ A topological space (X, \mathcal{T}) is *disconnected* if there are two disjoint non-empty open sets $A, B \in \mathcal{T}$ so that $X = A \cup B$.
- ▶ A topological space is *connected* if it is not disconnected.

Connectivity

- ▶ Open sets determine connectivity.
- ▶ A topological space (X, \mathcal{T}) is *disconnected* if there are two disjoint non-empty open sets $A, B \in \mathcal{T}$ so that $X = A \cup B$.
- ▶ A topological space is *connected* if it is not disconnected.
- ▶ Any **maximal** connected subsets of X is called a **connected component**.



Connectivity

- ▶ Open sets determine connectivity.
- ▶ A topological space (X, \mathcal{T}) is *disconnected* if there are two disjoint non-empty open sets $A, B \in \mathcal{T}$ so that $X = A \cup B$.
- ▶ A topological space is *connected* if it is not disconnected.
- ▶ Any **maximal** connected subsets of X is called a **connected component**.



Connectivity

- ▶ Open sets determine connectivity.
- ▶ A topological space (X, \mathcal{T}) is *disconnected* if there are two disjoint non-empty open sets $A, B \in \mathcal{T}$ so that $X = A \cup B$.
- ▶ A topological space is *connected* if it is not disconnected.
- ▶ Any *maximal* connected subsets of X is called a **connected component**.



Compactness

- ▶ This generalizes the notion of **closed** and **bounded** sets in Euclidean space
- ▶ Open cover: $\mathcal{U} = \{U_\alpha\}_{\alpha \in A}$ is an open cover for (X, \mathcal{T}) if $U_\alpha \in \mathcal{T}$ and $X = \bigcup_{\alpha \in A} U_\alpha$
- ▶ (X, \mathcal{T}) is called **compact** if for any open cover $\mathcal{U} = \{U_\alpha\}_{\alpha \in A}$ there exists a finite subcover, i.e., a finite set $A' \subseteq A$ such that $X = \bigcup_{\alpha \in A'} U_\alpha$
- ▶ Example: $(0,1)$ is not compact but $[0,1]$ is compact

Check-in: Where are we?

- ▶ Fundamental concepts
 - ▶ Topological space
 - ▶ Continuous maps
 - ▶ Homeomorphisms and homotopies
 - ▶ Manifolds

Check-in: Where are we?

▶ Fundamental concepts

- ▶ Topological space
- ▶ Continuous maps
- ▶ Homeomorphisms and homotopies
- ▶ Manifolds

How we mathematically talk about space of interest


Check-in: Where are we?

▶ Fundamental concepts

- ▶ Topological space → How we mathematically talk about space of interest
- ▶ Continuous maps → Now we need ways to connect different spaces!
- ▶ Homeomorphisms and homotopies
- ▶ Manifolds

Recall


- ▶ We want to rigorously define “continuous transformation”
 - ▶ A continuous map shouldn’t tear things apart
 - ▶ If S “contacts” x , under a continuous transformation, we want that $f(S)$ “contacts” $f(x)$



From <https://wgyory.wixsite.com/toolatetopologize/post/post-1>

Continuous function

- ▶ A function $f : X \rightarrow Y$ between two topological spaces is called **continuous** if for any subset $S \subset X$ we have that
 - ▶ $f(\bar{S}) \subset \overline{f(S)}$
- ▶ A formal way describing “If S contacts x , then $f(S)$ contacts $f(x)$ ”




From <https://wgyory.wixsite.com/toolatetopologize/post/post-1>

Continuous function

- ▶ A function $f : X \rightarrow Y$ between two topological spaces is called **continuous** if for any subset $S \subset X$ we have that
 - ▶ $f(\bar{S}) \subset \overline{f(S)}$
- ▶ A formal way describing “If S contacts x , then $f(S)$ contacts $f(x)$ ”

1. Closure of image contains image of closure
2. Continuous map does not tear things apart



From <https://wgyory.wixsite.com/toolatetopologize/post/post-1>


Exercise

- ▶ Please try to prove the notion of continuity in calculus is compatible with the new definition of continuity
 - ▶ $\lim_{x \rightarrow x_0} f(x) = f(x_0)$ for all x_0



Check-in: Where are we?

▶ Fundamental concepts

- ▶ Topological space → How we mathematically talk about space of interest
- ▶ Continuous maps → Now we need ways to connect different spaces!
- ▶ Homeomorphisms and homotopies → Describe relations of spaces
- ▶ Manifolds



\equiv



Homeomorphism = homoios + morphē = Similar shapes

Definition 5 (Homeomorphism) *Given two topological spaces X and Y , a homeomorphism between them is a map $h : X \rightarrow Y$ such that h is bijection and the inverse of h is also continuous.*

Two topological spaces are X and Y are homeomorphic, denoted by $X \cong Y$, if there is a homeomorphism between them.

Homeomorphism = homoios + morphē = Similar shapes

Definition 5 (Homeomorphism) *Given two topological spaces X and Y , a homeomorphism between them is a map $h : X \rightarrow Y$ such that h is bijection and the inverse of h is also continuous.*

Two topological spaces are X and Y are homeomorphic, denoted by $X \cong Y$, if there is a homeomorphism between them.

- ▶ Homeomorphic spaces are called *topologically equivalent*
 - ▶ Note that equivalent relations are transitive.
 - ▶ $X \cong Y$ and $Y \cong Z$ implies $X \cong Z$

Homeomorphism = homoios + morphē = Similar shapes

Definition 5 (Homeomorphism) *Given two topological spaces X and Y , a homeomorphism between them is a map $h : X \rightarrow Y$ such that h is bijection and the inverse of h is also continuous.*

Two topological spaces are X and Y are homeomorphic, denoted by $X \cong Y$, if there is a homeomorphism between them.

- ▶ Homeomorphic spaces are called *topologically equivalent*
 - ▶ Note that equivalent relations are transitive.
 - ▶ $X \cong Y$ and $Y \cong Z$ implies $X \cong Z$
- ▶ Layman's terms: two spaces are homeomorphic if one can continuously deform (stretch, compress) into the other without ever breaking or stitching them
 - ▶ Caveat: not always true

Homeomorphism = homoios + morphē = Similar shapes


Definition 5 (Homeomorphism) *Given two topological spaces X and Y , a homeomorphism between them is a map $h : X \rightarrow Y$ such that h is bijection and the inverse of h is also continuous.*

Two topological spaces are X and Y are homeomorphic, denoted by $X \cong Y$, if there is a homeomorphism between them.

- ▶ Homeomorphic spaces are called *topologically equivalent*
 - ▶ Note that equivalent relations are transitive.
 - ▶ $X \cong Y$ and $Y \cong Z$ implies $X \cong Z$
- ▶ Layman's terms: two spaces are homeomorphic if one can continuously deform (stretch, compress) into the other without ever breaking or stitching them
 - ▶ Caveat: not always true
- ▶ Homeomorphism preserves all topological quantities: number of connected components, number of holes, voids, etc.


Continuous bijective map may have discontinuous inverse

- ▶ $[0, 2\pi) \rightarrow \mathbb{S}^1$ by $\theta \mapsto e^{i\theta}$





Examples

Examples



Examples



Non-examples

Definition 5 (Homeomorphism) *Given two topological spaces X and Y , a homeomorphism between them is a map $h : X \rightarrow Y$ such that h is bijection and the inverse of h is also continuous.*

Two topological spaces are X and Y are homeomorphic, denoted by $X \cong Y$, if there is a homeomorphism between them.

Non-examples

Definition 5 (Homeomorphism) *Given two topological spaces X and Y , a homeomorphism between them is a map $h : X \rightarrow Y$ such that h is bijection and the inverse of h is also continuous.*

Two topological spaces are X and Y are homeomorphic, denoted by $X \cong Y$, if there is a homeomorphism between them.

- ▶ A trick: remove one point and check connected components
 - ▶ Y and I are not homeomorphic; X and Y are not homeomorphic
 - ▶ \mathbb{R} and \mathbb{R}^2 are not homeomorphic
 - ▶ What about \mathbb{R}^2 and \mathbb{R}^3 ?
- ▶ In general, hard to decide whether two spaces are homeomorphic or not!

More Examples and Non Examples

More Examples and Non Examples

- ▶ The Euclidean space \mathbb{R}^d is homeomorphic to any open ball $\mathbb{B}_o(c, r)$

More Examples and Non Examples

- ▶ The Euclidean space \mathbb{R}^d is homeomorphic to any open ball $\mathbb{B}_o(c, r)$
- ▶ Try to construct the homeomorphism by yourself

More Examples and Non Examples

- ▶ The Euclidean space \mathbb{R}^d is homeomorphic to any open ball $\mathbb{B}_o(c, r)$
- ▶ Try to construct the homeomorphism by yourself

More Examples and Non Examples

- ▶ The Euclidean space \mathbb{R}^d is homeomorphic to any open ball $\mathbb{B}_o(c, r)$
 - ▶ Try to construct the homeomorphism by yourself
- ▶ $[0,1]$, $(0,1]$, $(0,1)$

More Examples and Non Examples

- ▶ The Euclidean space \mathbb{R}^d is homeomorphic to any open ball $\mathbb{B}_o(c, r)$
 - ▶ Try to construct the homeomorphism by yourself
- ▶ $[0,1]$, $(0,1]$, $(0,1)$



More Examples and Non Examples

- ▶ The Euclidean space \mathbb{R}^d is homeomorphic to any open ball $\mathbb{B}_o(c, r)$
 - ▶ Try to construct the homeomorphism by yourself
- ▶ $[0,1]$, $(0,1]$, $(0,1)$
- ▶ \mathbb{R}^d is not homeomorphic to the half space $\mathbb{H}^d = \{(x_1, \dots, x_d) : x_1 \geq 0\}$

More Examples and Non Examples


- ▶ The Euclidean space \mathbb{R}^d is homeomorphic to any open ball $\mathbb{B}_o(c, r)$
 - ▶ Try to construct the homeomorphism by yourself
- ▶ $[0,1]$, $(0,1]$, $(0,1)$
- ▶ \mathbb{R}^d is not homeomorphic to the half space $\mathbb{H}^d = \{(x_1, \dots, x_d) : x_1 \geq 0\}$
 - ▶ We will see why after knowing homotopy

Another level of similarity





- ▶ They are not homeomorphic (why?)
- ▶ But they look very similar

A closer look





A closer look





Homotopy and Deformation Retraction

Definition 6 (Homotopy) First, two $f, g : X \rightarrow Y$ are homotopic if there is a continuous map $H : X \times [0, 1] \rightarrow Y$ such that $H_0 = f$ and $H_1 = g$. This map H is called a homotopy connecting f and g . If H exists, we say f and g are homotopic, denoted by $f \simeq g$



Homotopy and Deformation Retraction

Definition 6 (Homotopy) First, two $f, g : X \rightarrow Y$ are homotopic if there is a continuous map $H : X \times [0, 1] \rightarrow Y$ such that $H_0 = f$ and $H_1 = g$. This map H is called a homotopy connecting f and g . If H exists, we say f and g are homotopic, denoted by $f \simeq g$




Definition 7 (Deformation retraction) Let $A \subseteq X$ be a subspace of topological space X . A retraction (map) r is a continuous map $r : X \rightarrow A$ such that $r(x) = x$ for any $x \in A$.

We say that $A \subseteq X$ is a deformation retract of X if there is a retraction r that is homotopic to the identity map in X . This retraction map is called a deformation retraction.


Equivalently, a continuous map $R : X \times [0, 1] \rightarrow X$ is a deformation retraction of X onto A if for every $x \in X$ and $a \in A$, $R(x, 0) = x$; $R(x, 1) \in A$ and $R(a, 1) = a$.

Examples of Deformation Retraction



A weaker notion of similarity: Homotopy equivalent



- When X, Y are homeomorphic, there exists continuous functions $f: X \rightarrow Y, g: Y \rightarrow X$ such that $f \circ g = Id_Y$ and $g \circ f = Id_X$



- We say X, Y are **homotopy equivalent**, if there exists continuous functions $f: X \rightarrow Y, g: Y \rightarrow X$ such that $f \circ g \simeq Id_Y$ and $g \circ f \simeq Id_X$

A weaker notion of similarity: Homotopy equivalent

- ▶ Homeomorphism allows stretching and shrinking



- ▶ Homotopy allows stretching, shrinking and **crushing/collapsing**

A weaker notion: Homotopy equivalent

Definition 6 (Homotopy) First, two $f, g : X \rightarrow Y$ are homotopic if there is a continuous map $H : X \times [0, 1] \rightarrow Y$ such that $H_0 = f$ and $H_1 = g$. This map H is called a homotopy connecting f and g .

Definition 7 (Homotopy equivalence) Two spaces X and Y are homotopy equivalent if there is a continuous mapping $f : X \rightarrow Y$ and $g : Y \rightarrow X$ such that $f \circ g$ is homotopic to identity in Y and $g \circ f$ is homotopic to identity in X .

- ▶ **Theorem:**
 - ▶ Two homeomorphic spaces X and Y are also homotopy equivalent. But the inverse may not hold.

A weaker notion: Homotopy equivalent

Definition 6 (Homotopy) First, two $f, g : X \rightarrow Y$ are homotopic if there is a continuous map $H : X \times [0, 1] \rightarrow Y$ such that $H_0 = f$ and $H_1 = g$. This map H is called a homotopy connecting f and g .

Definition 7 (Homotopy equivalence) Two spaces X and Y are homotopy equivalent if there is a continuous mapping $f : X \rightarrow Y$ and $g : Y \rightarrow X$ such that $f \circ g$ is homotopic to identity in Y and $g \circ f$ is homotopic to identity in X .

- ▶ **Theorem:**
 - ▶ Two homeomorphic spaces X and Y are also homotopy equivalent. But the inverse may not hold.
- ▶ Homotopy equivalent relation is transitive.

- In general, hard to establish homotopy equivalent relation as well




Definition 7 (Deformation retraction) Let $A \subseteq X$ be a subspace of topological space X . A retraction (map) r is a continuous map $r : X \rightarrow A$ such that $r(x) = x$ for any $x \in A$.

We say that $A \subseteq X$ is a deformation retract of X if there is a retraction r that is homotopic to the identity map in X . This retraction map is called a deformation retraction.

Equivalently, a continuous map $R : X \times [0, 1] \rightarrow X$ is a deformation retraction of X onto A if for every $x \in X$ and $a \in A$, $R(x, 0) = x$; $R(x, 1) \in A$ and $R(a, 1) = a$.


Theorem:

- If Y is a deformation retract of X , then X and Y are homotopy equivalent.




Examples of homotopy equivalence

- ▶ X and Y are homotopy equivalent but not homeomorphic



- ▶ A disk and a point
- ▶ A tree and a point
- ▶ A punctured plane and a circle




$$R : [0,1] \times \mathbb{R}^2 \setminus 0 \rightarrow \mathbb{R}^2 \setminus 0$$




$$R(t, x) = (1 - t)x + t \frac{x}{\|x\|}$$

Examples of homotopy equivalence

- ▶ X and Y are homotopy equivalent but not homeomorphic



- ▶ A disk and a point
- ▶ A tree and a point
- ▶ A punctured plane and a circle



$$R : [0,1] \times \mathbb{R}^2 \setminus 0 \rightarrow \mathbb{R}^2 \setminus 0$$

$$R(t, x) = (1 - t)x + t \frac{x}{\|x\|}$$

Examples

- ▶ A punctured hollow torus \approx ?



Examples

- ▶ A punctured hollow torus \approx ?



Non Examples

Non Examples

- ▶ The Euclidean space \mathbb{R}^n is homotopy equivalent to a point

Non Examples

- ▶ The Euclidean space \mathbb{R}^n is homotopy equivalent to a point
- ▶ But a sphere \mathbb{S}^n is not homotopy equivalent to a point

Non Examples

- ▶ The Euclidean space \mathbb{R}^n is homotopy equivalent to a point
- ▶ But a sphere \mathbb{S}^n is not homotopy equivalent to a point
 - ▶ Surprisingly we need advanced machinery to prove this

Non Examples

- ▶ The Euclidean space \mathbb{R}^n is homotopy equivalent to a point
- ▶ But a sphere \mathbb{S}^n is not homotopy equivalent to a point
 - ▶ Surprisingly we need advanced machinery to prove this
 - ▶ Take this for granted at this moment

Non Examples

- ▶ The Euclidean space \mathbb{R}^n is homotopy equivalent to a point
- ▶ But a sphere \mathbb{S}^n is not homotopy equivalent to a point
 - ▶ Surprisingly we need advanced machinery to prove this
 - ▶ Take this for granted at this moment
- ▶ Are \mathbb{S}^n and \mathbb{S}^m homotopy equivalent?

Non Examples

- ▶ The Euclidean space \mathbb{R}^n is homotopy equivalent to a point
- ▶ But a sphere \mathbb{S}^n is not homotopy equivalent to a point
 - ▶ Surprisingly we need advanced machinery to prove this
 - ▶ Take this for granted at this moment
- ▶ Are \mathbb{S}^n and \mathbb{S}^m homotopy equivalent?
- ▶ Can we use computer to determine whether two topological spaces are homotopy equivalent or not?

Non Examples



- ▶ The Euclidean space \mathbb{R}^n is homotopy equivalent to a point
- ▶ But a sphere \mathbb{S}^n is not homotopy equivalent to a point
 - ▶ Surprisingly we need advanced machinery to prove this
 - ▶ Take this for granted at this moment
- ▶ Are \mathbb{S}^n and \mathbb{S}^m homotopy equivalent?

Both homeomorphism and homotopy equivalence can be hard to detect, and not computationally friendly in general. Later we will focus on homology, which is much easier to compute.

Here is what I promised

- ▶ \mathbb{R}^d is not homeomorphic to the half space $\mathbb{H}^d = \{(x_1, \dots, x_d) : x_1 \geq 0\}$


Summary



- ▶ Fundamental Questions
 - ▶ What is a topological space?
 - ▶ What is a “continuous” way of turning one space to another?
 - ▶ When can we say two spaces are the “same”?


Check-in: Where are we?

▶ Fundamental concepts




Why manifolds?

- ▶ We like Euclidean spaces. Why manifolds?



Why manifolds?

- ▶ We like Euclidean spaces. Why manifolds?
 - ▶ We live in non-Euclidean space



Why manifolds?


- ▶ We like Euclidean spaces. Why manifolds?

Why manifolds?

- ▶ We like Euclidean spaces. Why manifolds?
 - ▶ Data and the space where data are sampled from often live in a subspace within the ambient space they are embedded in
 - ▶ Ambient space:
 - the space where data or the object/space of interest are embedded in.
 - ▶ Intrinsic space:
 - the space where data are sampled from (**manifold hypothesis**)


Why manifolds?

- ▶ We like Euclidean spaces. Why manifolds?
 - ▶ Data and the space where data are sampled from often live in a subspace within the ambient space they are embedded in
 - ▶ Ambient space:
 - the space where data or the object/space of interest are embedded in.
 - ▶ Intrinsic space:
 - the space where data are sampled from (**manifold hypothesis**)




Why manifolds?

- ▶ We like Euclidean spaces. Why manifolds?
 - ▶ Data and the space where data are sampled from often live in a subspace within the ambient space they are embedded in
 - ▶ Ambient space:
 - the space where data or the object/space of interest are embedded in.
 - ▶ Intrinsic space:
 - the space where data are sampled from (**manifold hypothesis**)



7210414959
0690159734
9665407401
3134727121
1742351244




Why manifolds?

- ▶ We like Euclidean spaces. Why manifolds?
 - ▶ Data and the space where data are sampled from often live in a subspace within the ambient space they are embedded in
 - ▶ Ambient space:
 - the space where data or the object/space of interest are embedded in.
 - ▶ Intrinsic space:
 - the space where data are sampled from
 - ▶ The intrinsic space may not be a linear subspace of the ambient space
 - ▶ e.g., the surface of a bunny in \mathbb{R}^3

Why manifolds?

- ▶ We like Euclidean spaces. Why manifolds?
 - ▶ Data and the space where data are sampled from often live in a subspace within the ambient space they are embedded in
 - ▶ Ambient space:
 - the space where data or the object/space of interest are embedded in.
 - ▶ Intrinsic space:
 - the space where data are sampled from
 - ▶ The intrinsic space may not be a linear subspace of the ambient space
 - ▶ e.g., the surface of a bunny in \mathbb{R}^3



What are manifolds

- ▶ Non-linear, yet still well-behaved spaces!
- ▶ In particular, locally it behaves like a linear space.

What are manifolds


- ▶ Non-linear, yet still well-behaved spaces!
- ▶ In particular, locally it behaves like a linear space.

Definition 8 (Manifold). A topological space M is a *m-manifold*, or simply *manifold*, if every point $x \in M$ has a neighborhood homeomorphic to \mathbb{B}_o^m or \mathbb{H}^m . The *dimension* of M is m .

What are manifolds

- ▶ Non-linear, yet still well-behaved spaces!
- ▶ In particular, locally it behaves like a linear space.

Definition 8 (Manifold). A topological space M is a *m-manifold*, or simply *manifold*, if every point $x \in M$ has a neighborhood homeomorphic to \mathbb{B}_o^m or \mathbb{H}^m . The *dimension* of M is m .



What are manifolds

- ▶ Non-linear, yet still well-behaved spaces!
- ▶ In particular, locally it behaves like a linear space.

What are manifolds

- ▶ Non-linear, yet still well-behaved spaces!
- ▶ In particular, locally it behaves like a linear space.

Definition 8 (Manifold). A topological space M is a *m-manifold*, or simply *manifold*, if every point $x \in M$ has a neighborhood homeomorphic to \mathbb{B}_o^m or \mathbb{H}^m . The *dimension* of M is m .

What are manifolds

- ▶ Non-linear, yet still well-behaved spaces!
- ▶ In particular, locally it behaves like a linear space.

Definition 8 (Manifold). A topological space M is a *m-manifold*, or simply *manifold*, if every point $x \in M$ has a neighborhood homeomorphic to \mathbb{B}_o^m or \mathbb{H}^m . The *dimension* of M is m .

- ▶ Interior of M :

What are manifolds

- ▶ Non-linear, yet still well-behaved spaces!
- ▶ In particular, locally it behaves like a linear space.

Definition 8 (Manifold). A topological space M is a *m-manifold*, or simply *manifold*, if every point $x \in M$ has a neighborhood homeomorphic to \mathbb{B}_o^m or \mathbb{H}^m . The *dimension* of M is m .

- ▶ Interior of M :
 - ▶ those points with a neighborhood homeomorphic to \mathbb{B}_o^d

What are manifolds

- ▶ Non-linear, yet still well-behaved spaces!
- ▶ In particular, locally it behaves like a linear space.

Definition 8 (Manifold). A topological space M is a *m-manifold*, or simply *manifold*, if every point $x \in M$ has a neighborhood homeomorphic to \mathbb{B}_o^m or \mathbb{H}^m . The *dimension* of M is m .

- ▶ Interior of M :
 - ▶ those points with a neighborhood homeomorphic to \mathbb{B}_o^d
- ▶ Otherwise, boundary of M

What are manifolds

- ▶ Non-linear, yet still well-behaved spaces!
- ▶ In particular, locally it behaves like a linear space.

Definition 8 (Manifold). A topological space M is a *m-manifold*, or simply *manifold*, if every point $x \in M$ has a neighborhood homeomorphic to \mathbb{B}_o^m or \mathbb{H}^m . The *dimension* of M is *m*.

- ▶ Interior of M :
 - ▶ those points with a neighborhood homeomorphic to \mathbb{B}_o^d
- ▶ Otherwise, boundary of M
- ▶ When referring to a manifold, usually it comes with no boundary, i.e., all neighborhood homeomorphic to the open ball instead of the half space

What are manifolds

- ▶ Non-linear, yet still well-behaved spaces!
- ▶ In particular, locally it behaves like a linear space.

Definition 8 (Manifold). A topological space M is a *m-manifold*, or simply *manifold*, if every point $x \in M$ has a neighborhood homeomorphic to \mathbb{B}_o^m or \mathbb{H}^m . The *dimension* of M is *m*.

- ▶ Interior of M :
 - ▶ those points with a neighborhood homeomorphic to \mathbb{B}_o^d
- ▶ Otherwise, boundary of M
- ▶ When referring to a manifold, usually it comes with no boundary, i.e., all neighborhood homeomorphic to the open ball instead of the half space
- ▶ Will emphasize “manifold with boundary” when necessary

What are manifolds


- ▶ Non-linear, yet still well-behaved spaces!
- ▶ In particular, locally it behaves like a linear space.

Definition 8 (Manifold). A topological space M is a *m-manifold*, or simply *manifold*, if every point $x \in M$ has a neighborhood homeomorphic to \mathbb{B}_o^m or \mathbb{H}^m . The *dimension* of M is *m*.


- ▶ Interior of M :
 - ▶ those points with a neighborhood homeomorphic to \mathbb{B}_o^d
- ▶ Otherwise, boundary of M
- ▶ When referring to a manifold, usually it comes with no boundary, i.e., all neighborhood homeomorphic to the open ball instead of the half space
- ▶ Will emphasize “manifold with boundary” when necessary

1-dim manifold


1-dim manifold



1-dim manifold




1-dim manifold




- ▶ These are everything!
- ▶ Any connected 1-dim manifold (with or without boundary) is homeomorphic to one of the above

A non-example




2-dim manifold

Sphere




Klein bottle




2-dim manifold


Sphere




Klein bottle




Möbius strip




Charts of the sphere



=





U



Homeomorphism to \mathbb{R}^d

Stereographic projection (pic from wiki)





Is this a 2-d manifold?




Classification of 2d manifolds?

- ▶ A 2d manifold without boundary is also called a **surface**
- ▶ All connected compact surfaces can be classified into a specific class of surfaces constructed from polygons




Classification of 2d manifolds?


- ▶ A 2d manifold without boundary is also called a **surface**
- ▶ All connected compact surfaces can be classified into a specific class of surfaces constructed from polygons




sphere



real projective plane




torus




Klein bottle


Visualization of double Torus




Visualization of double Torus




Visualization of projective plane




Visualization of projective plane




Visualization of Klein bottle




Visualization of Klein bottle



Projective plane = Möbius strip + a disk




Projective plane = Möbius strip + a disk






Connected sum operation

- Given two compact surfaces M and S , the connected sum $M \# S$ intuitively “merge” the two by cutting off a small disk (cap) from each surface, and glue the remaining of the two surfaces along the boundary after the cutting.




Classification of compact surfaces


Theorem 2 (Classification Theorem) *The two infinite families \mathbb{S} , \mathbb{T} , $\mathbb{T}\#\mathbb{T}, \dots$, and \mathbb{P} , $\mathbb{P}\#\mathbb{P}, \dots$, exhaust the family of compact 2-manifold without boundary (upto homeomorphism). The first family of surfaces are all orientable; while the second family are all non-orientable. Furthermore, no two surfaces in these sequences are homeomorphic.*



How does $\mathbb{P} \# \mathbb{P}$ become the Klein bottle?



How does $\mathbb{P} \# \mathbb{P}$ become the Klein bottle?



Classification of compact surfaces

Theorem 2 (Classification Theorem) *The two infinite families \mathbb{S} , \mathbb{T} , $\mathbb{T} \# \mathbb{T}, \dots$, and \mathbb{P} , $\mathbb{P} \# \mathbb{P}, \dots$, exhaust the family of compact 2-manifold without boundary (upto homeomorphism). The first family of surfaces are all orientable; while the second family are all non-orientable. Furthermore, no two surfaces in these sequences are homeomorphic.*

- ▶ Intuitively
 - ▶ all orientable surfaces without boundaries can be generated by gluing handles to a sphere
- ▶ The number of \mathbb{T} or \mathbb{P} needed is called the genus g of the surface M
 - ▶ Sphere has genus 0, torus has genus 1, double-torus has genus 2.
- ▶ Hence the genus of a surface completely decides its topology upto homeomorphism
 - ▶ Any two compact surfaces with the same genus are homeomorphic

3d and beyond?

- ▶ Super complicated
- ▶ Way way way beyond the scope of this class
- ▶ Fields Medal level of mathematics

FIN