

Curvature and Vector Fields

Lecture Notes of Loring W. Tu [1]

Zhenhao Huang

Last updated: September 27, 2025

Contents

Lecture 1–Riemannian manifold	1
1.1 Inner Products on a Vector Space	1
1.2 Representations of Inner Products by Symmetric Matrices	2
1.3 Riemannian Metrics	3
1.4 Existence of a Riemannian Metric	3
1.5 Problems	3

❖ Lecture 1 (1/9)

1.1 Inner Products on a Vector Space

The **Euclidean inner product** on \mathbb{R}^n is defined by

$$\langle u, v \rangle = \sum_1^n u^i v^i, \quad (1)$$

and the length of a vector is

$$\|u\| = \sqrt{\langle u, u \rangle}, \quad (2)$$

the **angle** θ between two vectors is

$$\cos \theta = \frac{\langle u, v \rangle}{\|u\| \|v\|}, \quad (3)$$

the **arc length** of a curve $c(t) \in \mathbb{R}^n, a \leq t \leq b$ is

$$s = \int_a^b \|c'(t)\| dt \quad (4)$$

Definition 1.1. An inner product in a real vector space V is a postive-definite, bilinear and symmetric map: $\langle \cdot, \cdot \rangle : V \times V \rightarrow \mathbb{R}$ so that for $u, v, w \in V$ and $a, b \in \mathbb{R}$, satisfies

- (i) **Postive-definiteness** $\langle v, v \rangle = 0$ iff. $v = 0$
- (ii) **Symmetry** $\langle u, v \rangle = \langle v, u \rangle$
- (iii) **Bilinear** $\langle au + bv, w \rangle = a\langle u, w \rangle + b\langle v, w \rangle$

Proposition 1.2. If W is a subspace of V , then the restriction

$$\langle \cdot, \cdot \rangle_W := \langle \cdot, \cdot \rangle|_{W \times W} : W \times W \rightarrow \mathbb{R}, \quad (5)$$

of an inner product $\langle \cdot, \cdot \rangle$ on V is also an innver prodcut.

Proof. The subspace construction preserves the properites in Definition 1.1. \square

Proposition 1.3. The **nonnegative linear combination** of inner products $\langle \cdot, \cdot \rangle_i$ on V : $\langle \cdot, \cdot \rangle := \sum_{i=1}^r a_i \langle \cdot, \cdot \rangle_i, a_i \geq 0$ is again an inner product on V .

Proof. The **nonnegativity** of a_i preserves condition (i) in Definition 1.1, the linearity makes condition (ii), (iii) hold. \square

1.2 Representations of Inner Products by Symmetric Matrices

Let e_1, \dots, e_n be the basis of vector space V , each vector $x \in V$ can be represented as a column vector

$$x = \sum_{i=1}^n x^i e_i \leftrightarrow \mathbf{x} = \begin{bmatrix} x^1 \\ \vdots \\ x^n \end{bmatrix}. \quad (6)$$

Let A be an $n \times n$ matrix whose entries $a_{ij} = \langle e_i, e_j \rangle$, the matrix form of an inner product on V is

$$\langle x, y \rangle = \sum_{ij} x^i y^j \langle e_i, e_j \rangle = \mathbf{x}^\top A \mathbf{y}. \quad (7)$$

We find that, once a basis of V is chosen, the inner product on V determines a postive-definite symmetric matrix. Conversely, an $n \times n$ postive-definite symmetric matrix with a basis of V determines an inner product on V

It follows that there is an one-to-one correspondence

$$\left\{ \begin{array}{c} \text{inner product on a } n\text{-dimensional} \\ \text{vector space} \end{array} \right\} \leftrightarrow \left\{ \begin{array}{c} \text{An } n \times n \text{ postive-definite} \\ \text{symmetric matrix} \end{array} \right\}. \quad (8)$$

Let a basis of dual space $V^\vee := \text{Hom}(V, \mathbb{R})$ be $\alpha^1, \dots, \alpha^n$ w.r.t. the basis e_1, \dots, e_n of V , an inner product \langle, \rangle of $x, y \in V$ is

$$\begin{aligned}\langle x, y \rangle &= \sum_{i,j} a_{ij} x^i y^j = \sum_{i,j} a_{ij} \alpha^i(x) \alpha^j(y) \\ &= \sum_{i,j} a_{ij} \alpha^i \otimes \alpha^j(x, y)\end{aligned}$$

In terms of tensor product, an inner product on V may be written as

$$\langle, \rangle = \sum_{ij} a_{ij} \alpha^i \otimes \alpha^j \tag{9}$$

1.3 Riemannian Metrics

1.4 Existence of a Riemannian Metric

1.5 Problems

References

- [1] L.W. Tu. *An Introduction to Manifolds*. Universitext. Springer New York.