

A Ready To Use Web-Application Providing a Personalized Biopsy Schedule for Men With Low-Risk PCa Under Active Surveillance^{*}

Anirudh Tomer, MSc^{a,*}, Daan Nieboer, MSc^b, Monique J. Roobol, PhD^c, Anders Bjartell, MD, PhD^d, Ewout W. Steyerberg, PhD^{b,e}, Dimitris Rizopoulos, PhD^a, Movember Foundations Global Action Plan Prostate Cancer Active Surveillance (GAP3) consortium^f

^a*Department of Biostatistics, Erasmus University Medical Center, Rotterdam, the Netherlands*

^b*Department of Public Health, Erasmus University Medical Center, Rotterdam, the Netherlands*

^c*Department of Urology, Erasmus University Medical Center, Rotterdam, the Netherlands*

^d*Department of Urology, Skåne University Hospital, Malmö, Sweden*

^e*Department of Biomedical Data Sciences, Leiden University Medical Center, Leiden, the Netherlands*

^f*The Movember Foundations Global Action Plan Prostate Cancer Active Surveillance (GAP3) consortium members presented in Appendix A*

Abstract

Background: Prostate cancer active surveillance (AS) patients undergo repeat biopsies. Active treatment advised when biopsy Gleason grade group ≥ 2 (upgrading). Many patients never experience upgrading, yet undergo biopsies frequently. Personalized biopsy decisions based on upgrading risk may

*Word Count Abstract: 300; Word Count Text: 2500

*Corresponding author (Anirudh Tomer): Erasmus MC, kamer flex Na-2823, PO Box 2040, 3000 CA Rotterdam, the Netherlands. Tel: +31 10 70 43393

Email addresses: a.tomer@erasmusmc.nl (Anirudh Tomer, MSc), d.nieboer@erasmusmc.nl (Daan Nieboer, MSc), m.roobol@erasmusmc.nl (Monique J. Roobol, PhD), anders.bjartell@med.lu.se (Anders Bjartell, MD, PhD), e.w.steyerberg@lumc.nl (Ewout W. Steyerberg, PhD), d.rizopoulos@erasmusmc.nl (Dimitris Rizopoulos, PhD)

reduce patient burden.

Objective: Develop risk prediction model and web-application to assist patients/doctors in personalized biopsy decisions.

Design, Setting, and Participants: Model development: world's largest AS study PRIAS, 7813 patients, 1134 experienced upgrading; External validation: largest five cohorts of Movember Foundation's GAP3 database (>20,000 patients, 27 centers worldwide); Data: repeat prostate-specific antigen (PSA) and biopsy Gleason grade.

Outcome Measurements, and Statistical Analysis: Bayesian joint model fitted to PRIAS dataset. Model validated in GAP3 cohorts using risk prediction error, calibration, area under ROC (AUC). Model and personalized biopsy schedules based on predicted risks implemented in a web-application.

Results and Limitations: Upgrading rate at year five of follow-up: 35% in PRIAS, at most 50% in GAP3 cohorts. PSA velocity stronger predictor of upgrading (Hazard Ratio: 2.47, 95%CI: 1.93–2.99) than PSA value (Hazard Ratio: 0.99, 95%CI: 0.89–1.11). Validation: Moderate AUC (0.55–0.75) in PRIAS and GAP3 cohorts. Moderate prediction error (0.1–0.3) in GAP3 cohorts with upgrading rate similar to PRIAS, but large (0.3–0.45) otherwise. Recalibration advised for external cohorts.

Conclusions: We successfully developed and validated a model for predicting upgrading risk, and providing risk based personalized biopsy decisions, in prostate cancer AS. The model made available via a web-application, enables shared decision making of biopsy schedules by comparing fixed and personalized schedules on total biopsies and expected time delay in detecting upgrading.

Patient Summary: Personalized prostate biopsies are a novel alternative to fixed one-size-fits-all schedules. The underlying statistical models are made available through a user-friendly web-application and may help to reduce unnecessary prostate biopsies while maintaining cancer control.


Keywords: Active Surveillance, Biopsies, Personalized Medicine, Prostate Cancer, Shared Decision Making

1. Introduction

Patients with low- and very low-risk screening-detected localized prostate cancer are usually advised active surveillance (AS) instead of immediate radical treatment [1]. In AS, cancer progression is routinely monitored via prostate-specific antigen (PSA), digital rectal examination, and repeat biopsies. Among these, the strongest indicator of cancer-related outcomes is the biopsy Gleason grade group [2]. When the Gleason grade group increases from group 1 (Gleason 3+3) to 2 (Gleason 3+4) or higher, called *upgrading* [3], patients are commonly advised curative treatment [4].

In most AS protocols, biopsies are conducted periodically. Consequently, upgrading is always detected with a time delay (Figure 1). For detecting

A Biopsy every year



B Biopsy every 2 years




Figure 1: **Trade-off between the number of biopsies and time delay in detecting upgrading (Increase in Gleason grade group from 1 to 2 or higher):** The true time of upgrading for the patient in this figure is July 2008. When biopsies are scheduled annually (**Panel A**), upgrading is detected in January 2009 with a time delay of six months, and a total of four biopsies are scheduled. When biopsies are scheduled biennially (**Panel B**), upgrading is detected in January 2010 with a time delay of 18 months, and a total of three biopsies are scheduled. Since biopsies are conducted periodically, the time of upgrading is observed as an interval. For example, between Jan 2008–Jan 2009 in **Panel A** and between Jan 2008–Jan 2010 in **Panel B**.

12 upgrading timely, many AS programs schedule fixed and frequent biopsies
 13 (e.g., annually) for all patients [5, 6]. However, this leads to many unnec-
 14 essary biopsies in slow/non-progressing patients. Biopsies are invasive, may
 15 be painful, and are prone to medical complications such as bleeding and
 16 septicemia[7]. Thus, biopsy burden and patient non-compliance to frequent
 17 biopsies [8] has raised concerns regarding the optimal biopsy schedule [9, 10].
 18 To this end, infrequent schedules such as biennial biopsies have been pro-
 19 posed as an alternative [9, 11]. Although, biennial biopsies may still lead
 20 to five unnecessary biopsies over ten years (current study period of large
 21 AS programs) for slow/non-progressing patients. A promising alternative
 22 to fixed and frequent biopsies is personalized biopsy schedules based on the
 23 patient-specific risk of upgrading (Figure 2).

24 The first challenge in developing personalized biopsy schedules is consoli-
 25 dating accumulated patient data (e.g., PSA, previous biopsy results) into risk
 26 estimates for upgrading. Existing calculators for risk of upgrading [12, 13]
 27 use only the latest PSA measurement of a patient. In contrast, we intend to
 28 utilize all repeated measurements of PSA, previous biopsy results, and base-
 29 line characteristics of a patient. To this end, a suitable model is the joint
 30 model for time-to-event and longitudinal data [14, 15, 16]. A joint model pre-
 31 dicts risk of upgrading in a personalized manner. A subsequent challenge,
 32 however, is translating risks into clinical decisions. For example, a 10% risk
 33 of upgrading can be perceived high/low depending upon the patient age. Pa-
 34 tients may also weigh risks of upgrading with the potential *consequences* of
 35 another biopsy. Two relevant *consequences* of biopsies (Figure 1) are the tim-
 36 ing and total number of biopsies (burden), and the time delay in detecting




Figure 2: **Motivation for personalized risk-based decisions of biopsy:** Patient A (**Panel A**) and B (**Panel B**) had their latest biopsy at year one of follow-up (green vertical line). Patient A's prostate-specific antigen (PSA) profile remained stable until his current visit at year three, whereas patient B's profile has shown a rise. Consequently, patient B's cumulative risk of upgrading at the current visit (year three) is higher than that of patient A. This makes patient B a more suitable candidate for biopsy than Patient A. Risk estimates in this figure are only illustrative.

37 upgrading (smaller is beneficial). The relative importance of these *conse-*
 38 *quences* can vary between the patients, and also over the follow-up period
 39 for the same patient.

40 The goal of this work is to develop a robust, generalizable model that
 41 gives reliable estimates for individualized risk of upgrading, and to create
 42 personalized biopsy schedules based on these risks. To further assist patients
 43 and doctors in shared decision making of a biopsy schedule, we also aim to
 44 provide quantitative estimates of *consequences* of opting for a personalized
 45 biopsy schedule versus the standard fixed protocol. For developing our model
 46 we will use the worlds largest AS dataset PRIAS. Subsequently, we will ex-
 47 ternally validate our model in the largest five AS cohorts from the Movember
 48 Foundation’s GAP3 database [17]. Last, we intend to implement our model
 49 and methodology in a web-application.

50 **2. Patients and Methods**

51 *2.1. Study Cohort*

52 For developing a statistical model to power our web-application, we used
 53 the Prostate Cancer International Active Surveillance (PRIAS) database. It
 54 is an ongoing (December 2006 – to date) prospective cohort study of men with
 55 low- and very-low risk prostate cancer diagnoses [4]. More than 100 medical
 56 centers from 17 countries contributed to PRIAS, using a common protocol
 57 (<https://www.prias-project.org>). Upon inclusion in PRIAS, PSA was
 58 measured quarterly for the first two years of follow-up and semiannually
 59 thereafter. Biopsies were scheduled at year one, four, seven, and ten of
 60 follow-up. Additional yearly biopsies were scheduled when PSA doubling

61 time was between zero and ten years.

62 We selected all 7813 patients who had Gleason grade group 1 [2] at the
 63 time of inclusion in PRIAS (Table 1). Our primary event of interest is in-
 64 crease in this Gleason grade group upon repeat biopsy, called *upgrading* (1134
 65 patients). Upgrading is a trigger for treatment advice in PRIAS. Although,
 66 2250 patients were provided treatment on the basis of their PSA, or num-
 67 ber of biopsy cores with cancer, or anxiety/other reasons. Our reasons for
 68 focusing solely on upgrading are, namely, upgrading is strongly associated
 69 with cancer-related outcomes, and other triggers for treatment vary between
 70 cohorts.

71 2.2. Statistical Model

72 To create personalized biopsy schedules based on patient-specific risk of
 73 upgrading, we aimed to develop a risk prediction model. Available data was
 74 patient age at inclusion in AS, longitudinally measured PSA, timing of repeat
 75 biopsies and corresponding Gleason grades, and observed time of upgrading.
 76 Analysis of this data required modeling the within-patient correlation for
 77 PSA, association between the Gleason grades and PSA profiles of a patient,
 78 and handling missing PSA measurements after a patient experienced up-
 79 grading. In such situations, a commonly used model is the joint model for
 80 time-to-event and longitudinal data [14, 15, 16].

81 Our joint model consisted of two sub-models. First, a linear mixed
 82 model [18] for longitudinally measured PSA (log-transformed). Second, a
 83 relative-risk model (similar to Cox model) for obtaining the risk of upgrad-
 84 ing. In the model for PSA, we fitted a curve to PSA measurements (Panel A,
 85 Figure 3). From each patient’s fitted PSA profile, we extracted the instan-

Table 1: **Summary of the PRIAS dataset.** The primary event of interest is upgrading, that is, increase in Gleason grade group from group 1 to 2 or higher. IQR: interquartile range, PSA: prostate-specific antigen.

Characteristic	Value
Total patients	7813
Upgrading (primary event)	1134
Treatment	2250
Watchful waiting	334
Loss to follow-up	249
Death (unrelated to prostate cancer)	95
Death (related to prostate cancer)	2
Median age at diagnosis (years)	66 (IQR: 61–71)
Median follow-up period per patient (years)	1.8 (IQR: 0.9–4.0)
Total PSA measurements	67578
Median number of PSA measurements per patient	6 (IQR: 4–12)
Median PSA value (ng/mL)	5.7 (IQR: 4.1–7.7)
Total biopsies	15686
Median number of biopsies per patient	2 (IQR: 1–2)

86 taneous PSA velocity. This velocity varies over time (Panel B, Figure 3).
 87 Consequently, it is more precise than the currently used constant PSA ve-
 88 locity assumption [19]. We connected the two sub-models by using the fitted
 89 PSA and instantaneous velocity as predictors in the sub-model for risk of up-
 90 grading (Panel C, Figure 3). Patient age was included in both sub-models.
 91 The parameters of the two sub-models were estimated jointly (Supplemen-
 92 tary A) using the R package **JMbayes** [20].

93 *2.3. Risk of Upgrading Based Personalized Biopsies*

94 The key component in personalized schedules is the cumulative-risk of
 95 upgrading. Given a patient’s accumulated PSA measurements and biopsy
 96 results, our joint model predicted the cumulative-risk of upgrading at his cur-
 97 rent as well as future visit times (Panel C, Figure 3). This cumulative-risk is
 98 updated with more patient data over follow-up (Figure 5, Supplementary B).

99 In PRIAS, patient PSA was measured every six months. If during a PSA
 100 visit, a patient’s predicted cumulative-risk of upgrading was more than a cer-
 101 tain threshold (e.g., 10%), we scheduled an immediate biopsy. We scheduled
 102 future biopsies too because our model predicts patient’s cumulative-risk at his
 103 future follow-up visits as well. We achieved this by repeatedly applying the
 104 same risk threshold rule at each future follow-up visit (Supplementary C).
 105 We maintained a minimum gap of one year between consecutive biopsies
 106 (PRIAS recommendation). Example personalized schedules based on 5%
 107 and 10% risk thresholds are shown in Panel B, Figure 4. Due to the cur-
 108 rently limited follow-up period of PRIAS, we were able to schedule biopsies
 109 during the first six years of follow-up only (Table 12, Supplementary C).

110 The choice of the risk threshold in the personalized schedule dictates the




Figure 3: **Illustration of the joint model on a real PRIAS patient.** **Panel A:** Observed PSA (blue dots) and fitted PSA (solid blue line), log-transformed. **Panel B:** Estimated instantaneous velocity of PSA (log-transformed). **Panel C:** Predicted cumulative-risk of upgrading (95% credible interval shaded). Upgrading is defined as an increase in Gleason grade group from group 1 to 2 or higher. This risk of upgrading is available starting from the time of the latest negative biopsy (vertical green line at year 1 of follow-up). The joint model estimated it by combining the fitted value and fitted instantaneous velocity of PSA (log scale), and time of the latest negative biopsy. Black dashed line at year 3 denotes the time of current visit.




Figure 4: **Illustration of personalized and fixed schedules of biopsies.** The PSA profile of this patient is shown in Figure 3. **Panel A:** Predicted cumulative-risk of upgrading (95% credible interval shaded). **Panel B:** Personalized and fixed schedules of biopsies, with a red 'B' indicating a scheduled biopsy. Green vertical line at year 1 denotes the time of latest negative biopsy. Black dashed line at year 3 denotes time of current visit. **Panel C:** Expected time delay in detecting upgrading (months) for different schedules.

111 *consequences* of following that schedule. *Consequences* are the timing and the
 112 total number of biopsies, and the expected time delay in detecting upgrading.
 113 Our model estimated *consequences* in a personalized manner (Panel B,C in
 114 Figure 4, and Figure 9–11 in Supplementary C) given any schedule of biopsies.
 115 Thus, patients can compare personalized schedules based on different risk
 116 thresholds, with fixed schedules, before making a choice.

117 2.4. Model Validation

118 We validated the PRIAS based model internally using the PRIAS cohort,
 119 and externally using the largest five GAP3 database [17] cohorts. These were,
 120 namely, University of Toronto AS (Toronto), Johns Hopkins AS (Hopkins),
 121 Memorial Sloan Kettering Cancer Center AS (MSKCC), King’s College Lon-
 122 don AS (KCL), and Michigan Urological Surgery Improvement Collaborative
 123 AS (MUSIC). We assessed our model’s ability to discriminate between pa-
 124 tients who experience/do not experience upgrading, via the area under the
 125 receiver operating characteristic curve or AUC [21]. We employed calibra-
 126 tion plots [22, 23] and mean absolute prediction error [21] to graphically and
 127 quantitatively evaluate the prediction accuracy of our model. Due to the
 128 longitudinal nature of AS studies, the AUC and prediction error varies over
 129 follow-up (Supplementary B.1). Lastly, to resolve any potential model mis-
 130 calibration in external GAP3 cohorts we aimed to recalibrate our model’s
 131 baseline hazard of upgrading, individually for each GAP3 cohort (Supple-
 132 mentary B.1).

133 IMPORTATN...make this a subsection and move web-application to re-
 134 sults Finally we aim to transform our model into a user friendly web ap-
 135 plication including visualization of different biopsy risk thresholds and the

136 consequence (i.e dealy time???)

137 2.5. Web-Application

138 We implemented our methodology in a web-application https://emcbiostatistics.shinyapps.io/prias_biopsy_recommender/. It utilizes the joint model fit-
 139 ted to the PRIAS dataset. Currently, the web-application supports PRIAS
 140 and the five external cohorts in which we validated our model. Patient
 141 data can be entered manually or can be uploaded in Microsoft Excel format.
 142 Predictions for risk of upgrading are shown for a currently limited, cohort-
 143 specific, follow-up period (Table 12, Supplementary C). The web-application
 144 allows comparison of the *consequences* of following these schedules: person-
 145 alized schedules based on 5%, 10%, and 15% risk threshold; annual biopsies;
 146 biennial biopsies; and PRIAS schedule.

148 3. Results

149 The rate of upgrading at year five of follow-up was 35% in PRIAS, and
 150 at most 50% in the five validation GAP3 cohorts (Panel B, Figure 5). That
 151 is, many patients do not require any biopsy in the first five years of AS.

152 In the fitted joint model, when patient age increased from 61 to 71
 153 years (25-th to 75-th percentile), the adjusted hazard ratio of upgrading
 154 was 1.45 (95%CI: 1.30–1.63). When fitted PSA value (log scale) increased
 155 from 2.36 to 3.07 (25-th to 75-th percentile), the adjusted hazard ratio was
 156 0.99 (95%CI: 0.89–1.11). When estimated instantaneous PSA (log scale) ve-
 157 locity increased from -0.09 to 0.31 (25-th to 75-th percentile), the adjusted
 158 hazard ratio was 2.47 (95%CI: 1.93–2.99). Hence, instantaneous velocity of

159 PSA was a stronger predictor of upgrading than PSA value. Detailed pa-
 160 rameter estimates are in Supplementary A.2.

161 The time-varying mean absolute prediction error, time-varying AUC, and
 162 calibration plot of our model in different cohorts are shown in Panel B, Fig-
 163 ure 8, Supplementary B; Panel A, Figure 5; and Panel B, Figure 5, respec-
 164 tively. The AUC was moderate (0.55 to 0.75) in all cohorts. Mean absolute
 165 prediction error was large (0.3 to 0.45) in cohorts with rate of upgrading dif-
 166 ferent from PRIAS, and moderate (0.1 to 0.3) otherwise. Our model required
 167 recalibration of baseline hazard of upgrading in all cohorts (Figure 6, Supple-
 168 mentary B). Although, calibration was fine in Johns Hopkins cohort, whose
 169 rate of upgrading was similar to PRIAS (Panel B, Figure 5). The risk pre-
 170 dictions from the recalibrated models were as good as risk predictions from
 171 joint models fitted separately to each cohort (Figure 7, Supplementary B).
 172 Comprehensive validation results are in Supplementary B.

173 Various personalized and fixed biopsy schedules for a demonstration pa-
 174 tient in Figure 4 show that a personalized schedule based on 10% risk thresh-
 175 old leads to one less biopsy than other schedules. At the same time, the cor-
 176 responding time delay in detection of upgrading is expected to be only one
 177 month more than other schedules. A compulsory biopsy was scheduled at
 178 year six (maximum biopsy scheduling time in PRIAS, Supplementary C) in
 179 all schedules for a meaningful comparison between them. Additional demon-
 180 strations are in Figure 9–11, Supplementary C.

181 4. Discussion

182 First mention overall result and performance




Figure 5: **Model Validation Results.** **Panel A:** time dependent area under the receiver operating characteristic curve or AUC (measure of discrimination). **Panel B:** calibration-at-large [22, 23], with solid lines depicting the non-parametric estimate of the cumulative-risk of upgrading [24], and dashed lines showing the average cumulative-risk of upgrading obtained using the joint model fitted to the PRIAS dataset. Full names of Cohorts are *PRIAS*: Prostate Cancer International Active Surveillance, *Toronto*: University of Toronto Active Surveillance, *Hopkins*: Johns Hopkins Active Surveillance, *MSKCC*: Memorial Sloan Kettering Cancer Center Active Surveillance, *KCL*: King's College London Active Surveillance, *MUSIC*: Michigan Urological Surgery Improvement Collaborative Active Surveillance.

183 We developed a web-application for assisting patients/doctors in mak-
 184 ing biopsy decisions during prostate cancer active surveillance (AS). Our
 185 web-application provides the patient’s current and future risks of upgrading
 186 (increase in Gleason grade [2] from grade 1 to 2 or higher), and personal-
 187 ized biopsy schedules based on this risk. Our work has four novel features
 188 over earlier risk calculators [15, 25]. First, for personalized biopsy sched-
 189 ules, we developed a statistical model using the world’s largest AS dataset
 190 PRIAS. Second, for following any biopsy schedule, fixed or personalized, our
 191 model predicts the corresponding time delay in detection of upgrading (less
 192 is beneficial). Thus, patients/doctors can compare schedules before mak-
 193 ing a choice. Third, we externally validated our model in the largest five
 194 GAP3 database [17] AS cohorts. Fourth, we implemented our methodology
 195 in a web-application ([https://emcbiostatistics.shinyapps.io/prias_](https://emcbiostatistics.shinyapps.io/prias_biopsy_recommender/)
 196 [biopsy_recommender/](https://emcbiostatistics.shinyapps.io/prias_biopsy_recommender/)) for PRIAS and validated GAP3 cohorts.

197 Currently, biopsies are decided either according to fixed schedules (e.g., an-
 198 nual biopsies) or utilize PSA. Both approaches have drawbacks [19, 8]. In
 199 particular, PSA has not been exploited fully and correctly. For example,
 200 using observed PSA is incorrect because it has measurement error. Other
 201 approaches utilize only the latest PSA, and/or when they utilize all PSA
 202 data, they assume constant PSA velocity. In contrast, our model employs all
 203 PSA measurements to build a patient-specific profile of PSA. This profile is
 204 allowed to increase/decrease non-linearly over time (non-constant PSA veloc-
 205 ity). Subsequently, the model consolidates the PSA profile, previous biopsy
 206 results, and baseline characteristics of a patient, into a single personalized
 207 risk of upgrading. This risk also gets updated as more patient data becomes

208 available over follow-up. Due to currently limited magnetic resonance imag-
209 ing (MRI) data, we could not incorporate it into our model. However, MRI
210 data can be added as a predictor in our model in the future. Decisions based
211 on information combined from multiple sources can yield better results than
212 based on MRI or PSA alone.

213 MRI is used a lot today and more and more before the first biopsy and a
214 lot in AS patients to avoid re-biopsy. I would suggest to write that it is used
215 in revised protocols and that the value of it is still to be determined when
216 more follow-up data has been collected.

217 Our model is useful for a large number of patients from PRIAS (model
218 development), and the largest five GAP3 database AS cohorts (model ex-
219 ternal validation). These are the University of Toronto AS, Johns Hopkins
220 AS, Memorial Sloan Kettering Cancer Center AS, King's College London
221 AS, and Michigan Urological Surgery Improvement Collaborative AS. Dur-
222 ing validation, we required recalibration of our model's baseline hazard of
223 upgrading, individually for all validation cohorts. Our model's prediction
224 error was moderate in cohorts with rate of upgrading similar to PRIAS, and
225 large otherwise. Both prediction error and AUC can be improved with newer
226 biomarkers or MRI data in the future.

227 Our work has important clinical implications. The rate of upgrading af-
228 ter five years of follow-up was at most 50% in all cohorts (Figure 5). That
229 is, a large number of patients do not require any biopsy during the first
230 five years of follow-up. Given the non-compliance and burden of frequent
231 biopsies [8], the availability of our methodology as a web-application may
232 encourage patients/doctors to consider personalized schedules instead. To

assist them in this decision making, the web-application provides an estimate of time delay in detection of upgrading for both personalized and fixed schedules, in a personalized manner. We hope this will objectively address patient apprehensions regarding adverse outcomes in AS.

This work has certain limitations. Due to currently limited follow-up period of PRIAS and GAP3 cohorts, the proposed model is valid only for a restricted period (Table 12, Supplementary C). This problem can be mitigated by refitting the model with new follow-up data in the future. While we focused only on upgrading, the number of positive biopsy cores can also be used to trigger treatment. We did not consider such additional criteria because they differ between cohorts [5], whereas upgrading is used widely. Upgrading is susceptible to inter-observer variation too. Models which account for this variation [15, 26] will be interesting to investigate further. However, the methodology for personalized scheduling, and for comparison of various schedules need not change.

5. Conclusions

Prediction tool for risk of upgrading presented in a web-based application (https://emcbiostatistics.shinyapps.io/prias_biopsy_recommender/), for assisting patients/doctors in making biopsy decisions during prostate cancer AS. Our web-application provides the patient's current and future risks of upgrading, and personalized biopsy schedules based on this risk. Our web-application enables shared decision making of biopsy schedule by comparing fixed and personalized schedules on the total biopsies and expected time delay in detection of upgrading.

Currently supported cohorts are the worlds largest AS cohort PRIAS (model development), and the largest five GAP3 database cohorts (model external validation). Risk prediction accuracy in validation cohorts was better only if they had the rate of reclassification similar to PRIAS

Instead of mentionre currently supported cohorts say: Not in conclusion, more a general remarks that calibration to a specific setting is adviasble or something like that.

Author Contributions

Anirudh Tomer had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Study concept and design: Tomer, Nieboer, Roobol, Bjartell, and Rizopoulos

Acquisition of data: Tomer, Nieboer, and Roobol

Analysis and interpretation of data: Tomer, Nieboer, and Rizopoulos

Drafting of the manuscript: Tomer, and Rizopoulos

Critical revision of the manuscript for important intellectual content: Tomer, Nieboer, Roobol, Bjartell, Steyerberg, and Rizopoulos

Statistical analyses: Tomer, Nieboer, Steyerberg, and Rizopoulos

Obtaining funding: Roobol, and Rizopoulos

Administrative, technical or material support: Nieboer

Supervision: Roobol, and Rizopoulos

Other: none

280 **Acknowledgments**

281 The first and last authors would like to acknowledge support by Neder-
 282 landse Organisatie voor Wetenschappelijk Onderzoek (the national research
 283 council of the Netherlands) VIDI grant nr. 016.146.301, and Erasmus Uni-
 284 versity Medical Center funding. Part of this work was carried out on the
 285 Dutch national e-infrastructure with the support of SURF Cooperative. The
 286 authors also thank the Erasmus University Medical Center's Cancer Com-
 287 putational Biology Center for giving access to their IT-infrastructure and
 288 software that was used for the computations and data analysis in this study.

289 This work was supported by the Movember Foundation. The funder did
 290 not play any role in the study design, collection, analysis or interpretation of
 291 data, or in the drafting of this paper.

292 **Appendix A. Members of The Movember Foundations Global Ac-** 293 **tion Plan Prostate Cancer Active Surveillance (GAP3) consortium**

294 *Principle Investigators:* Bruce Trock (Johns Hopkins University, The
 295 James Buchanan Brady Urological Institute, Baltimore, USA), Behfar Ehdaie
 296 (Memorial Sloan Kettering Cancer Center, New York, USA), Peter Car-
 297 roll (University of California San Francisco, San Francisco, USA), Christo-
 298 pher Filson (Emory University School of Medicine, Winship Cancer Insti-
 299 tute, Atlanta, USA), Jeri Kim / Christopher Logothetis (MD Anderson
 300 Cancer Centre, Houston, USA), Todd Morgan (University of Michigan and
 301 Michigan Urological Surgery Improvement Collaborative (MUSIC), Michi-
 302 gan, USA), Laurence Klotz (University of Toronto, Sunnybrook Health Sci-
 303 ences Centre, Toronto, Ontario, Canada), Tom Pickles (University of British

304 Columbia, BC Cancer Agency, Vancouver, Canada), Eric Hyndman (Uni-
 305 versity of Calgary, Southern Alberta Institute of Urology, Calgary, Canada),
 306 Caroline Moore (University College London & University College London
 307 Hospital Trust, London, UK), Vincent Gnanapragasam (University of Cam-
 308 bridge & Cambridge University Hospitals NHS Foundation Trust, Cam-
 309 bridge, UK), Mieke Van Hemelrijck (King's College London, London, UK
 310 & Guys and St Thomas NHS Foundation Trust, London, UK), Prokar Das-
 311 gupta (Guys and St Thomas NHS Foundation Trust, London, UK), Chris
 312 Bangma (Erasmus Medical Center, Rotterdam, The Netherlands/ represen-
 313 tative of Prostate cancer Research International Active Surveillance (PRIAS)
 314 consortium), Monique Roobol (Erasmus Medical Center, Rotterdam, The
 315 Netherlands/ representative of Prostate cancer Research International Active
 316 Surveillance (PRIAS) consortium), Arnauld Villers (Lille University Hospi-
 317 tal Center, Lille, France), Antti Rannikko (Helsinki University and Helsinki
 318 University Hospital, Helsinki, Finland), Riccardo Valdagni (Department of
 319 Oncology and Hemato-oncology, Universit degli Studi di Milano, Radia-
 320 tion Oncology 1 and Prostate Cancer Program, Fondazione IRCCS Istituto
 321 Nazionale dei Tumori, Milan, Italy), Antoinette Perry (University College
 322 Dublin, Dublin, Ireland), Jonas Hugosson (Sahlgrenska University Hospital,
 323 Gteborg, Sweden), Jose Rubio-Briones (Instituto Valenciano de Oncologia,
 324 Valencia, Spain), Anders Bjartell (Skne University Hospital, Malm, Swe-
 325 den), Lukas Hefermehl (Kantonsspital Baden, Baden, Switzerland), Lee Lui
 326 Shiong (Singapore General Hospital, Singapore, Singapore), Mark Fryden-
 327 berg (Monash Health; Monash University, Melbourne, Australia), Yoshiyuki
 328 Kakehi / Mikio Sugimoto (Kagawa University Faculty of Medicine, Kagawa,

329 Japan), Byung Ha Chung (Gangnam Severance Hospital, Yonsei University
330 Health System, Seoul, Republic of Korea)

331 *Pathologist:* Theo van der Kwast (Princess Margaret Cancer Centre,
332 Toronto, Canada). Technology Research Partners: Henk Obbink (Royal
333 Philips, Eindhoven, the Netherlands), Wim van der Linden (Royal Philips,
334 Eindhoven, the Netherlands), Tim Hulsen (Royal Philips, Eindhoven, the
335 Netherlands), Cees de Jonge (Royal Philips, Eindhoven, the Netherlands).

336 *Advisory Regional statisticians:* Mike Kattan (Cleveland Clinic, Cleve-
337 land, Ohio, USA), Ji Xinge (Cleveland Clinic, Cleveland, Ohio, USA), Ken-
338 neth Muir (University of Manchester, Manchester, UK), Artitaya Lophatananon
339 (University of Manchester, Manchester, UK), Michael Fahey (Epworth Health-
340 Care, Melbourne, Australia), Ewout Steyerberg (Erasmus Medical Center,
341 Rotterdam, The Netherlands), Daan Nieboer (Erasmus Medical Center, Rot-
342 terdam, The Netherlands); Liying Zhang (University of Toronto, Sunnybrook
343 Health Sciences Centre, Toronto, Ontario, Canada)

344 *Executive Regional statisticians:* Ewout Steyerberg (Erasmus Medical
345 Center, Rotterdam, The Netherlands), Daan Nieboer (Erasmus Medical Cen-
346 ter, Rotterdam, The Netherlands); Kerri Beckmann (King's College London,
347 London, UK & Guys and St Thomas NHS Foundation Trust, London, UK),
348 Brian Denton (University of Michigan, Michigan, USA), Andrew Hayen (Uni-
349 versity of Technology Sydney, Australia), Paul Boutros (Ontario Institute of
350 Cancer Research, Toronto, Ontario, Canada).

351 *Clinical Research Partners IT Experts:* Wei Guo (Johns Hopkins Uni-
352 versity, The James Buchanan Brady Urological Institute, Baltimore, USA),
353 Nicole Benfante (Memorial Sloan Kettering Cancer Center, New York, USA),

354 Janet Cowan (University of California San Francisco, San Francisco, USA),
 355 Dattatraya Patil (Emory University School of Medicine, Winship Cancer In-
 356 stitute, Atlanta, USA), Emily Tolosa (MD Anderson Cancer Centre, Hous-
 357 ton, Texas, USA), Tae-Kyung Kim (University of Michigan and Michigan
 358 Urological Surgery Improvement Collaborative, Ann Arbor, Michigan, USA),
 359 Alexandre Mamedov (University of Toronto, Sunnybrook Health Sciences
 360 Centre, Toronto, Ontario, Canada), Vincent LaPointe (University of British
 361 Columbia, BC Cancer Agency, Vancouver, Canada), Trafford Crump (Uni-
 362 versity of Calgary, Southern Alberta Institute of Urology, Calgary, Canada),
 363 Vasilis Stavriniades (University College London & University College Lon-
 364 don Hospital Trust, London, UK), Jenna Kimberly-Duffell (University of
 365 Cambridge & Cambridge University Hospitals NHS Foundation Trust, Cam-
 366 bridge, UK), Aida Santaolalla (King's College London, London, UK & Guys
 367 and St Thomas NHS Foundation Trust, London, UK), Daan Nieboer (Eras-
 368 mus Medical Center, Rotterdam, The Netherlands), Jonathan Olivier (Lille
 369 University Hospital Center, Lille, France), Tiziana Rancati (Fondazione IR-
 370 CCS Istituto Nazionale dei Tumori di Milano, Milan, Italy), Heln Ahlgren
 371 (Sahlgrenska University Hospital, Gteborg, Sweden), Juanma Mascars (Insti-
 372 tuto Valenciano de Oncologa, Valencia, Spain), Annica Lfgren (Skne Univer-
 373 sity Hospital, Malm, Sweden), Kurt Lehmann (Kantonsspital Baden, Baden,
 374 Switzerland), Catherine Han Lin (Monash University and Epworth Health-
 375 Care, Melbourne, Australia), Hiromi Hiram (Kagawa University, Kagawa,
 376 Japan), Kwang Suk Lee (Yonsei University College of Medicine, Gangnam
 377 Severance Hospital, Seoul, Korea).

378 *Research Advisory Committee:* Guido Jenster (Erasmus MC, Rotterdam,

the Netherlands), Anssi Auvinen (University of Tampere, Tampere, Finland), Anders Bjartell (Skne University Hospital, Malm, Sweden), Masoom Haider (University of Toronto, Toronto, Canada), Kees van Bochove (The Hyve B.V. Utrecht, Utrecht, the Netherlands), Ballentine Carter (Johns Hopkins University, Baltimore, USA until 2018).

Management team: Sam Gledhill (Movember Foundation, Melbourne, Australia), Mark Buzza / Michelle Kouspou (Movember Foundation, Melbourne, Australia), Chris Bangma (Erasmus Medical Center, Rotterdam, The Netherlands), Monique Roobol (Erasmus Medical Center, Rotterdam, The Netherlands), Sophie Bruinsma / Jozien Helleman (Erasmus Medical Center, Rotterdam, The Netherlands).

References

1. Briganti A, Fossati N, Catto JW, Cornford P, Montorsi F, Mottet N, Wirth M, Van Poppel H. Active surveillance for low-risk prostate cancer: the European Association of Urology position in 2018. *European urology* 2018;74(3):357–68.
2. Epstein JI, Egevad L, Amin MB, Delahunt B, Srigley JR, Humphrey PA. The 2014 international society of urological pathology (isup) consensus conference on gleason grading of prostatic carcinoma. *The American journal of surgical pathology* 2016;40(2):244–52.
3. Bruinsma SM, Roobol MJ, Carroll PR, Klotz L, Pickles T, Moore CM, Gnanapragasam VJ, Villers A, Rannikko A, Valdagni R, et al. Expert consensus document: semantics in active surveillance for men with lo-

- 402 calized prostate cancer results of a modified delphi consensus procedure.
 403 *Nature reviews urology* 2017;14(5):312.
- 404 4. Bul M, Zhu X, Valdagni R, Pickles T, Kakehi Y, Rannikko A, Bjartell A,
 405 Van Der Schoot DK, Cornel EB, Conti GN, et al. Active surveillance for
 406 low-risk prostate cancer worldwide: the prias study. *European urology*
 407 2013;63(4):597–603.
- 408 5. Nieboer D, Tomer A, Rizopoulos D, Roobol MJ, Steyerberg EW. Active
 409 surveillance: a review of risk-based, dynamic monitoring. *Translational*
 410 *andrology and urology* 2018;7(1):106–15.
- 411 6. Loeb S, Carter HB, Schwartz M, Fagerlin A, Braithwaite RS, Lepor H.
 412 Heterogeneity in active surveillance protocols worldwide. *Reviews in*
 413 *urology* 2014;16(4):202–3.
- 414 7. Loeb S, Vellekoop A, Ahmed HU, Catto J, Emberton M, Nam R, Rosario
 415 DJ, Scattoni V, Lotan Y. Systematic review of complications of prostate
 416 biopsy. *European urology* 2013;64(6):876–92.
- 417 8. Bokhorst LP, Alberts AR, Rannikko A, Valdagni R, Pickles T, Kakehi Y,
 418 Bangma CH, Roobol MJ, PRIAS study group . Compliance rates with
 419 the Prostate Cancer Research International Active Surveillance (PRIAS)
 420 protocol and disease reclassification in noncompliers. *European Urology*
 421 2015;68(5):814–21.
- 422 9. Inoue LY, Lin DW, Newcomb LF, Leonardson AS, Ankerst D, Gulati R,
 423 Carter HB, Trock BJ, Carroll PR, Cooperberg MR, et al. Comparative

- 424 analysis of biopsy upgrading in four prostate cancer active surveillance
425 cohorts. *Annals of internal medicine* 2018;168(1):1–9.
- 426 10. Bratt O, Carlsson S, Holmberg E, Holmberg L, Johansson E, Josefsson
427 A, Nilsson A, Nyberg M, Robinson D, Sandberg J, et al. The study of
428 active monitoring in sweden (sams): a randomized study comparing two
429 different follow-up schedules for active surveillance of low-risk prostate
430 cancer. *Scandinavian journal of urology* 2013;47(5):347–55.
- 431 11. de Carvalho TM, Heijnsdijk EA, de Koning HJ. Estimating the risks
432 and benefits of active surveillance protocols for prostate cancer: a mi-
433 crosimulation study. *BJU international* 2017;119(4):560–6.
- 434 12. Partin AW, Yoo J, Carter HB, Pearson JD, Chan DW, Epstein JI, Walsh
435 PC. The use of prostate specific antigen, clinical stage and gleason score
436 to predict pathological stage in men with localized prostate cancer. *The*
437 *Journal of urology* 1993;150(1):110–4.
- 438 13. Makarov DV, Trock BJ, Humphreys EB, Mangold LA, Walsh PC, Ep-
439 stein JI, Partin AW. Updated nomogram to predict pathologic stage of
440 prostate cancer given prostate-specific antigen level, clinical stage, and
441 biopsy gleason score (partin tables) based on cases from 2000 to 2005.
442 *Urology* 2007;69(6):1095–101.
- 443 14. Tomer A, Nieboer D, Roobol MJ, Steyerberg EW, Rizopoulos D. Per-
444 sonalized schedules for surveillance of low-risk prostate cancer patients.
445 *Biometrics* 2019;75(1):153–62.

- 446 15. Coley RY, Zeger SL, Mamawala M, Pienta KJ, Carter HB. Prediction
 447 of the pathologic gleason score to inform a personalized management
 448 program for prostate cancer. *European urology* 2017;72(1):135–41.
- 449 16. Rizopoulos D. Joint Models for Longitudinal and Time-to-Event Data:
 450 With Applications in R. CRC Press; 2012. ISBN 9781439872864.
- 451 17. Bruinsma SM, Zhang L, Roobol MJ, Bangma CH, Steyerberg EW,
 452 Nieboer D, Van Hemelrijck M, consortium MFGAPPCASG, Trock B,
 453 Ehdaie B, et al. The movember foundation’s gap3 cohort: a profile of
 454 the largest global prostate cancer active surveillance database to date.
 455 *BJU international* 2018;121(5):737–44.
- 456 18. Laird NM, Ware JH, et al. Random-effects models for longitudinal data.
 457 *Biometrics* 1982;38(4):963–74.
- 458 19. Vickers AJ, Savage C, O’Brien MF, Lilja H. Systematic review of pre-
 459 treatment prostate-specific antigen velocity and doubling time as pre-
 460 dictors for prostate cancer. *Journal of Clinical Oncology* 2009;27(3):398.
- 461 20. Rizopoulos D. The R package JMbayses for fitting joint models for lon-
 462 gitudinal and time-to-event data using MCMC. *Journal of Statistical*
 463 *Software* 2016;72(7):1–46.
- 464 21. Rizopoulos D, Molenberghs G, Lesaffre EM. Dynamic predictions with
 465 time-dependent covariates in survival analysis using joint modeling and
 466 landmarking. *Biometrical Journal* 2017;59(6):1261–76.
- 467 22. Royston P, Altman DG. External validation of a cox prognostic

- 468 model: principles and methods. *BMC medical research methodology*
469 2013;13(1):33.
- 470 23. Steyerberg EW, Vickers AJ, Cook NR, Gerds T, Gonen M, Obuchowski
471 N, Pencina MJ, Kattan MW. Assessing the performance of prediction
472 models: a framework for some traditional and novel measures. *Epidemi-*
473 *ology (Cambridge, Mass)* 2010;21(1):128.
- 474 24. Turnbull BW. The empirical distribution function with arbitrarily
475 grouped, censored and truncated data. *Journal of the Royal Statisti-*
476 *cal Society Series B (Methodological)* 1976;38(3):290–5.
- 477 25. Ankerst DP, Xia J, Thompson Jr IM, Hoefler J, Newcomb LF, Brooks
478 JD, Carroll PR, Ellis WJ, Gleave ME, Lance RS, et al. Precision
479 medicine in active surveillance for prostate cancer: development of the
480 canary–early detection research network active surveillance biopsy risk
481 calculator. *European urology* 2015;68(6):1083–8.
- 482 26. Balasubramanian R, Lagakos SW. Estimation of a failure time distribu-
483 tion based on imperfect diagnostic tests. *Biometrika* 2003;90(1):171–82.