

A Ready To Use Web-Application Providing a Personalized Biopsy Schedule for Men With Low-Risk PCa Under Active Surveillance*

Anirudh Tomer, MSc^{a,*}, Daan Nieboer, MSc^{b,c}, Monique J. Roobol, PhD^c, Anders Bjartell, MD, PhD^d, Ewout W. Steyerberg, PhD^{b,e}, Dimitris Rizopoulos, PhD^a, Movember Foundation's Global Action Plan Prostate Cancer Active Surveillance (GAP3) consortium^f

^a*Department of Biostatistics, Erasmus University Medical Center, Rotterdam, the Netherlands*

^b*Department of Public Health, Erasmus University Medical Center, Rotterdam, the Netherlands*

^c*Department of Urology, Erasmus University Medical Center, Rotterdam, the Netherlands*

^d*Department of Urology, Skåne University Hospital, Malmö, Sweden*

^e*Department of Biomedical Data Sciences, Leiden University Medical Center, Leiden, the Netherlands*

^f*The Movember Foundation's Global Action Plan Prostate Cancer Active Surveillance (GAP3) consortium members presented in Appendix A*

Abstract

Background: Prostate cancer active surveillance (AS) patients undergo repeat biopsies. Active treatment is advised when biopsy Gleason grade group ≥ 2 (*upgrading*). Many patients never experience upgrading, yet undergo biopsies frequently. Personalized biopsy decisions based on upgrading-risk may

*Word count: abstract (headings excluded) 300; Text 2488; Total 2788

*Corresponding author (Anirudh Tomer): Erasmus MC, kamer flex Na-2823, PO Box 2040, 3000 CA Rotterdam, the Netherlands. Tel: +31 10 70 43393

Email addresses: a.tomer@erasmusmc.nl (Anirudh Tomer, MSc), d.nieboer@erasmusmc.nl (Daan Nieboer, MSc), m.roobol@erasmusmc.nl (Monique J. Roobol, PhD), anders.bjartell@med.lu.se (Anders Bjartell, MD, PhD), e.w.steyerberg@lumc.nl (Ewout W. Steyerberg, PhD), d.rizopoulos@erasmusmc.nl (Dimitris Rizopoulos, PhD)

reduce patient burden.

Objective: Develop a risk prediction model and web-application to assist patients/doctors in personalized biopsy decisions.

Design, Setting, and Participants: Model development: world's largest AS study PRIAS, 7813 patients, 1134 experienced upgrading; External validation: largest six cohorts of Movember Foundation's GAP3 database ($> 20,000$ patients, 27 centers worldwide); Data: repeat prostate-specific antigen (PSA) and biopsy Gleason grade group.

Outcome Measurements, and Statistical Analysis: A Bayesian joint model fitted to PRIAS dataset. This model was validated in GAP3 cohorts using risk prediction error, calibration, area under ROC (AUC). Model and personalized biopsy schedules based on predicted risks were implemented in a web-application.

Results and Limitations: Cause-specific cumulative upgrading-risk at year five of follow-up: 35% in PRIAS, at most 50% in GAP3 cohorts. PRIAS based model: PSA velocity was stronger predictor of upgrading (Hazard Ratio: 2.47, 95%CI: 1.93–2.99) than PSA value (Hazard Ratio: 0.99, 95%CI: 0.89–1.11). Validation: Moderate AUC (0.55–0.75) in PRIAS and GAP3 cohorts. Moderate prediction error (0.1–0.3) in GAP3 cohorts where impact of PSA value and velocity on upgrading-risk was similar to PRIAS, but large (0.3–0.45) otherwise. Recalibration of baseline risk required for external cohorts.

Conclusions: We successfully developed and validated a model for predicting upgrading-risk, and providing risk-based personalized biopsy decisions, in prostate cancer AS. The model made available via a web-application enables shared decision making of biopsy schedules by comparing fixed and personalized schedules on total biopsies and expected time delay in detecting upgrading.

Patient Summary: Personalized prostate biopsies are a novel alternative to fixed one-size-fits-all schedules. The underlying statistical models are made available through a user-friendly web-application and may help to reduce unnecessary prostate biopsies while maintaining cancer control.


Keywords: Active Surveillance, Biopsies, Personalized Medicine, Prostate Cancer, Shared Decision Making

1. Introduction

Patients with low- and very low-risk screening-detected localized prostate cancer are usually recommended active surveillance (AS) instead of immediate radical treatment [1]. In AS, cancer progression is routinely monitored via prostate-specific antigen (PSA), digital rectal examination, and repeat biopsies. Among these, the strongest indicator of cancer-related outcomes is the biopsy Gleason grade group [2]. When it increases from group 1 (Gleason 3+3) to 2 (Gleason 3+4) or higher, called *upgrading* [3], patients are commonly advised curative treatment [4].

Usually, AS protocols schedule biopsies periodically. Consequently, up-

A Biopsy every year



B Biopsy every 2 years




Figure 1: **Trade-off between the number of biopsies and time delay in detecting upgrading (Increase in Gleason grade group from 1 to 2 or higher):** The true time of upgrading for the patient in this figure is July 2008. When biopsies are scheduled annually (**Panel A**), upgrading is detected in January 2009 with a time delay of six months, and a total of four biopsies are scheduled. When biopsies are scheduled biennially (**Panel B**), upgrading is detected in January 2010 with a time delay of 18 months, and a total of three biopsies are scheduled. Since biopsies are conducted periodically, the time of upgrading is observed as an interval. For example, between Jan 2008–Jan 2009 in **Panel A** and between Jan 2008–Jan 2010 in **Panel B**. The phrase ‘Gleason grade group’ is shortened to ‘Gleason grade’ for brevity.

grading is always detected with a time delay (Figure 1). For detecting upgrading timely, many AS programs schedule fixed and frequent biopsies (e.g., annually) for all patients [5, 6]. However, this leads to unnecessary biopsies in slow/non-progressing patients. Biopsies are invasive, may be painful, and are prone to medical complications such as bleeding and septicemia[7]. Biopsy burden and patient non-compliance to frequent biopsies [8] have raised concerns regarding the optimal biopsy schedule [9, 10]. In this regard, in some cohorts, magnetic resonance imaging (MRI) is employed for targeted biopsies and to study its value for tumor monitoring. Although, due to currently limited AS data, MRI’s value is not clear. Others have proposed the option of scheduling biopsies infrequently (e.g., biennially) [9, 11]. However, due to differences in baseline upgrading-risk across cohorts [9], fixed biopsy schemes can still lead to many unnecessary biopsies. A promising alternative to fixed schedules are personalized biopsy schedules based on the patient-specific upgrading-risk (Figure 2).

The first challenge in creating personalized biopsy schedules is developing a statistical model to consolidate accumulated patient data (e.g., PSA, previous biopsy results) into predictions for upgrading-risk. Existing upgrading-risk [12, 13] calculators use only the latest PSA measurement of a patient. Comparatively, more information is captured by considering all repeatedly measured PSA, previous biopsy results, and baseline characteristics of a patient. To this end, a suitable model is the joint model for time-to-event and longitudinal data [14, 15, 16]. A joint model predicts upgrading-risk in a personalized manner. However, a subsequent challenge is translating predicted risks into clinical decisions. For example, a 10% upgrading-risk can

A Should a biopsy be conducted at current visit?




Figure 2: **Motivation for personalized upgrading-risk based decisions of biopsy:** Patient A (**Panel A**) and B (**Panel B**) had their latest biopsy at year one of follow-up (green vertical line). Patient A's prostate-specific antigen (PSA) profile remained stable until his current visit at year two, whereas patient B's profile has shown a rise. Consequently, patient B's upgrading-risk at the current visit (year two) is higher than that of patient A. This makes patient B a more suitable candidate for biopsy than Patient A. Risk estimates in this figure are only illustrative.

36 be perceived high/low depending upon the patient’s age. Patients may also
 37 weigh risks of upgrading with the potential *consequences* of another biopsy.
 38 Two such relevant consequences (Figure 1) are the timing and the total num-
 39 ber of planned biopsies (burden), and the time delay in detecting upgrading
 40 (smaller is beneficial). The relative importance of these consequences can
 41 vary between the patients, and also within a patient over the follow-up pe-
 42 riod.

43 The goal of this work is two-fold. First, to develop a robust, general-
 44 izable model that gives reliable estimates for individual upgrading-risk in
 45 AS. Second, to utilize the predicted upgrading-risks to create personalized
 46 biopsy schedules. To facilitate shared decision making of biopsy schedules,
 47 we also intend to provide quantitative estimates of the aforementioned *con-*
 48 *sequences* of opting for a personalized versus the standard fixed schedule.
 49 For developing our model, we will use the world’s largest AS dataset PRIAS.
 50 Subsequently, we want to externally validate our model in the largest six
 51 AS cohorts from the Movember Foundation’s GAP3 database [17]. Last, we
 52 intend to implement our model and methodology in a web-application.

53 2. Patients and Methods

54 2.1. Study Cohort

55 For developing a statistical model to predict upgrading-risk, we used the
 56 world’s largest AS dataset, Prostate Cancer International Active Surveillance
 57 or PRIAS [4], dated April 2019 (Table 1). In PRIAS, PSA was measured
 58 quarterly for the first two years of follow-up and semiannually thereafter.
 59 Biopsies were scheduled at year one, four, seven, and ten of follow-up. Addi-

60 tional yearly biopsies were scheduled when PSA doubling time was between
 61 zero and ten years.

62 We selected all 7813 patients who had Gleason grade group 1 at the time
 63 of inclusion in PRIAS. Our primary event of interest is an increase in this
 64 Gleason grade group observed upon repeat biopsy, called *upgrading* (1134 pa-
 65 tients). Upgrading is a trigger for treatment advice in PRIAS. Also, 2250 pa-
 66 tients were provided treatment based on their PSA, number of biopsy cores
 67 with cancer, or anxiety/other reasons. Our reasons for focusing solely on
 68 upgrading are, namely, upgrading is strongly associated with cancer-related
 69 outcomes, and other treatment triggers vary between cohorts [5].

70 For model validation, we selected the following largest (by number of
 71 repeated measurements) six cohorts from Movember Foundation’s GAP3
 72 database version 3.1 [17]: University of California San Francisco AS (UCSF,
 73 version 3.2), University of Toronto AS (Toronto), Johns Hopkins AS (Hop-
 74 kins), Memorial Sloan Kettering Cancer Center AS (MSKCC), King’s College
 75 London AS (KCL), and Michigan Urological Surgery Improvement Collabo-
 76 rative AS (MUSIC). Only patients with a Gleason grade group 1 at the time
 77 of inclusion in these cohorts were selected. Summary statistics are presented
 78 in Supplementary A.2.

79 2.2. Statistical Model

80 For developing an upgrading-risk prediction model, the data we utilized
 81 from the PRIAS cohort was patient age at inclusion in AS, longitudinally
 82 measured PSA, timing of repeat biopsies and Gleason grades, and observed
 83 time of upgrading. Analysis of this data required modeling the within-patient
 84 correlation for PSA, the association between the Gleason grades and PSA

Table 1: **Summary of the PRIAS dataset as of April 2019.** The primary event of interest is upgrading, that is, increase in Gleason grade group from group 1 [2] to 2 or higher. IQR: interquartile range, PSA: prostate-specific antigen. Study protocol URL: <https://www.prias-project.org>

Characteristic	Value
Total centers	> 100
Total patients	7813
Upgrading (primary event)	1134
Treatment	2250
Watchful waiting	334
Loss to follow-up	249
Death (unrelated to prostate cancer)	95
Death (related to prostate cancer)	2
Median age at diagnosis (years)	66 (IQR: 61–71)
Median maximum follow-up per patient (years)	1.8 (IQR: 0.9–4.0)
Total PSA measurements	67578
Median number of PSA measurements per patient	6 (IQR: 4–12)
Median PSA value (ng/mL)	5.7 (IQR: 4.1–7.7)
Total biopsies	15686
Median number of biopsies per patient	2 (IQR: 1–2)

85 profiles of a patient, and handling missing PSA measurements after a patient
 86 experienced upgrading. In such situations, a commonly used model is the
 87 joint model for time-to-event and longitudinal data [14, 15, 16].

88 Our joint model consisted of two sub-models. First, a linear mixed sub-
 89 model [18] for longitudinally measured PSA (log-transformed). Second, a
 90 relative-risk sub-model (similar to the Cox model) for obtaining the cause-
 91 specific upgrading-risk. We included patient age in both sub-models. In the
 92 PSA sub-model, we fitted a unique curve to the PSA measurements of each
 93 patient (Panel A, Figure 3). Subsequently, we calculated the mathematical
 94 derivative of the patient’s fitted PSA profile (Equation 2, Supplementary A),
 95 to obtain his follow-up time specific instantaneous PSA velocity (Panel B,
 96 Figure 3). This instantaneous velocity is a stronger predictor of upgrading
 97 than the widely used average PSA velocity [19]. We modeled the impact
 98 of PSA on upgrading-risk by employing fitted PSA value and instantaneous
 99 velocity as predictors in the risk sub-model. Also, we included the time of
 100 the latest negative biopsy in the risk sub-model (Panel C, Figure 3). The
 101 parameters of the two sub-models were estimated jointly (Supplementary A)
 102 using the R package **JMbayes** [20].

103 *2.3. Risk Prediction and Model Validation*

104 Our model provides predictions for upgrading-risk over the entire fu-
 105 ture follow-up period of a patient. Predictions also automatically update
 106 over follow-up as more patient data becomes available (Figure 5, Supple-
 107 mentary B). We validated our PRIAS based model internally in the PRIAS
 108 cohort, and externally in the largest six GAP3 database cohorts. We em-
 109 ployed calibration plots [21, 22] and follow-up *time-dependent* mean absolute




Figure 3: **Illustration of the joint model on a real PRIAS patient.** **Panel A:** Observed PSA (blue dots) and fitted PSA (solid blue line), log-transformed from ng/mL. **Panel B:** Estimated instantaneous velocity of PSA (log-transformed). **Panel C:** Predicted cause-specific cumulative upgrading-risk (95% credible interval shaded). Upgrading is defined as an increase in the Gleason grade group from group 1 [2] to 2 or higher. This upgrading-risk is calculated starting from the time of the latest negative biopsy (vertical green line at year one of follow-up). The joint model estimated it by combining the fitted PSA (log scale) value and instantaneous velocity, and time of the latest negative biopsy. Black dashed line at year two denotes the time of current visit.

110 risk prediction error or MAPE [23] to graphically and quantitatively evaluate
 111 our model’s risk prediction accuracy, respectively. We assessed our model’s
 112 ability to discriminate between patients who experience/do not experience
 113 upgrading via the time-dependent area under the receiver operating charac-
 114 teristic curve or AUC [23].

115 The aforementioned *time-dependent* AUC and MAPE [23] are temporal
 116 extensions of their standard versions [22] in a longitudinal setting. Specif-
 117 ically, at every six months of follow-up, we calculated a unique AUC and
 118 MAPE for predicting upgrading-risk in the subsequent one year (Supplemen-
 119 tary B.1). For emulating a realistic situation, we calculated the AUC and
 120 MAPE at each follow-up using only the validation data available until that
 121 follow-up. Last, to resolve any potential model miscalibration in validation
 122 cohorts, we aimed to recalibrate our model’s baseline hazard of upgrading
 123 (Supplementary B.1), individually for each cohort.

124 3. Results

125 The cause-specific cumulative upgrading-risk at year five of follow-up was
 126 35% in PRIAS and at most 50% in validation cohorts (Panel B, Figure 4).
 127 Hence, many patients do not require all biopsies planned in the first five
 128 years of AS. In the fitted PRIAS model, the adjusted hazard ratio (aHR)
 129 of upgrading for an increase in patient age from 61 to 71 years (25-th to
 130 75-th percentile) was 1.45 (95%CI: 1.30–1.63). For an increase in fitted PSA
 131 value from 2.36 to 3.07 (25-th to 75-th percentile, log scale), the aHR was
 132 0.99 (95%CI: 0.89–1.11). The strongest predictor of upgrading-risk was in-
 133 stantaneous PSA velocity, with an increase from -0.09 to 0.31 (25-th to 75-th

percentile), giving an aHR of 2.47 (95%CI: 1.93–2.99). The aHR for PSA value and velocity varied between GAP3 cohorts (Supplementary Table 8).

The time-dependent AUC, calibration plot, and time-dependent MAPE of our model are shown in Figure 4, and Supplementary Figure 8. In all cohorts, time-dependent AUC was moderate (0.55 to 0.75) over the whole follow-up period. Time-dependent MAPE was large (0.3 to 0.45) in those cohorts where the impact of PSA on upgrading-risk was different from PRIAS (e.g., MUSIC cohort, Supplementary Table 8), and moderate (0.1 to 0.3) otherwise. In all cohorts, the MAPE decreased rapidly after year one of follow-up. Our model was miscalibrated for validation cohorts (Panel B, Figure 4). Recalibrating the baseline hazard of upgrading in validation cohorts resolved this issue (Supplementary Figure 6). We compared risk predictions from the recalibrated models, with predictions from separately fitted cohort-specific joint models (Supplementary Figure 7). The difference in predictions was lowest in Johns Hopkins cohort (impact of PSA on upgrading-risk similar to PRIAS). Comprehensive results are in Supplementary A.2 and B.

3.1. *Personalized Biopsy Schedules*

We employed the PRIAS based fitted model to create personalized biopsy schedules for real PRIAS patients. Specifically, first using the model and patient’s observed data, we predicted his cumulative upgrading-risk (Figure 5) on all of his future follow-up visits (biannually in PRIAS). Subsequently, we planned biopsies on those future visits where his conditional cumulative upgrading-risk was more than a certain threshold (Supplementary Figure 9). Example personalized schedules based on 5% and 10% risk thresholds are shown in Figure 5, and in Supplementary Figure 10–12. For both personal-




Figure 4: **Model Validation Results.** **Panel A:** time-dependent area under the receiver operating characteristic curve or AUC (measure of discrimination). **Panel B:** calibration-at-large indicates model miscalibration. This is because solid lines depicting the non-parametric estimate of the cause-specific cumulative upgrading-risk [24], and dashed lines showing the average cause-specific cumulative upgrading-risk obtained using the joint model fitted to the PRIAS dataset, are not overlapping. Recalibrating the baseline hazard of upgrading resolved this issue (Supplementary Figure 6). Full names of Cohorts are *PRIAS*: Prostate Cancer International Active Surveillance, *Toronto*: University of Toronto Active Surveillance, *Hopkins*: Johns Hopkins Active Surveillance, *MSKCC*: Memorial Sloan Kettering Cancer Center Active Surveillance, *KCL*: King’s College London Active Surveillance, *MUSIC*: Michigan Urological Surgery Improvement Collaborative Active Surveillance, *UCSF*: University of California San Francisco AS.

159 ized and fixed schedules, we estimated the expected time delay in detecting
 160 upgrading if the patient progresses before the time of the last planned biopsy
 161 (Panel C, Figure 5). This delay is also personalized (Supplementary C.1).
 162 That is, even if two different patients are prescribed the same biopsy schedule,
 163 their expected delays will depend on their individual upgrading-risk profiles.
 164 Patients/doctors can utilize the expected delay and schedule of biopsies as
 165 criteria to compare fixed, and different risk-based personalized schedules.

166 3.2. *Web-Application*

167 We implemented our model and personalized schedules in a user-friendly
 168 web-application [https://emcbiostatistics.shinyapps.io/prias_biopsy_](https://emcbiostatistics.shinyapps.io/prias_biopsy_recommender/)
 169 **recommender/**. Currently, the web-application supports PRIAS and the six
 170 validation cohorts. Patient data can be entered manually and in Microsoft
 171 Excel format. Predictions for upgrading-risk are available for a currently
 172 limited, cohort-specific, follow-up period (Supplementary Table 9). The web-
 173 application visualizes the timing of biopsies, and expected time delay in de-
 174 tecting upgrading, for personalized schedules based on 5%, 10%, and 15%
 175 risk threshold; annual biopsies; biennial biopsies; and PRIAS schedule.

176 4. Discussion

177 We successfully developed and externally validated a model for predicting
 178 upgrading-risk [3] in prostate cancer AS, and providing risk-based personal-
 179 ized biopsy decisions. Our work has four novel features over earlier risk
 180 calculators [15, 25]. First, our model was fitted to the world’s largest AS
 181 dataset PRIAS and externally validated in the largest six cohorts of the
 182 Movember Foundation’s GAP3 database [17]. Second, the model predicts a




Figure 5: **Illustration of personalized and fixed schedules of biopsies for patient from Figure 3.** **Panel A:** Predicted cumulative upgrading-risk (95% credible interval shaded). **Panel B:** Different biopsy schedules with a red 'B' indicating a future biopsy. Risk: 5% and Risk: 10% are personalized schedules in which a biopsy is planned whenever the conditional cause-specific cumulative upgrading-risk is above 5% or 10% risk, respectively. Green vertical line at year one is the time of the latest negative biopsy. Black dashed line at year two denotes the time of the current visit. **Panel C:** Expected time delay in detecting upgrading (years) if patient progresses before year six. A compulsory biopsy was scheduled at year six (maximum biopsy scheduling time in PRIAS, Supplementary C) in all schedules for a meaningful comparison between them.

183 patient’s current and future upgrading-risk in a personalized manner. Third,
 184 using the predicted risks, we created personalized biopsy schedules and also
 185 calculated the expected time delay in detecting upgrading (less is beneficial)
 186 if that schedule was followed. Thus, patients/doctors can compare sched-
 187 ules before making a choice. Fourth, we implemented our methodology in a
 188 user-friendly web-application ([https://emcbiostatistics.shinyapps.io/
 189 prias_biopsy_recommender/](https://emcbiostatistics.shinyapps.io/prias_biopsy_recommender/)) for both PRIAS and validated cohorts.

190 Our model is useful for numerous patients from PRIAS and validated
 191 cohorts. The discrimination ability of our model, exhibited by the *time-*
 192 *dependent* AUC, was moderate (0.55–0.75). This is possible because, unlike
 193 the standard AUC [22], the time-dependent AUC utilizes only the validation
 194 data available until the time at which it is calculated. The same holds for the
 195 time-dependent MAPE (mean absolute prediction error). Although, MAPE
 196 varied much more between cohorts than AUC. In cohorts where the effect
 197 size for the impact of PSA value and velocity on upgrading-risk was similar
 198 to that for PRIAS (e.g., Hopkins cohort), MAPE was moderate. Otherwise,
 199 MAPE was large (e.g., KCL and MUSIC cohorts). In all cohorts, MAPE
 200 decreased rapidly after year one of follow-up. A plausible reason is that at
 201 year one, the validation data also contains those patients who may have been
 202 misclassified as Gleason grade group 1 at the time of inclusion in AS. This
 203 issue can be obviated by scheduling a compulsory biopsy at year one for all
 204 patients (current PRIAS recommendation). Last, we required recalibration
 205 of our model’s baseline hazard of upgrading for all validation cohorts.

206 The clinical implications of our work are as follows. First, the cause-
 207 specific cumulative upgrading-risk at year five of follow-up was at most 50%

208 in all cohorts (Panel B, Figure 4). That is, many patients may not require
 209 all biopsies planned in the first five years of AS. Given the non-compliance
 210 and burden of frequent biopsies [8], the availability of our methodology as a
 211 web-application may encourage patients/doctors to consider upgrading-risk
 212 based personalized schedules instead. An additional advantage of personal-
 213 ized schedules is that they update as more patient data becomes available
 214 over follow-up. We have shown via a simulation study [26] that personalized
 215 schedules may reduce, on average, six biopsies compared to annual schedule
 216 and two biopsies compared to PRIAS schedule in slow/non-progressing AS
 217 patients, while maintaining almost the same time delay in detecting upgrad-
 218 ing as PRIAS schedule. Personalized schedules with different risk thresholds
 219 indeed have different performance. In this regard, to assist patients/doctors
 220 in choosing between fixed schedules and personalized schedules based on
 221 different risk thresholds, the web-application provides a patient-specific esti-
 222 mate of the expected time delay in detecting upgrading, for both personalized
 223 and fixed schedules. We hope that this will objectively address patient ap-
 224 prehensions regarding adverse outcomes in AS.

225 This work has certain limitations. Predictions for upgrading-risk, and
 226 personalized schedules are available only for a currently limited, cohort-
 227 specific, follow-up period (Supplementary Table 9). This problem can be
 228 mitigated by refitting the model with new follow-up data in the future. Re-
 229 cently, some cohorts started utilizing MRI to explore the possibility of tar-
 230 geting visible lesions by biopsy. Presently, the GAP3 database has limited
 231 MRI follow-up data available. As more such data becomes available, the cur-
 232 rent model can be extended to include MRI based predictors. We scheduled

233 biopsies using cause-specific cumulative upgrading-risk, which ignores com-
 234 peting events such as treatment based on the number of positive biopsy cores.
 235 Employing a competing-risk model may lead to improved personalized sched-
 236 ules. Upgrading is susceptible to inter-observer variation too. Models which
 237 account for this variation [15, 27] will be interesting to investigate further.
 238 However, even with an enhanced risk prediction model, the methodology for
 239 personalized scheduling and calculation of expected time delay (Supplemen-
 240 tary C) need not change.

241 5. Conclusions

242 We successfully developed and externally validated a model for predict-
 243 ing upgrading-risk, and providing risk-based personalized biopsy decisions,
 244 in prostate cancer AS. The model made available via a user-friendly web-
 245 application (https://emcbiostatistics.shinyapps.io/prias_biopsy_recommender/)
 246 enables shared decision making of biopsy schedules by comparing fixed and
 247 personalized schedules on total biopsies and expected time delay in detecting
 248 upgrading. Novel biomarkers and MRI data can be added as predictors in
 249 the model to improve predictions in the future. Recalibration of baseline
 250 upgrading-risk is advised for external cohorts.

251 Author Contributions

252 Anirudh Tomer had full access to all the data in the study and takes
 253 responsibility for the integrity of the data and the accuracy of the data anal-
 254 ysis.

255 *Study concept and design:* Tomer, Nieboer, Roobol, Bjartell, and Rizopoulos
 256 *Acquisition of data:* Tomer, Nieboer, and Roobol
 257 *Analysis and interpretation of data:* Tomer, Nieboer, and Rizopoulos
 258 *Drafting of the manuscript:* Tomer, and Rizopoulos
 259 *Critical revision of the manuscript for important intellectual content:* Tomer,
 260 Nieboer, Roobol, Bjartell, Steyerberg, and Rizopoulos
 261 *Statistical analyses:* Tomer, Nieboer, Steyerberg, and Rizopoulos
 262 *Obtaining funding:* Roobol, Steyerberg, and Rizopoulos
 263 *Administrative, technical or material support:* Nieboer
 264 *Supervision:* Roobol, and Rizopoulos
 265 *Other:* none

266 **Acknowledgments**

267 We thank Jozien Helleman from the Department of Urology, Erasmus
 268 University Medical Center, for coordinating the project. The first and last
 269 authors would like to acknowledge support by Nederlandse Organisatie voor
 270 Wetenschappelijk Onderzoek (the national research council of the Nether-
 271 lands) VIDI grant nr. 016.146.301, and Erasmus University Medical Cen-
 272 ter funding. Part of this work was carried out on the Dutch national e-
 273 infrastructure with the support of SURF Cooperative. The authors also
 274 thank the Erasmus University Medical Center's Cancer Computational Biol-
 275 ogy Center for giving access to their IT-infrastructure and software that was
 276 used for the computations and data analysis in this study.

277 The PRIAS website is funded by the Prostate Cancer Research Foun-
 278 dation, Rotterdam (SWOP). This work was supported by the Movember

Foundation. The funder did not play any role in the study design, collection,
analysis or interpretation of data, or in the drafting of this paper.

Appendix A. Members of The Movember Foundations Global Action Plan Prostate Cancer Active Surveillance (GAP3) consortium

Principle Investigators: Bruce Trock (Johns Hopkins University, The James Buchanan Brady Urological Institute, Baltimore, USA), Behfar Ehdaie (Memorial Sloan Kettering Cancer Center, New York, USA), Peter Carroll (University of California San Francisco, San Francisco, USA), Christopher Filson (Emory University School of Medicine, Winship Cancer Institute, Atlanta, USA), Jeri Kim / Christopher Logothetis (MD Anderson Cancer Centre, Houston, USA), Todd Morgan (University of Michigan and Michigan Urological Surgery Improvement Collaborative (MUSIC), Michigan, USA), Laurence Klotz (University of Toronto, Sunnybrook Health Sciences Centre, Toronto, Ontario, Canada), Tom Pickles (University of British Columbia, BC Cancer Agency, Vancouver, Canada), Eric Hyndman (University of Calgary, Southern Alberta Institute of Urology, Calgary, Canada), Caroline Moore (University College London & University College London Hospital Trust, London, UK), Vincent Gnanapragasam (University of Cambridge & Cambridge University Hospitals NHS Foundation Trust, Cambridge, UK), Mieke Van Hemelrijck (King's College London, London, UK & Guys and St Thomas NHS Foundation Trust, London, UK), Prokar Dasgupta (Guys and St Thomas NHS Foundation Trust, London, UK), Chris Bangma (Erasmus Medical Center, Rotterdam, The Netherlands/ representative of Prostate cancer Research International Active Surveillance (PRIAS)

consortium), Monique Roobol (Erasmus Medical Center, Rotterdam, The Netherlands/ representative of Prostate cancer Research International Active Surveillance (PRIAS) consortium), Arnauld Villers (Lille University Hospital Center, Lille, France), Antti Rannikko (Helsinki University and Helsinki University Hospital, Helsinki, Finland), Riccardo Valdagni (Department of Oncology and Hemato-oncology, Universit degli Studi di Milano, Radiation Oncology 1 and Prostate Cancer Program, Fondazione IRCCS Istituto Nazionale dei Tumori, Milan, Italy), Antoinette Perry (University College Dublin, Dublin, Ireland), Jonas Hugosson (Sahlgrenska University Hospital, Gteborg, Sweden), Jose Rubio-Briones (Instituto Valenciano de Oncologia, Valencia, Spain), Anders Bjartell (Skne University Hospital, Malm, Sweden), Lukas Hefermehl (Kantonsspital Baden, Baden, Switzerland), Lee Lui Shiong (Singapore General Hospital, Singapore, Singapore), Mark Frydenberg (Monash Health; Monash University, Melbourne, Australia), Yoshiyuki Kakehi / Mikio Sugimoto (Kagawa University Faculty of Medicine, Kagawa, Japan), Byung Ha Chung (Gangnam Severance Hospital, Yonsei University Health System, Seoul, Republic of Korea)

Pathologist: Theo van der Kwast (Princess Margaret Cancer Centre, Toronto, Canada). Technology Research Partners: Henk Obbink (Royal Philips, Eindhoven, the Netherlands), Wim van der Linden (Royal Philips, Eindhoven, the Netherlands), Tim Hulsén (Royal Philips, Eindhoven, the Netherlands), Cees de Jonge (Royal Philips, Eindhoven, the Netherlands).

Advisory Regional statisticians: Mike Kattan (Cleveland Clinic, Cleveland, Ohio, USA), Ji Xinge (Cleveland Clinic, Cleveland, Ohio, USA), Kenneth Muir (University of Manchester, Manchester, UK), Artitaya Lophatananon

328 (University of Manchester, Manchester, UK), Michael Fahey (Epworth Health-
 329 Care, Melbourne, Australia), Ewout Steyerberg (Erasmus Medical Center,
 330 Rotterdam, The Netherlands), Daan Nieboer (Erasmus Medical Center, Rot-
 331 terdam, The Netherlands); Liying Zhang (University of Toronto, Sunnybrook
 332 Health Sciences Centre, Toronto, Ontario, Canada)

333 *Executive Regional statisticians:* Ewout Steyerberg (Erasmus Medical
 334 Center, Rotterdam, The Netherlands), Daan Nieboer (Erasmus Medical Cen-
 335 ter, Rotterdam, The Netherlands); Kerri Beckmann (King's College London,
 336 London, UK & Guys and St Thomas NHS Foundation Trust, London, UK),
 337 Brian Denton (University of Michigan, Michigan, USA), Andrew Hayen (Uni-
 338 versity of Technology Sydney, Australia), Paul Boutros (Ontario Institute of
 339 Cancer Research, Toronto, Ontario, Canada).

340 *Clinical Research Partners IT Experts:* Wei Guo (Johns Hopkins Uni-
 341 versity, The James Buchanan Brady Urological Institute, Baltimore, USA),
 342 Nicole Benfante (Memorial Sloan Kettering Cancer Center, New York, USA),
 343 Janet Cowan (University of California San Francisco, San Francisco, USA),
 344 Dattatraya Patil (Emory University School of Medicine, Winship Cancer In-
 345 stitute, Atlanta, USA), Emily Tolosa (MD Anderson Cancer Centre, Hous-
 346 ton, Texas, USA), Tae-Kyung Kim (University of Michigan and Michigan
 347 Urological Surgery Improvement Collaborative, Ann Arbor, Michigan, USA),
 348 Alexandre Mamedov (University of Toronto, Sunnybrook Health Sciences
 349 Centre, Toronto, Ontario, Canada), Vincent LaPointe (University of British
 350 Columbia, BC Cancer Agency, Vancouver, Canada), Trafford Crump (Uni-
 351 versity of Calgary, Southern Alberta Institute of Urology, Calgary, Canada),
 352 Vasilis Stavrinos (University College London & University College Lon-

353 don Hospital Trust, London, UK), Jenna Kimberly-Duffell (University of
 354 Cambridge & Cambridge University Hospitals NHS Foundation Trust, Cam-
 355 bridge, UK), Aida Santaolalla (King's College London, London, UK & Guys
 356 and St Thomas NHS Foundation Trust, London, UK), Daan Nieboer (Eras-
 357 mus Medical Center, Rotterdam, The Netherlands), Jonathan Olivier (Lille
 358 University Hospital Center, Lille, France), Tiziana Rancati (Fondazione IR-
 359 CCS Istituto Nazionale dei Tumori di Milano, Milan, Italy), Heln Ahlgren
 360 (Sahlgrenska University Hospital, Gteborg, Sweden), Juanma Mascars (Insti-
 361 tuto Valenciano de Oncologa, Valencia, Spain), Annica Lfgren (Skne Univer-
 362 sity Hospital, Malm, Sweden), Kurt Lehmann (Kantonsspital Baden, Baden,
 363 Switzerland), Catherine Han Lin (Monash University and Epworth Health-
 364 Care, Melbourne, Australia), Hiromi Hiram (Kagawa University, Kagawa,
 365 Japan), Kwang Suk Lee (Yonsei University College of Medicine, Gangnam
 366 Severance Hospital, Seoul, Korea).

367 *Research Advisory Committee:* Guido Jenster (Erasmus MC, Rotterdam,
 368 the Netherlands), Anssi Auvinen (University of Tampere, Tampere, Finland),
 369 Anders Bjartell (Skne University Hospital, Malm, Sweden), Masoom Haider
 370 (University of Toronto, Toronto, Canada), Kees van Bochove (The Hyve
 371 B.V. Utrecht, Utrecht, the Netherlands), Ballentine Carter (Johns Hopkins
 372 University, Baltimore, USA until 2018).

373 *Management team:* Sam Gledhill (Movember Foundation, Melbourne,
 374 Australia), Mark Buzza / Michelle Kouspou (Movember Foundation, Mel-
 375 bourne, Australia), Chris Bangma (Erasmus Medical Center, Rotterdam,
 376 The Netherlands), Monique Roobol (Erasmus Medical Center, Rotterdam,
 377 The Netherlands), Sophie Bruinsma / Jozien Helleman (Erasmus Medical

Center, Rotterdam, The Netherlands).

References

1. Briganti A, Fossati N, Catto JW, Cornford P, Montorsi F, Mottet N, Wirth M, Van Poppel H. Active surveillance for low-risk prostate cancer: the European Association of Urology position in 2018. *European urology* 2018;74(3):357–68.
2. Epstein JI, Egevad L, Amin MB, Delahunt B, Srigley JR, Humphrey PA. The 2014 international society of urological pathology (isup) consensus conference on gleason grading of prostatic carcinoma. *The American journal of surgical pathology* 2016;40(2):244–52.
3. Bruinsma SM, Roobol MJ, Carroll PR, Klotz L, Pickles T, Moore CM, Gnanapragasam VJ, Villers A, Rannikko A, Valdagni R, et al. Expert consensus document: semantics in active surveillance for men with localized prostate cancer results of a modified delphi consensus procedure. *Nature reviews urology* 2017;14(5):312.
4. Bul M, Zhu X, Valdagni R, Pickles T, Kakehi Y, Rannikko A, Bjartell A, Van Der Schoot DK, Cornel EB, Conti GN, et al. Active surveillance for low-risk prostate cancer worldwide: the prias study. *European urology* 2013;63(4):597–603.
5. Nieboer D, Tomer A, Rizopoulos D, Roobol MJ, Steyerberg EW. Active surveillance: a review of risk-based, dynamic monitoring. *Translational andrology and urology* 2018;7(1):106–15.

- 400 6. Loeb S, Carter HB, Schwartz M, Fagerlin A, Braithwaite RS, Lepor H.
401 Heterogeneity in active surveillance protocols worldwide. *Reviews in*
402 *urology* 2014;16(4):202–3.
- 403 7. Loeb S, Vellekoop A, Ahmed HU, Catto J, Emberton M, Nam R, Rosario
404 DJ, Scattoni V, Lotan Y. Systematic review of complications of prostate
405 biopsy. *European urology* 2013;64(6):876–92.
- 406 8. Bokhorst LP, Alberts AR, Rannikko A, Valdagni R, Pickles T, Kakehi Y,
407 Bangma CH, Roobol MJ, PRIAS study group . Compliance rates with
408 the Prostate Cancer Research International Active Surveillance (PRIAS)
409 protocol and disease reclassification in noncompliers. *European Urology*
410 2015;68(5):814–21.
- 411 9. Inoue LY, Lin DW, Newcomb LF, Leonardson AS, Ankerst D, Gulati R,
412 Carter HB, Trock BJ, Carroll PR, Cooperberg MR, et al. Comparative
413 analysis of biopsy upgrading in four prostate cancer active surveillance
414 cohorts. *Annals of internal medicine* 2018;168(1):1–9.
- 415 10. Bratt O, Carlsson S, Holmberg E, Holmberg L, Johansson E, Josefsson
416 A, Nilsson A, Nyberg M, Robinson D, Sandberg J, et al. The study of
417 active monitoring in sweden (sams): a randomized study comparing two
418 different follow-up schedules for active surveillance of low-risk prostate
419 cancer. *Scandinavian journal of urology* 2013;47(5):347–55.
- 420 11. de Carvalho TM, Heijnsdijk EA, de Koning HJ. Estimating the risks
421 and benefits of active surveillance protocols for prostate cancer: a mi-
422 crosimulation study. *BJU international* 2017;119(4):560–6.

- 423 12. Partin AW, Yoo J, Carter HB, Pearson JD, Chan DW, Epstein JI, Walsh
 424 PC. The use of prostate specific antigen, clinical stage and gleason score
 425 to predict pathological stage in men with localized prostate cancer. *The*
 426 *Journal of urology* 1993;150(1):110–4.
- 427 13. Makarov DV, Trock BJ, Humphreys EB, Mangold LA, Walsh PC, Ep-
 428 stein JI, Partin AW. Updated nomogram to predict pathologic stage of
 429 prostate cancer given prostate-specific antigen level, clinical stage, and
 430 biopsy gleason score (partin tables) based on cases from 2000 to 2005.
 431 *Urology* 2007;69(6):1095–101.
- 432 14. Tomer A, Nieboer D, Roobol MJ, Steyerberg EW, Rizopoulos D. Per-
 433 sonalized schedules for surveillance of low-risk prostate cancer patients.
 434 *Biometrics* 2019;75(1):153–62.
- 435 15. Coley RY, Zeger SL, Mamawala M, Pienta KJ, Carter HB. Prediction
 436 of the pathologic gleason score to inform a personalized management
 437 program for prostate cancer. *European urology* 2017;72(1):135–41.
- 438 16. Rizopoulos D. Joint Models for Longitudinal and Time-to-Event Data:
 439 With Applications in R. CRC Press; 2012. ISBN 9781439872864.
- 440 17. Bruinsma SM, Zhang L, Roobol MJ, Bangma CH, Steyerberg EW,
 441 Nieboer D, Van Hemelrijck M, consortium MFGAPPCASG, Trock B,
 442 Ehdaie B, et al. The movember foundation’s gap3 cohort: a profile of
 443 the largest global prostate cancer active surveillance database to date.
 444 *BJU international* 2018;121(5):737–44.

- 445 18. Laird NM, Ware JH, et al. Random-effects models for longitudinal data.
446 *Biometrics* 1982;38(4):963–74.
- 447 19. Cooperberg MR, Brooks JD, Faino AV, Newcomb LF, Kearns JT, Car-
448 roll PR, Dash A, Etzioni R, Fabrizio MD, Gleave ME, et al. Refined
449 analysis of prostate-specific antigen kinetics to predict prostate cancer
450 active surveillance outcomes. *European urology* 2018;74(2):211–7.
- 451 20. Rizopoulos D. The R package JMBayes for fitting joint models for lon-
452 gitudinal and time-to-event data using MCMC. *Journal of Statistical*
453 *Software* 2016;72(7):1–46.
- 454 21. Royston P, Altman DG. External validation of a cox prognostic
455 model: principles and methods. *BMC medical research methodology*
456 2013;13(1):33.
- 457 22. Steyerberg EW, Vickers AJ, Cook NR, Gerds T, Gonen M, Obuchowski
458 N, Pencina MJ, Kattan MW. Assessing the performance of prediction
459 models: a framework for some traditional and novel measures. *Epidemi-*
460 *ology (Cambridge, Mass)* 2010;21(1):128.
- 461 23. Rizopoulos D, Molenberghs G, Lesaffre EM. Dynamic predictions with
462 time-dependent covariates in survival analysis using joint modeling and
463 landmarking. *Biometrical Journal* 2017;59(6):1261–76.
- 464 24. Turnbull BW. The empirical distribution function with arbitrarily
465 grouped, censored and truncated data. *Journal of the Royal Statisti-*
466 *cal Society Series B (Methodological)* 1976;38(3):290–5.

- 467 25. Ankerst DP, Xia J, Thompson Jr IM, Hoeffler J, Newcomb LF, Brooks
468 JD, Carroll PR, Ellis WJ, Gleave ME, Lance RS, et al. Precision
469 medicine in active surveillance for prostate cancer: development of the
470 canary–early detection research network active surveillance biopsy risk
471 calculator. *European urology* 2015;68(6):1083–8.
- 472 26. Tomer A, Rizopoulos D, Nieboer D, Drost FJ, Roobol MJ, Steyerberg
473 EW. Personalized decision making for biopsies in prostate cancer active
474 surveillance programs. *Medical Decision Making* 2019;39(5):499–508.
- 475 27. Balasubramanian R, Lagakos SW. Estimation of a failure time distribu-
476 tion based on imperfect diagnostic tests. *Biometrika* 2003;90(1):171–82.