

A Ready To Use Web-Application Providing a Personalized Biopsy Schedule for Men With Low-Risk PCa Under Active Surveillance*

Anirudh Tomer, MSc^{a,*}, Daan Nieboer, MSc^{b,c}, Monique J. Roobol, PhD^c, Anders Bjartell, MD, PhD^d, Ewout W. Steyerberg, PhD^{b,e}, Dimitris Rizopoulos, PhD^a, Movember Foundations Global Action Plan Prostate Cancer Active Surveillance (GAP3) consortium^f

^a*Department of Biostatistics, Erasmus University Medical Center, Rotterdam, the Netherlands*

^b*Department of Public Health, Erasmus University Medical Center, Rotterdam, the Netherlands*

^c*Department of Urology, Erasmus University Medical Center, Rotterdam, the Netherlands*

^d*Department of Urology, Skåne University Hospital, Malmö, Sweden*

^e*Department of Biomedical Data Sciences, Leiden University Medical Center, Leiden, the Netherlands*

^f*The Movember Foundations Global Action Plan Prostate Cancer Active Surveillance (GAP3) consortium members presented in Appendix A*

Abstract

Background: Prostate cancer active surveillance (AS) patients undergo repeat biopsies. Active treatment is advised when biopsy Gleason grade group ≥ 2 (*upgrading*). Many patients never experience upgrading, yet undergo biopsies frequently. Personalized biopsy decisions based on upgrading-risk may

*Word count abstract (headings excluded): 300; Word count text: 2551

*Corresponding author (Anirudh Tomer): Erasmus MC, kamer flex Na-2823, PO Box 2040, 3000 CA Rotterdam, the Netherlands. Tel: +31 10 70 43393

Email addresses: a.tomer@erasmusmc.nl (Anirudh Tomer, MSc), d.nieboer@erasmusmc.nl (Daan Nieboer, MSc), m.roobol@erasmusmc.nl (Monique J. Roobol, PhD), anders.bjartell@med.lu.se (Anders Bjartell, MD, PhD), e.w.steyerberg@lumc.nl (Ewout W. Steyerberg, PhD), d.rizopoulos@erasmusmc.nl (Dimitris Rizopoulos, PhD)

reduce patient burden.

Objective: Develop a risk prediction model and web-application to assist patients/doctors in personalized biopsy decisions.

Design, Setting, and Participants: Model development: world's largest AS study PRIAS, 7813 patients, 1134 experienced upgrading; External validation: largest six cohorts of Movember Foundation's GAP3 database ($> 20,000$ patients, 27 centers worldwide); Data: repeat prostate-specific antigen (PSA) and biopsy Gleason grade.

Outcome Measurements, and Statistical Analysis: A Bayesian joint model fitted to the PRIAS dataset. This model was validated in GAP3 cohorts using risk prediction error, calibration, area under ROC (AUC). Model and personalized biopsy schedules based on predicted risks were implemented in a web-application.

Results and Limitations: Cause-specific cumulative upgrading-risk at year five of follow-up: 35% in PRIAS, at most 50% in GAP3 cohorts. PRIAS based model: PSA velocity was stronger predictor of upgrading (Hazard Ratio: 2.47, 95%CI: 1.93–2.99) than PSA value (Hazard Ratio: 0.99, 95%CI: 0.89–1.11). Validation: Moderate AUC (0.55–0.75) in PRIAS and GAP3 cohorts. Moderate prediction error (0.1–0.3) in GAP3 cohorts where impact of PSA value and velocity on upgrading-risk was similar to PRIAS, but large (0.3–0.45) otherwise. Recalibration of baseline risk required for external cohorts.

Conclusions: We successfully developed and validated a model for predicting upgrading-risk, and providing risk-based personalized biopsy decisions, in prostate cancer AS. The model made available via a web-application enables shared decision making of biopsy schedules by comparing fixed and personalized schedules on total biopsies and expected time delay in detecting upgrading.

Patient Summary: Personalized prostate biopsies are a novel alternative to fixed one-size-fits-all schedules. The underlying statistical models are made available through a user-friendly web-application and may help to reduce unnecessary prostate biopsies while maintaining cancer control.


Keywords: Active Surveillance, Biopsies, Personalized Medicine, Prostate Cancer, Shared Decision Making

1. Introduction

Patients with low- and very low-risk screening-detected localized prostate cancer are usually recommended active surveillance (AS) instead of immediate radical treatment [1]. In AS, cancer progression is routinely monitored via prostate-specific antigen (PSA), digital rectal examination, and repeat biopsies. Among these, the strongest indicator of cancer-related outcomes is the biopsy Gleason grade group [2]. When it increases from group 1 (Gleason 3+3) to 2 (Gleason 3+4) or higher, called *upgrading* [3], patients are commonly advised curative treatment [4].

Usually, AS protocols schedule biopsies periodically. Consequently, up-

A Biopsy every year



B Biopsy every 2 years




Figure 1: **Trade-off between the number of biopsies and time delay in detecting upgrading (Increase in Gleason grade group from 1 to 2 or higher):** The true time of upgrading for the patient in this figure is July 2008. When biopsies are scheduled annually (**Panel A**), upgrading is detected in January 2009 with a time delay of six months, and a total of four biopsies are scheduled. When biopsies are scheduled biennially (**Panel B**), upgrading is detected in January 2010 with a time delay of 18 months, and a total of three biopsies are scheduled. Since biopsies are conducted periodically, the time of upgrading is observed as an interval. For example, between Jan 2008–Jan 2009 in **Panel A** and between Jan 2008–Jan 2010 in **Panel B**. The phrase ‘Gleason grade group’ is shortened to ‘Gleason grade’ for brevity.

grading is always detected with a time delay (Figure 1). For detecting upgrading timely, many AS programs schedule fixed and frequent biopsies (e.g., annually) for all patients [5, 6]. However, this leads to unnecessary biopsies in slow/non-progressing patients. Biopsies are invasive, may be painful, and are prone to medical complications such as bleeding and septicemia[7]. Biopsy burden and patient non-compliance to frequent biopsies [8] have raised concerns regarding the optimal biopsy schedule [9, 10]. In this regard, in some cohorts, magnetic resonance imaging (MRI) is employed for targeted biopsies and to study the value for tumor monitoring. Although, due to currently limited AS data, MRI’s value is not clear. Others have proposed the option of scheduling biopsies infrequently (e.g., biennially) [9, 11]. However, due to differences in baseline upgrading-risk across cohorts [9], fixed biopsy schemes can still lead to many unnecessary biopsies. A promising alternative to fixed and frequent biopsies is personalized biopsy schedules based on the patient-specific upgrading-risk (Figure 2).

The first challenge in creating personalized biopsy schedules is developing a statistical model to consolidate accumulated patient data (e.g., PSA, previous biopsy results) into estimates for upgrading-risk. Existing upgrading-risk [12, 13] calculators use only the latest PSA measurement of a patient. Comparatively, more information is captured by considering all repeatedly measured PSA, previous biopsy results, and baseline characteristics of a patient. To this end, a suitable model is the joint model for time-to-event and longitudinal data [14, 15, 16]. A joint model predicts the upgrading-risk in a personalized manner. However, a subsequent challenge is translating predicted risks into clinical decisions. For example, a 10% upgrading-risk can

A Should a biopsy be conducted at current visit?




Figure 2: **Motivation for personalized upgrading-risk based decisions of biopsy:** Patient A (**Panel A**) and B (**Panel B**) had their latest biopsy at year one of follow-up (green vertical line). Patient A's prostate-specific antigen (PSA) profile remained stable until his current visit at year two, whereas patient B's profile has shown a rise. Consequently, patient B's upgrading-risk at the current visit (year two) is higher than that of patient A. This makes patient B a more suitable candidate for biopsy than Patient A. Risk estimates in this figure are only illustrative.

36 be perceived high/low depending upon the patient’s age. Patients may also
 37 weigh risks of upgrading with the potential *consequences* of another biopsy.
 38 Two relevant consequences of biopsies (Figure 1) are the timing and the to-
 39 tal number of planned biopsies (burden), and the time delay in detecting
 40 upgrading (smaller is beneficial). The relative importance of these conse-
 41 quences can vary between the patients, and also over the follow-up period
 42 for the same patient.

43 The goal of this work is two-fold. First, to develop a robust, general-
 44 izable model that gives reliable estimates for individual upgrading-risk in
 45 AS. Second, to utilize the predicted upgrading-risks to create personalized
 46 biopsy schedules. To facilitate shared decision making of biopsy schedules,
 47 we also intend to provide quantitative estimates of the aforementioned *con-*
 48 *sequences* of opting for a personalized versus the standard fixed schedule.
 49 For developing our model, we will use the world’s largest AS dataset PRIAS.
 50 Subsequently, we want to externally validate our model in the largest six
 51 AS cohorts from the Movember Foundation’s GAP3 database [17]. Last, we
 52 intend to implement our model and methodology in a web-application.

53 2. Patients and Methods

54 2.1. Study Cohort

55 For developing a statistical model to predict upgrading-risk, we used the
 56 world’s largest AS dataset, Prostate Cancer International Active Surveillance
 57 or PRIAS [4] available as of April 2019 (Table 1). In PRIAS, PSA was
 58 measured quarterly for the first two years of follow-up and semiannually
 59 thereafter. Biopsies were scheduled at year one, four, seven, and ten of

60 follow-up. Additional yearly biopsies were scheduled when PSA doubling
 61 time was between zero and ten years.

62 We selected all 7813 patients who had Gleason grade group 1 at the time
 63 of inclusion in PRIAS. Our primary event of interest is an increase in this
 64 Gleason grade group upon repeat biopsy, called *upgrading* (1134 patients).
 65 Upgrading is a trigger for treatment advice in PRIAS. Also, 2250 patients
 66 were provided treatment based on their PSA, number of biopsy cores with
 67 cancer, or anxiety/other reasons. Our reasons for focusing solely on up-
 68 grading are, namely, upgrading is strongly associated with cancer-related
 69 outcomes, and other triggers for treatment vary between cohorts [5].

70 For model validation, we selected the following largest (by number of
 71 repeated measurements) six cohorts from Movember Foundation’s GAP3
 72 database version 3.1 [17]: University of California San Francisco AS (UCSF,
 73 version 3.2), University of Toronto AS (Toronto), Johns Hopkins AS (Hop-
 74 kins), Memorial Sloan Kettering Cancer Center AS (MSKCC), King’s College
 75 London AS (KCL), and Michigan Urological Surgery Improvement Collabo-
 76 rative AS (MUSIC). Only patients with a Gleason grade group 1 at the time
 77 of inclusion in these cohorts were selected. Summary statistics are presented
 78 in Supplementary A.2.

79 2.2. Statistical Model

80 For developing an upgrading-risk prediction model, the available data in
 81 the PRIAS cohort was patient age at inclusion in AS, longitudinally mea-
 82 sured PSA, timing of repeat biopsies and Gleason grades, and observed time
 83 of upgrading. Analysis of this data required modeling the within-patient
 84 correlation for PSA, the association between the Gleason grades and PSA

Table 1: **Summary of the PRIAS dataset as of April 2019.** The primary event of interest is upgrading, that is, increase in Gleason grade group from group 1 [2] to 2 or higher. IQR: interquartile range, PSA: prostate-specific antigen. Study protocol URL: <https://www.prias-project.org>

Characteristic	Value
Total centers	> 100
Total patients	7813
Upgrading (primary event)	1134
Treatment	2250
Watchful waiting	334
Loss to follow-up	249
Death (unrelated to prostate cancer)	95
Death (related to prostate cancer)	2
Median age at diagnosis (years)	66 (IQR: 61–71)
Median maximum follow-up per patient (years)	1.8 (IQR: 0.9–4.0)
Total PSA measurements	67578
Median number of PSA measurements per patient	6 (IQR: 4–12)
Median PSA value (ng/mL)	5.7 (IQR: 4.1–7.7)
Total biopsies	15686
Median number of biopsies per patient	2 (IQR: 1–2)

85 profiles of a patient, and handling missing PSA measurements after a patient
 86 experienced upgrading. In such situations, a commonly used model is the
 87 joint model for time-to-event and longitudinal data [14, 15, 16].

88 Our joint model consisted of two sub-models. First, a linear mixed sub-
 89 model [18] for longitudinally measured PSA (log-transformed). Second, a
 90 relative-risk sub-model (similar to the Cox model) for obtaining the cause-
 91 specific upgrading-risk. We included patient age in both sub-models. In the
 92 PSA sub-model, we fitted a unique curve to the PSA measurements of each
 93 patient (Panel A, Figure 3). Subsequently, we calculated the mathematical
 94 derivative of the patient’s fitted PSA profile (Equation 2, Supplementary A),
 95 to obtain his follow-up time specific instantaneous PSA velocity (Panel B,
 96 Figure 3). This instantaneous velocity is a stronger predictor of upgrading
 97 than the widely used average PSA velocity [19]. We modeled the impact
 98 of PSA on upgrading-risk by including fitted PSA value and instantaneous
 99 velocity as predictors in the relative-risk model. Last, the time of the latest
 100 negative biopsy was utilized in the relative-risk sub-model (Panel C, Fig-
 101 ure 3). The parameters of the two sub-models were estimated jointly (Sup-
 102 plementary A) using the R package **JMbayes** [20].

103 2.3. Risk Prediction and Model Validation

104 The predictions for upgrading-risk from our model are made for the en-
 105 tire future follow-up period of a patient. These predictions automatically
 106 update over follow-up as more patient data becomes available (Figure 5,
 107 Supplementary B). We validated our PRIAS based model internally in the
 108 PRIAS cohort, and externally in the largest six GAP3 database cohorts. We
 109 employed calibration plots [21, 22] and follow-up *time-dependent* mean ab-




Figure 3: **Illustration of the joint model on a real PRIAS patient.** **Panel A:** Observed PSA (blue dots) and fitted PSA (solid blue line), log-transformed from ng/mL. **Panel B:** Estimated instantaneous velocity of PSA (log-transformed). **Panel C:** Predicted cause-specific cumulative upgrading-risk (95% credible interval shaded). Upgrading is defined as an increase in the Gleason grade group from group 1 [2] to 2 or higher. This upgrading-risk is calculated starting from the time of the latest negative biopsy (vertical green line at year 1 of follow-up). The joint model estimated it by combining the fitted PSA (log scale) value and instantaneous velocity, and time of the latest negative biopsy. Black dashed line at year 3 denotes the time of current visit.

110 solute risk prediction error or MAPE [23] to graphically and quantitatively
 111 evaluate our model’s risk prediction accuracy, respectively. We assessed our
 112 model’s ability to discriminate between patients who experience/do not ex-
 113 perience upgrading via the time-dependent area under the receiver operating
 114 characteristic curve or AUC [23].

115 The aforementioned *time-dependent* AUC and MAPE [23] are temporal
 116 extensions of their standard versions [22] in a longitudinal setting. More
 117 specifically, at every six months of follow-up we calculated a unique AUC
 118 and MAPE for predicting upgrading-risk in the subsequent one year (Sup-
 119 plementary B.1). For emulating a realistic situation, we calculated the AUC
 120 and MAPE at each follow-up using only the validation data available un-
 121 til that follow-up. Lastly, to resolve any potential model miscalibration in
 122 validation cohorts, we aimed to recalibrate our model’s baseline hazard of
 123 upgrading (Supplementary A), individually for each cohort.

124 3. Results

125 The cause-specific cumulative upgrading-risk at year five of follow-up was
 126 35% in PRIAS, and at most 50% in the six validation cohorts (Panel B,
 127 Figure 4). That is, many patients may not require all biopsies planned in the
 128 first five years of AS. In the PRIAS based fitted model, the adjusted hazard
 129 ratio (aHR) of upgrading for an increase in patient age from 61 to 71 years
 130 (25-th to 75-th percentile) was 1.45 (95%CI: 1.30–1.63). For an increase in
 131 fitted PSA value (log scale) from 2.36 to 3.07 (25-th to 75-th percentile), the
 132 aHR was 0.99 (95%CI: 0.89–1.11). The strongest predictor of upgrading-risk
 133 was instantaneous PSA velocity, with an increase from -0.09 to 0.31 (25-th to

134 75-th percentile), leading to an aHR of 2.47 (95%CI: 1.93–2.99). The aHR for
 135 PSA value and velocity varied between cohorts (Table 8, Supplementary A.2).

136 The time-dependent MAPE; time-dependent AUC; and calibration plot
 137 of our model in different validation cohorts are shown in Panel B, Figure 8,
 138 Supplementary B; Panel A, Figure 4; and Panel B, Figure 4, respectively. In
 139 all cohorts, time-dependent AUC was moderate (0.55 to 0.75) over the whole
 140 follow-up period. Time-dependent MAPE was large (0.3 to 0.45) in those
 141 cohorts where the impact of PSA value and velocity on upgrading-risk was
 142 different from PRIAS (e.g., MUSIC cohort, Table 8, Supplementary A.2),
 143 and moderate (0.1 to 0.3) otherwise. In all cohorts, the MAPE decreased
 144 rapidly after year one of follow-up. Our model was miscalibrated for vali-
 145 dation cohorts (Panel B, Figure 4). We resolved this by recalibrating the
 146 baseline hazard of upgrading in all cohorts (Figure 6, Supplementary B). We
 147 compared risk predictions from the recalibrated models with predictions from
 148 separately fitted joint models to each cohort (Figure 7, Supplementary B).
 149 The difference in predictions was lowest in Johns Hopkins cohort (impact of
 150 PSA similar to PRIAS, Table 5, Supplementary A.2). Comprehensive results
 151 are in Supplementary A.2 and B.

152 3.1. *Personalized Biopsy Schedules*

153 We utilized the PRIAS based model to create personalized biopsy sched-
 154 ules. Specifically, first, we utilized the model to predict a patient’s cause-
 155 specific cumulative upgrading-risk on his future follow-up visits (usually ev-
 156 ery six months in PRIAS) based on his available data (Figure 5). We then
 157 planned biopsies on those visits where the patient’s conditional cumulative
 158 upgrading-risk was more than a certain threshold (Supplementary C). Exam-




Figure 4: **Model Validation Results.** **Panel A:** time-dependent area under the receiver operating characteristic curve or AUC (measure of discrimination). **Panel B:** calibration-at-large indicates model miscalibration. This is because solid lines depicting the non-parametric estimate of the cause-specific cumulative upgrading-risk [24], and dashed lines showing the average cause-specific cumulative upgrading-risk obtained using the joint model fitted to the PRIAS dataset, are not overlapping. Same plot after recalibration is shown in Figure 6, Supplementary B. Full names of Cohorts are *PRIAS*: Prostate Cancer International Active Surveillance, *Toronto*: University of Toronto Active Surveillance, *Hopkins*: Johns Hopkins Active Surveillance, *MSKCC*: Memorial Sloan Kettering Cancer Center Active Surveillance, *KCL*: King's College London Active Surveillance, *MUSIC*: Michigan Urological Surgery Improvement Collaborative Active Surveillance, *UCSF*: University of California San Francisco AS.

159 ple personalized schedules based on 5% and 10% risk thresholds are shown in
 160 Figure 5, and in Figure 9–11, Supplementary C. For both personalized and
 161 fixed schedules, we estimated the expected time delay in detecting progres-
 162 sion if a patient follows them (Panel C, Figure 5). The delay is calculated in a
 163 patient-specific manner (Supplementary C) and is updated as more data be-
 164 comes available over follow-up. Using expected delay and schedule of biopsies
 165 as criteria, patients/doctors can compare fixed schedules with personalized
 166 schedules based on different risk thresholds.

167 3.2. *Web-Application*

168 We implemented our model and personalized schedules in a user-friendly
 169 web-application [https://emcbiostatistics.shinyapps.io/prias_biopsy_](https://emcbiostatistics.shinyapps.io/prias_biopsy_recommender/)
 170 [recommender/](https://emcbiostatistics.shinyapps.io/prias_biopsy_recommender/). Currently, the web-application supports PRIAS and the six
 171 validation cohorts. Patient data can be entered manually and in Microsoft
 172 Excel format. Predictions for upgrading-risk are available for a currently
 173 limited, cohort-specific, follow-up period (Table 7, Supplementary C). The
 174 web-application visualizes the timing of biopsies, and expected time delay in
 175 detecting upgrading, for personalized schedules based on 5%, 10%, and 15%
 176 risk threshold; annual biopsies; biennial biopsies; and PRIAS schedule.

177 4. Discussion

178 We successfully developed and externally validated a model for predict-
 179 ing upgrading-risk [3] in prostate cancer AS, and providing risk-based per-
 180 sonalized biopsy decisions. Our work has four novel features over earlier
 181 risk calculators [15, 25]. First, our model was fitted to the world’s largest
 182 AS dataset PRIAS and externally validated in the largest six cohorts of




Figure 5: **Illustration of personalized and fixed schedules of biopsies.** Due to a lack of space, the PSA profile of this patient is shown in Figure 3. **Panel A:** Predicted cumulative upgrading-risk (95% credible interval shaded). **Panel B:** Personalized and fixed schedules of biopsies, with a red 'B' indicating a scheduled biopsy. The green vertical line at year 1 denotes the time of the latest negative biopsy. Black dashed line at year 3 denotes the time of the current visit. **Panel C:** Expected time delay in detecting upgrading (months) for different schedules. A compulsory biopsy was scheduled at year six (maximum biopsy scheduling time in PRIAS, Supplementary C) in all schedules for a meaningful comparison between them.

183 the Movember Foundation’s GAP3 database [17]. Second, the model pre-
 184 dicts a patient’s current and future upgrading-risk in a personalized man-
 185 ner. Third, we use the risks to make a personalized schedule, and to also
 186 calculate expected time delay in detecting upgrading (less is beneficial) if
 187 that schedule is followed. Thus, patients/doctors can compare schedules
 188 before making a choice. Fourth, we implemented our methodology in a
 189 user-friendly web-application ([https://emcbiostatistics.shinyapps.io/
 190 prias_biopsy_recommender/](https://emcbiostatistics.shinyapps.io/prias_biopsy_recommender/)) for PRIAS and validated cohorts.

191 Our model is useful for a large number of patients from PRIAS and
 192 validated cohorts. Across these cohorts, our model had a moderate time-
 193 dependent AUC (0.55–0.75), a measure of discrimination. The moderate
 194 AUC can be explained by the fact that, unlike the standard AUC [22], the
 195 time-dependent AUC utilizes only the validation data available until the time
 196 at which it is calculated. The same holds for the time-dependent MAPE
 197 (mean absolute prediction error), although it varied much more between co-
 198 horts than AUC. It was moderate in cohorts where the effect size for impact
 199 of PSA value and velocity on upgrading-risk was similar to that for PRIAS
 200 (e.g., Hopkins cohort). Otherwise, as in the case of KCL or MUSIC cohorts,
 201 the MAPE was large. In all cohorts, MAPE decreased rapidly after year one
 202 of follow-up. This may be explained by the fact at year one the validation
 203 data also consists of those patients who may have been misclassified incor-
 204 rectly as Gleason grade group 1 at the time of inclusion in AS. The currently
 205 practiced compulsory biopsy at year one of follow-up for all patients may
 206 obviate this issue. Last, we required recalibration of our model’s baseline
 207 hazard of upgrading for all validation cohorts.

208 The clinical implications of our work are as follows. First, the cause-
 209 specific cumulative upgrading-risk at year five of follow-up was at most 50%
 210 in all cohorts (Panel B, Figure 4). That is, many patients may not require
 211 all biopsies planned in the first five years of AS. Given the non-compliance
 212 and burden of frequent biopsies [8], the availability of our methodology as a
 213 web-application may encourage patients/doctors to consider upgrading-risk
 214 based personalized schedules instead. An additional advantage of personal-
 215 ized schedules is that they update as more patient data becomes available
 216 over follow-up. We have shown via a simulation study [26] that personalized
 217 schedules may reduce on average six biopsies compared to annual sched-
 218 ule and two biopsies compared to PRIAS schedule in slow/non-progressing
 219 AS patients, while maintaining almost the same time delay in detecting
 220 progression as PRIAS schedule. Personalized schedules with different risk
 221 thresholds indeed have different performance. In this regard, to assist pa-
 222 tients/doctors in choosing between various personalized, and fixed schedules,
 223 the web-application provides a patient-specific estimate of expected time de-
 224 lay in detecting upgrading, for both personalized and fixed schedules. We
 225 hope that this will objectively address patient apprehensions regarding ad-
 226 verse outcomes in AS.

227 This work has certain limitations. Predictions for upgrading-risk and per-
 228 sonalized schedules are available only for a currently limited, cohort-specific,
 229 follow-up period (Table 7, Supplementary C). This problem can be mitigated
 230 by refitting the model with new follow-up data in the future. Recently, some
 231 cohorts utilize MRI to explore the possibility of targeting visible lesions by
 232 biopsy. Currently GAP3 database has limited MRI follow-up data available.

As more such data becomes available, the current model can be extended to include MRI based predictors. We scheduled biopsies using cause-specific cumulative upgrading-risk. Accounting for competing events, such as treatment based on the number of positive biopsy cores, may lead to improved personalized biopsy decisions. Although, in this work, we did not consider such additional triggers for treatment because, unlike upgrading, they differ between cohorts [5]. Upgrading is susceptible to inter-observer variation too. Models which account for this variation [15, 27] will be interesting to investigate further. However, the methodology for personalized scheduling, and for comparison of various schedules need not change.

5. Conclusions

We successfully developed and validated a model for predicting upgrading-risk, and providing risk-based personalized biopsy decisions, in prostate cancer AS. The model made available via a user-friendly web-application (https://emcbiostatistics.shinyapps.io/prias_biopsy_recommender/) enables shared decision making of biopsy schedules by comparing fixed and personalized schedules on total biopsies and expected time delay in detecting upgrading. Novel biomarkers and MRI data can be added as predictors in the model to improve predictions in the future. Recalibration of baseline upgrading-risk is advised for external cohorts.

Author Contributions

Anirudh Tomer had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data anal-

256 ysis.

257 *Study concept and design:* Tomer, Nieboer, Roobol, Bjartell, and Ri-
258 zopoulos

259 *Acquisition of data:* Tomer, Nieboer, and Roobol

260 *Analysis and interpretation of data:* Tomer, Nieboer, and Rizopoulos

261 *Drafting of the manuscript:* Tomer, and Rizopoulos

262 *Critical revision of the manuscript for important intellectual content:* Tomer,
263 Nieboer, Roobol, Bjartell, Steyerberg, and Rizopoulos

264 *Statistical analyses:* Tomer, Nieboer, Steyerberg, and Rizopoulos

265 *Obtaining funding:* Roobol, Steyerberg, and Rizopoulos

266 *Administrative, technical or material support:* Nieboer

267 *Supervision:* Roobol, and Rizopoulos

268 *Other:* none

269 **Acknowledgments**

270 The first and last authors would like to acknowledge support by Neder-
271 landse Organisatie voor Wetenschappelijk Onderzoek (the national research
272 council of the Netherlands) VIDI grant nr. 016.146.301, and Erasmus Uni-
273 versity Medical Center funding. Part of this work was carried out on the
274 Dutch national e-infrastructure with the support of SURF Cooperative. The
275 authors also thank the Erasmus University Medical Center's Cancer Com-
276 putational Biology Center for giving access to their IT-infrastructure and
277 software that was used for the computations and data analysis in this study.

278 The PRIAS website is funded by the Rotterdam Prostate Cancer Research
279 Foundation Rotterdam (SWOP). This work was supported by the Movember

Foundation. The funder did not play any role in the study design, collection,
analysis or interpretation of data, or in the drafting of this paper.

Appendix A. Members of The Movember Foundations Global Action Plan Prostate Cancer Active Surveillance (GAP3) consortium

Principle Investigators: Bruce Trock (Johns Hopkins University, The James Buchanan Brady Urological Institute, Baltimore, USA), Behfar Ehdaie (Memorial Sloan Kettering Cancer Center, New York, USA), Peter Carroll (University of California San Francisco, San Francisco, USA), Christopher Filson (Emory University School of Medicine, Winship Cancer Institute, Atlanta, USA), Jeri Kim / Christopher Logothetis (MD Anderson Cancer Centre, Houston, USA), Todd Morgan (University of Michigan and Michigan Urological Surgery Improvement Collaborative (MUSIC), Michigan, USA), Laurence Klotz (University of Toronto, Sunnybrook Health Sciences Centre, Toronto, Ontario, Canada), Tom Pickles (University of British Columbia, BC Cancer Agency, Vancouver, Canada), Eric Hyndman (University of Calgary, Southern Alberta Institute of Urology, Calgary, Canada), Caroline Moore (University College London & University College London Hospital Trust, London, UK), Vincent Gnanapragasam (University of Cambridge & Cambridge University Hospitals NHS Foundation Trust, Cambridge, UK), Mieke Van Hemelrijck (King's College London, London, UK & Guys and St Thomas NHS Foundation Trust, London, UK), Prokar Dasgupta (Guys and St Thomas NHS Foundation Trust, London, UK), Chris Bangma (Erasmus Medical Center, Rotterdam, The Netherlands/ representative of Prostate cancer Research International Active Surveillance (PRIAS)

304 consortium), Monique Roobol (Erasmus Medical Center, Rotterdam, The
 305 Netherlands/ representative of Prostate cancer Research International Active
 306 Surveillance (PRIAS) consortium), Arnauld Villers (Lille University Hospi-
 307 tal Center, Lille, France), Antti Rannikko (Helsinki University and Helsinki
 308 University Hospital, Helsinki, Finland), Riccardo Valdagni (Department of
 309 Oncology and Hemato-oncology, Universit degli Studi di Milano, Radia-
 310 tion Oncology 1 and Prostate Cancer Program, Fondazione IRCCS Istituto
 311 Nazionale dei Tumori, Milan, Italy), Antoinette Perry (University College
 312 Dublin, Dublin, Ireland), Jonas Hugosson (Sahlgrenska University Hospital,
 313 Gteborg, Sweden), Jose Rubio-Briones (Instituto Valenciano de Oncologa,
 314 Valencia, Spain), Anders Bjartell (Skne University Hospital, Malm, Swe-
 315 den), Lukas Hefermehl (Kantonsspital Baden, Baden, Switzerland), Lee Lui
 316 Shiong (Singapore General Hospital, Singapore, Singapore), Mark Fryden-
 317 berg (Monash Health; Monash University, Melbourne, Australia), Yoshiyuki
 318 Kakehi / Mikio Sugimoto (Kagawa University Faculty of Medicine, Kagawa,
 319 Japan), Byung Ha Chung (Gangnam Severance Hospital, Yonsei University
 320 Health System, Seoul, Republic of Korea)

321 *Pathologist:* Theo van der Kwast (Princess Margaret Cancer Centre,
 322 Toronto, Canada). Technology Research Partners: Henk Obbink (Royal
 323 Philips, Eindhoven, the Netherlands), Wim van der Linden (Royal Philips,
 324 Eindhoven, the Netherlands), Tim Hulsen (Royal Philips, Eindhoven, the
 325 Netherlands), Cees de Jonge (Royal Philips, Eindhoven, the Netherlands).

326 *Advisory Regional statisticians:* Mike Kattan (Cleveland Clinic, Cleve-
 327 land, Ohio, USA), Ji Xinge (Cleveland Clinic, Cleveland, Ohio, USA), Ken-
 328 neth Muir (University of Manchester, Manchester, UK), Artitaya Lophatananon

329 (University of Manchester, Manchester, UK), Michael Fahey (Epworth Health-
 330 Care, Melbourne, Australia), Ewout Steyerberg (Erasmus Medical Center,
 331 Rotterdam, The Netherlands), Daan Nieboer (Erasmus Medical Center, Rot-
 332 terdam, The Netherlands); Liying Zhang (University of Toronto, Sunnybrook
 333 Health Sciences Centre, Toronto, Ontario, Canada)

334 *Executive Regional statisticians:* Ewout Steyerberg (Erasmus Medical
 335 Center, Rotterdam, The Netherlands), Daan Nieboer (Erasmus Medical Cen-
 336 ter, Rotterdam, The Netherlands); Kerri Beckmann (King's College London,
 337 London, UK & Guys and St Thomas NHS Foundation Trust, London, UK),
 338 Brian Denton (University of Michigan, Michigan, USA), Andrew Hayen (Uni-
 339 versity of Technology Sydney, Australia), Paul Boutros (Ontario Institute of
 340 Cancer Research, Toronto, Ontario, Canada).

341 *Clinical Research Partners IT Experts:* Wei Guo (Johns Hopkins Uni-
 342 versity, The James Buchanan Brady Urological Institute, Baltimore, USA),
 343 Nicole Benfante (Memorial Sloan Kettering Cancer Center, New York, USA),
 344 Janet Cowan (University of California San Francisco, San Francisco, USA),
 345 Dattatraya Patil (Emory University School of Medicine, Winship Cancer In-
 346 stitute, Atlanta, USA), Emily Tolosa (MD Anderson Cancer Centre, Hous-
 347 ton, Texas, USA), Tae-Kyung Kim (University of Michigan and Michigan
 348 Urological Surgery Improvement Collaborative, Ann Arbor, Michigan, USA),
 349 Alexandre Mamedov (University of Toronto, Sunnybrook Health Sciences
 350 Centre, Toronto, Ontario, Canada), Vincent LaPointe (University of British
 351 Columbia, BC Cancer Agency, Vancouver, Canada), Trafford Crump (Uni-
 352 versity of Calgary, Southern Alberta Institute of Urology, Calgary, Canada),
 353 Vasilis Stavrinos (University College London & University College Lon-

354 don Hospital Trust, London, UK), Jenna Kimberly-Duffell (University of
 355 Cambridge & Cambridge University Hospitals NHS Foundation Trust, Cam-
 356 bridge, UK), Aida Santaolalla (King's College London, London, UK & Guys
 357 and St Thomas NHS Foundation Trust, London, UK), Daan Nieboer (Eras-
 358 mus Medical Center, Rotterdam, The Netherlands), Jonathan Olivier (Lille
 359 University Hospital Center, Lille, France), Tiziana Rancati (Fondazione IR-
 360 CCS Istituto Nazionale dei Tumori di Milano, Milan, Italy), Heln Ahlgren
 361 (Sahlgrenska University Hospital, Gteborg, Sweden), Juanma Mascars (Insti-
 362 tuto Valenciano de Oncologa, Valencia, Spain), Annica Lfgren (Skne Univer-
 363 sity Hospital, Malm, Sweden), Kurt Lehmann (Kantonsspital Baden, Baden,
 364 Switzerland), Catherine Han Lin (Monash University and Epworth Health-
 365 Care, Melbourne, Australia), Hiromi Hiram (Kagawa University, Kagawa,
 366 Japan), Kwang Suk Lee (Yonsei University College of Medicine, Gangnam
 367 Severance Hospital, Seoul, Korea).

368 *Research Advisory Committee:* Guido Jenster (Erasmus MC, Rotterdam,
 369 the Netherlands), Anssi Auvinen (University of Tampere, Tampere, Finland),
 370 Anders Bjartell (Skne University Hospital, Malm, Sweden), Masoom Haider
 371 (University of Toronto, Toronto, Canada), Kees van Bochove (The Hyve
 372 B.V. Utrecht, Utrecht, the Netherlands), Ballentine Carter (Johns Hopkins
 373 University, Baltimore, USA until 2018).

374 *Management team:* Sam Gledhill (Movember Foundation, Melbourne,
 375 Australia), Mark Buzza / Michelle Koussou (Movember Foundation, Mel-
 376 bourne, Australia), Chris Bangma (Erasmus Medical Center, Rotterdam,
 377 The Netherlands), Monique Roobol (Erasmus Medical Center, Rotterdam,
 378 The Netherlands), Sophie Bruinsma / Jozen Helleman (Erasmus Medical

Center, Rotterdam, The Netherlands).

References

1. Briganti A, Fossati N, Catto JW, Cornford P, Montorsi F, Mottet N, Wirth M, Van Poppel H. Active surveillance for low-risk prostate cancer: the European Association of Urology position in 2018. *European urology* 2018;74(3):357–68.
2. Epstein JI, Egevad L, Amin MB, Delahunt B, Srigley JR, Humphrey PA. The 2014 international society of urological pathology (isup) consensus conference on gleason grading of prostatic carcinoma. *The American journal of surgical pathology* 2016;40(2):244–52.
3. Bruinsma SM, Roobol MJ, Carroll PR, Klotz L, Pickles T, Moore CM, Gnanapragasam VJ, Villers A, Rannikko A, Valdagni R, et al. Expert consensus document: semantics in active surveillance for men with localized prostate cancer results of a modified delphi consensus procedure. *Nature reviews urology* 2017;14(5):312.
4. Bul M, Zhu X, Valdagni R, Pickles T, Kakehi Y, Rannikko A, Bjartell A, Van Der Schoot DK, Cornel EB, Conti GN, et al. Active surveillance for low-risk prostate cancer worldwide: the prias study. *European urology* 2013;63(4):597–603.
5. Nieboer D, Tomer A, Rizopoulos D, Roobol MJ, Steyerberg EW. Active surveillance: a review of risk-based, dynamic monitoring. *Translational andrology and urology* 2018;7(1):106–15.

- 401 6. Loeb S, Carter HB, Schwartz M, Fagerlin A, Braithwaite RS, Lepor H.
402 Heterogeneity in active surveillance protocols worldwide. *Reviews in*
403 *urology* 2014;16(4):202–3.
- 404 7. Loeb S, Vellekoop A, Ahmed HU, Catto J, Emberton M, Nam R, Rosario
405 DJ, Scattoni V, Lotan Y. Systematic review of complications of prostate
406 biopsy. *European urology* 2013;64(6):876–92.
- 407 8. Bokhorst LP, Alberts AR, Rannikko A, Valdagni R, Pickles T, Kakehi Y,
408 Bangma CH, Roobol MJ, PRIAS study group . Compliance rates with
409 the Prostate Cancer Research International Active Surveillance (PRIAS)
410 protocol and disease reclassification in noncompliers. *European Urology*
411 2015;68(5):814–21.
- 412 9. Inoue LY, Lin DW, Newcomb LF, Leonardson AS, Ankerst D, Gulati R,
413 Carter HB, Trock BJ, Carroll PR, Cooperberg MR, et al. Comparative
414 analysis of biopsy upgrading in four prostate cancer active surveillance
415 cohorts. *Annals of internal medicine* 2018;168(1):1–9.
- 416 10. Bratt O, Carlsson S, Holmberg E, Holmberg L, Johansson E, Josefsson
417 A, Nilsson A, Nyberg M, Robinson D, Sandberg J, et al. The study of
418 active monitoring in sweden (sams): a randomized study comparing two
419 different follow-up schedules for active surveillance of low-risk prostate
420 cancer. *Scandinavian journal of urology* 2013;47(5):347–55.
- 421 11. de Carvalho TM, Heijnsdijk EA, de Koning HJ. Estimating the risks
422 and benefits of active surveillance protocols for prostate cancer: a mi-
423 crosimulation study. *BJU international* 2017;119(4):560–6.

- 424 12. Partin AW, Yoo J, Carter HB, Pearson JD, Chan DW, Epstein JI, Walsh
 425 PC. The use of prostate specific antigen, clinical stage and gleason score
 426 to predict pathological stage in men with localized prostate cancer. *The*
 427 *Journal of urology* 1993;150(1):110–4.
- 428 13. Makarov DV, Trock BJ, Humphreys EB, Mangold LA, Walsh PC, Ep-
 429 stein JI, Partin AW. Updated nomogram to predict pathologic stage of
 430 prostate cancer given prostate-specific antigen level, clinical stage, and
 431 biopsy gleason score (partin tables) based on cases from 2000 to 2005.
 432 *Urology* 2007;69(6):1095–101.
- 433 14. Tomer A, Nieboer D, Roobol MJ, Steyerberg EW, Rizopoulos D. Per-
 434 sonalized schedules for surveillance of low-risk prostate cancer patients.
 435 *Biometrics* 2019;75(1):153–62.
- 436 15. Coley RY, Zeger SL, Mamawala M, Pienta KJ, Carter HB. Prediction
 437 of the pathologic gleason score to inform a personalized management
 438 program for prostate cancer. *European urology* 2017;72(1):135–41.
- 439 16. Rizopoulos D. Joint Models for Longitudinal and Time-to-Event Data:
 440 With Applications in R. CRC Press; 2012. ISBN 9781439872864.
- 441 17. Bruinsma SM, Zhang L, Roobol MJ, Bangma CH, Steyerberg EW,
 442 Nieboer D, Van Hemelrijck M, consortium MFGAPPCASG, Trock B,
 443 Ehdaie B, et al. The movember foundation’s gap3 cohort: a profile of
 444 the largest global prostate cancer active surveillance database to date.
 445 *BJU international* 2018;121(5):737–44.

- 446 18. Laird NM, Ware JH, et al. Random-effects models for longitudinal data.
447 *Biometrics* 1982;38(4):963–74.
- 448 19. Cooperberg MR, Brooks JD, Faino AV, Newcomb LF, Kearns JT, Car-
449 roll PR, Dash A, Etzioni R, Fabrizio MD, Gleave ME, et al. Refined
450 analysis of prostate-specific antigen kinetics to predict prostate cancer
451 active surveillance outcomes. *European urology* 2018;74(2):211–7.
- 452 20. Rizopoulos D. The R package JMBayes for fitting joint models for lon-
453 gitudinal and time-to-event data using MCMC. *Journal of Statistical*
454 *Software* 2016;72(7):1–46.
- 455 21. Royston P, Altman DG. External validation of a cox prognostic
456 model: principles and methods. *BMC medical research methodology*
457 2013;13(1):33.
- 458 22. Steyerberg EW, Vickers AJ, Cook NR, Gerds T, Gonen M, Obuchowski
459 N, Pencina MJ, Kattan MW. Assessing the performance of prediction
460 models: a framework for some traditional and novel measures. *Epidemi-*
461 *ology (Cambridge, Mass)* 2010;21(1):128.
- 462 23. Rizopoulos D, Molenberghs G, Lesaffre EM. Dynamic predictions with
463 time-dependent covariates in survival analysis using joint modeling and
464 landmarking. *Biometrical Journal* 2017;59(6):1261–76.
- 465 24. Turnbull BW. The empirical distribution function with arbitrarily
466 grouped, censored and truncated data. *Journal of the Royal Statisti-*
467 *cal Society Series B (Methodological)* 1976;38(3):290–5.

- 468 25. Ankerst DP, Xia J, Thompson Jr IM, Hoeffler J, Newcomb LF, Brooks
469 JD, Carroll PR, Ellis WJ, Gleave ME, Lance RS, et al. Precision
470 medicine in active surveillance for prostate cancer: development of the
471 canary–early detection research network active surveillance biopsy risk
472 calculator. *European urology* 2015;68(6):1083–8.
- 473 26. Tomer A, Rizopoulos D, Nieboer D, Drost FJ, Roobol MJ, Steyerberg
474 EW. Personalized decision making for biopsies in prostate cancer active
475 surveillance programs. *Medical Decision Making* 2019;39(5):499–508.
- 476 27. Balasubramanian R, Lagakos SW. Estimation of a failure time distribu-
477 tion based on imperfect diagnostic tests. *Biometrika* 2003;90(1):171–82.