

Personalized Decision Making for Biopsies in Prostate Cancer Active Surveillance Programs

Medical Decision Making

XX(X):3–19

©The Author(s) 2018

Reprints and permission:

sagepub.co.uk/journalsPermissions.nav

DOI: 10.1177/ToBeAssigned

www.sagepub.com/

SAGE

Anirudh Tomer¹, Dimitris Rizopoulos¹, Daan Nieboer²,
Frank-Jan Drost³, Monique J. Roobol³, and
Ewout W. Steyerberg^{2,4}

Abstract

Background. Low-risk prostate cancer patients enrolled in active surveillance (AS) programs commonly undergo biopsies for examination of cancer progression. Biopsies are conducted as per a fixed and frequent schedule (e.g. annual biopsies), same for all patients. Since biopsies are burdensome, patients do not always comply with the schedule, which increases the risk of delayed detection of cancer progression.

Objective. Our aim is to better balance the number of biopsies (burden) and the delay in detection of cancer progression (less is beneficial), by personalizing the decision of conducting biopsies.

Data Sources. We use patient data of the world's largest AS program called PRIAS. It has 5270 patients, 866 cancer progressions, and an average of 9 prostate-specific antigen (PSA) and 5 digital rectal examination (DRE) measurements per patient.

Methods. Using joint models for time-to-event and longitudinal data, we jointly model the historical DRE and PSA measurements, and biopsy results of a patient at each follow-up visit. This results in a visit and patient-specific, cumulative risk of cancer progression. If this risk is above a certain threshold, we schedule a biopsy. We compare this personalized approach with the currently practiced biopsy schedules via an extensive and realistic simulation study, based on a replica of the patients from the PRIAS program.

Results. Compared to the currently practiced schedules, the personalized approach saves one to seven biopsies per patient, depending upon their time of cancer progression. However, the corresponding delay in the detection of cancer progression due to the personalized approach remains comparable with that of the biopsy schedule of the PRIAS program.

Conclusions. We conclude that the personalized schedules better balance the number of biopsies per detected cancer progression.

Keywords

Active surveillance, biopsy, joint models, personalized medical decisions, prostate cancer

* *

Introduction

Prostate cancer is the second most frequently diagnosed cancer in men worldwide¹. In prostate cancer screening programs, many of the diagnosed tumors are clinically insignificant (over-diagnosed)². To avoid further over-treatment, patients diagnosed with low-grade prostate cancer are commonly advised to join active surveillance (AS) programs. In AS, invasive treatments such as surgery are delayed until cancer progresses. Cancer progression is routinely examined via serum prostate-specific antigen (PSA) measurements, a protein biomarker; digital rectal examination (DRE) measurements, a measure of the size and location of the tumor; medical imaging; and biopsies.

While larger values for PSA and/or DRE, may indicate cancer progression, biopsies are the most reliable cancer progression examination technique used in AS. When a patient's biopsy Gleason score becomes larger than 6 (positive biopsy, cancer progression detected), AS is stopped and the patient is advised treatment³. However, biopsies are invasive, painful, and prone to medical complications^{4,5}. Hence, they are conducted intermittently until a positive biopsy. Consequently, at the time of a positive biopsy, cancer progression is observed with a delay of an unknown duration. This delay is defined as the difference between the time of the positive biopsy and the unobserved true time of cancer progression. Thus, the decision to conduct biopsies requires a fine compromise between the burden of biopsy and the potential delay in the detection of cancer progression.

¹Department of Biostatistics, Erasmus University Medical Center, the Netherlands

²Department of Public Health, Erasmus University Medical Center, the Netherlands

³Department of Urology, Erasmus University Medical Center, the Netherlands

⁴Department of Biomedical Data Sciences, Leiden University Medical Center, the Netherlands

Corresponding author:

Anirudh Tomer, Erasmus MC, t.a.v. Anirudh Tomer / kamer flex Na-2823, PO Box 2040, 3000 CA Rotterdam, the Netherlands.

Email: a.tomer@erasmusmc.nl

*Financial support for this study was provided Netherlands Organization for Scientific Research's VIDI grant nr. 016.146.301, and Erasmus MC funding. The funding agreement ensured the authors independence in designing the study, interpreting the data, writing, and publishing the report.

Word count: 3972

In AS, a delay in the detection of cancer progression around 12 to 14 months is assumed to be unlikely to substantially increase risk for adverse downstream outcomes^{6,7}. However, for biopsies, there is little consensus on the time gap between them⁸. Majority of the AS programs focus on minimizing only the delay in detection of cancer progression, by scheduling biopsies annually (most frequent schedule) for all patients. A drawback of annual biopsies, and other fixed/heuristic schedules^{6,9}, is that they ignore the large variation in the time of cancer progression of AS patients. While they may work well for patients who progress early (*fast progressing*) in AS, but for a large proportion of patients who do not progress until late (*slow progressing*) in AS, many unnecessary burdensome biopsies are scheduled. To mediate the burden between the *fast* and *slow progressing* patients, the world's largest AS program, Prostate Cancer Research International Active Surveillance¹⁰, abbreviated as PRIAS, schedules annual biopsies only for patients with a low PSA doubling time³. For everyone else, PRIAS schedules biopsies at following fixed follow-up times: year one, four, seven, and ten, and every five years thereafter. Despite this effort in PRIAS, patients may get scheduled for four to ten biopsies over a period of ten years. Therefore, compliance for biopsies is low in PRIAS³. This can lead to the original problem of delayed detection of cancer progression, and reduce the effectiveness of AS.

This article is motivated by the need to better balance the number of biopsies (more are burdensome), and the delay in detection of cancer progression (less is beneficial), than currently practiced schedules (see Figure 1). We intend to achieve this by personalizing the decision to conduct biopsies (see Figure 2). For logistical reasons, these decisions are made at a patient's pre-scheduled follow-up visits for DRE and PSA measurements. To develop the personalized decision making methodology, we utilize the data of the patients of the PRIAS study. We model this data, and develop the personalized approach using joint models for time-to-event and longitudinal data^{11,12}. We conduct an extensive and realistic simulation study to compare the personalized approach with the currently practiced schedules.




Figure 1. Burden-benefit frontier: Estimated median number of biopsies, and median delay in detection of cancer progression, due to various currently practiced fixed/heuristic biopsy schedules (red squares), over a follow-up of ten years. These are based on cancer progression times simulated using inversion sampling from the cumulative risk of cancer progression observed in PRIAS study (see Figure 3 and Study Population section). Using personalized decision making for biopsies, we intend to better balance the number of biopsies and the delay (green region), than currently practiced schedules. An ideal biopsy schedule (blue circle) will schedule only one biopsy, exactly at the true time of cancer progression.

Methods

Study Population

To develop our methodology we use the data of prostate cancer patients from the world's largest AS study called PRIAS¹⁰. More than 100 medical centers from 17 countries worldwide contribute to the collection of data, utilizing a common study protocol and a web-based tool, both available at www.prias-project.org. We use data collected over a period of ten years, between December 2006 (beginning of PRIAS study) and December 2016. It consists of 5270 patients. The primary event of interest in this work is cancer progression detected upon a positive biopsy. It




Figure 2. The personalized decision making problem: Available data of a patient j , who had his latest negative biopsy at $t = 2.6$ years. The shaded region shows the time period in which the patient is at risk of cancer progression. His current pre-scheduled follow-up visit for measurement of DRE and PSA is at $s = 4$ years. Using his entire history of DRE $\mathcal{Y}_{dj}(s)$ and PSA $\mathcal{Y}_{pj}(s)$ measurements up to the current visit s , and the time of the latest biopsy t , we intend to make a decision on scheduling a biopsy at the current visit.

is observed in 866 patients, although the time of cancer progression is interval censored because biopsies are scheduled periodically. Biopsies are scheduled as per the PRIAS protocol (see [Introduction](#)). There are three types of competing events, namely death of 63 patients, of which 61 died from reasons unrelated to prostate cancer; removal of 464 patients from AS on the basis of their observed DRE and PSA measurements; and loss to follow-up of 685 patients because of patient anxiety or unknown reasons. In this work, we assume these three types of events to be censoring (see Appendix A.5 for details). However, our model allows removal of patients to depend on observed longitudinal data and baseline covariates of the patient. Under the aforementioned assumption of censoring, Figure 3

shows the cumulative risk of cancer progression over the study follow-up period.




Figure 3. Estimated cumulative risk of cancer progression in AS for patients in the Prostate Cancer Research International Active Surveillance (PRIAS) dataset. Nearly 50% patients (*slow progressing*) do not progress in the ten year follow-up period. Cumulative risk is estimated using nonparametric maximum likelihood estimation¹³, to account for interval censored cancer progression times observed in PRIAS program. Censoring includes death, removal from AS on the basis of observed longitudinal data, and patient dropout.

For all patients, PSA measurements (ng/mL) are scheduled every 3 months for the first 2 years and every 6 months thereafter. The DRE measurements (ordinal scale) are scheduled every 6 months. We use the DRE measurements after converting them to a binary scale, namely $DRE > T1c$ and $DRE = T1c$. A DRE measurement equal to $T1c$ ¹⁴ indicates a clinically inapparent tumor which is not palpable or visible by imaging. Tumors with $DRE > T1c$ are palpable. On average 5 DRE and 9 PSA measurements have been recorded per patient.

A Bivariate Joint Model for the Longitudinal PSA, and DRE Measurements, and Time of Cancer Progression

Let T_i^* denote the true cancer progression time of the i -th patient included in PRIAS. Since biopsies are conducted periodically, T_i^* is observed with interval censoring $l_i < T_i^* \leq r_i$. When progression is observed for the patient at his latest biopsy time r_i , then l_i denotes the time of the second latest biopsy. Otherwise, l_i denotes the time of the latest biopsy and $r_i = \infty$. Let \mathbf{y}_{di} and \mathbf{y}_{pi} denote his observed DRE and PSA longitudinal measurements, respectively. The observed data of all n patients is denoted by $\mathcal{D}_n = \{l_i, r_i, \mathbf{y}_{di}, \mathbf{y}_{pi}; i = 1, \dots, n\}$.

In our joint model, the patient-specific DRE and PSA measurements over time are modeled using a bivariate generalized linear mixed effects sub-model. The sub-model for DRE is given by (see Panel A, Figure 4):

$$\begin{aligned} \text{logit}[\Pr\{y_{di}(t) > \text{T1c}\}] &= \beta_{0d} + b_{0di} + (\beta_{1d} + b_{1di})t \\ &\quad + \beta_{2d}(\text{Age}_i - 70) + \beta_{3d}(\text{Age}_i - 70)^2 \end{aligned} \quad (1)$$

where, t denotes the follow-up visit time, and Age_i is the age of the i -th patient at the time of inclusion in AS. The fixed effect parameters are denoted by $\{\beta_{0d}, \dots, \beta_{3d}\}$, and $\{b_{0di}, b_{1di}\}$ are the patient specific random effects. With this definition, we assume that the patient-specific log odds of obtaining a DRE measurement larger than T1c remain linear over time.

The mixed effects sub-model for PSA is given by (see Panel B, Figure 4):

$$\begin{aligned} \log_2 \{y_{pi}(t) + 1\} &= m_{pi}(t) + \varepsilon_{pi}(t), \\ m_{pi}(t) &= \beta_{0p} + b_{0pi} + \sum_{k=1}^4 (\beta_{kp} + b_{kpi}) B_k(t, \mathcal{K}) \\ &\quad + \beta_{5p}(\text{Age}_i - 70) + \beta_{6p}(\text{Age}_i - 70)^2, \end{aligned} \quad (2)$$

where, $m_{pi}(t)$ denotes the underlying measurement error free value of $\log_2(\text{PSA} + 1)$ transformed^{15,16} measurements at time t . We model it non-linearly over time using B-splines¹⁷. To this end, our B-spline basis function $B_k(t, \mathcal{K})$ has 3 internal knots at $\mathcal{K} = \{0.1, 0.7, 4\}$ years, and boundary knots at 0 and 5.42 years (95-th percentile of the observed follow-up times). The fixed effect parameters are denoted by $\{\beta_{0p}, \dots, \beta_{6p}\}$, and $\{b_{0pi}, \dots, b_{4pi}\}$ are the patient specific random effects.




Figure 4. Illustration of the joint model fitted to the PRIAS dataset. **Panel A:** shows the observed DRE measurements and the fitted probability of obtaining DRE > T1c (Equation 1) for i -th patient. **Panel B:** shows the observed and fitted $\log_2(\text{PSA} + 1)$ measurements (Equation 2). **Panel C:** shows the estimated $\log_2(\text{PSA} + 1)$ velocity (velocity cannot be observed directly) over time. The hazard function (Equation 3) shown in **Panel D**, depends on the fitted log odds of having a DRE > T1c, and the fitted $\log_2(\text{PSA} + 1)$ value and velocity.

The error $\varepsilon_{pi}(t)$ is assumed to be t-distributed with three degrees of freedom (see Appendix B.1) and scale σ , and is independent of the random effects.

To account for the correlation between the DRE and PSA measurements of a patient, we link their corresponding random effects. More specifically, the complete vector of random effects $\mathbf{b}_i = (b_{0di}, b_{1di}, b_{0pi}, \dots, b_{4pi})^T$ is assumed to follow a multivariate normal distribution with mean zero and variance-covariance matrix \mathbf{D} .

To model the impact of DRE and PSA measurements on the risk of cancer progression, our joint model uses a relative risk sub-model. More specifically, the hazard of cancer progression $h_i(t)$ at a time t is given by (see Panel D, Figure 4):

$$h_i(t) = h_0(t) \exp \left(\gamma_1(\text{Age}_i - 70) + \gamma_2(\text{Age}_i - 70)^2 + \alpha_{1d} \text{logit}[\text{Pr}\{y_{di}(t) > \text{T1c}\}] + \alpha_{1p} m_{pi}(t) + \alpha_{2p} \frac{\partial m_{pi}(t)}{\partial t} \right), \quad (3)$$

where, γ_1, γ_2 are the parameters for the effect of age. The parameter α_{1d} models the impact of log odds of obtaining a DRE > T1c on the hazard of cancer progression. The impact of PSA on the hazard of cancer progression is modeled in two ways: a) the impact of the error free underlying PSA value $m_{pi}(t)$ (see Panel B, Figure 4), and b) the impact of the underlying PSA velocity $\partial m_{pi}(t)/\partial t$ (see Panel C, Figure 4). The corresponding parameters are α_{1p} and α_{2p} , respectively. Lastly, $h_0(t)$ is the baseline hazard at time t , and is modeled flexibly using P-splines¹⁸. The detailed specification of the baseline hazard $h_0(t)$, and the joint parameter estimation of the two sub-models using the Bayesian approach (R package **JMbayes**¹⁹) are presented in Appendix A of the supplementary material.

Personalized Decisions for Biopsy

Let us assume that a decision of conducting a biopsy is to be made for a new patient j shown in Figure 2, at his current follow-up visit time s . Let $t \leq s$ be the time of his latest negative biopsy. Let $\mathcal{Y}_{dj}(s)$ and $\mathcal{Y}_{pj}(s)$ denote his observed DRE and PSA measurements up to the current visit, respectively. From the observed measurements we want to extract the underlying measurement error free trend of $\log_2(\text{PSA} + 1)$ values and velocity, and the log odds of obtaining DRE > T1c. We intend to combine them to inform us when the cancer progression is to be expected (see

Figure 5), and to further guide the decision making on whether to conduct a biopsy at the current follow-up visit. The combined information is given by the following posterior predictive distribution $g(T_j^*)$ of his time of cancer progression $T_j^* > t$ (see Appendix A.4 for details):

$$g(T_j^*) = p\{T_j^* \mid T_j^* > t, \mathcal{Y}_{dj}(s), \mathcal{Y}_{pj}(s), \mathcal{D}_n\}. \quad (4)$$

The distribution $g(T_j^*)$ is not only patient-specific, but also updates as extra information is recorded at future follow-up visits.

A key ingredient in the decision of conducting a biopsy for patient j at the current follow-up visit time s is the personalized cumulative risk of observing a cancer progression at time s (illustrated in Figure 5). This risk can be derived from the posterior predictive distribution $g(T_j^*)$ ²⁰, and is given by:

$$R_j(s \mid t) = \Pr\{T_j^* \leq s \mid T_j^* > t, \mathcal{Y}_{dj}(s), \mathcal{Y}_{pj}(s), \mathcal{D}_n\}, \quad s \geq t. \quad (5)$$

A simple and straightforward approach to decide upon conducting a biopsy for patient j at the current follow-up visit would be to do so if his personalized cumulative risk of cancer progression at the visit is higher than a certain threshold $0 \leq \kappa \leq 1$. For example, as shown in Panel B of Figure 5, biopsy at a visit may be scheduled if the personalized cumulative risk is higher than 10% (example risk threshold). This decision making process is iterated over the follow-up period, incorporating on each subsequent visit the newly observed data, until a positive biopsy is observed. Subsequently, an entire personalized schedule of biopsies for each patient can be obtained.

The choice of the risk threshold dictates the schedule of biopsies and has to be made on each subsequent follow-up visit of a patient. In this regard, a straightforward approach is choosing a fixed risk threshold, such as 5% or 10% risk, at all follow-up visits. Fixed risk thresholds may be chosen by patients and/or doctors according to how they weigh the relative harms of doing an unnecessary biopsy versus a missed cancer progression (e.g., 10% threshold means a 1:9 ratio) if the biopsy is not conducted²¹. An alternative approach is that at each follow-up visit a unique threshold is chosen on the basis of its classification accuracy. More specifically, given the time of latest biopsy t of patient j , and his current visit time s we find a visit-specific biopsy threshold κ , which gives the highest cancer progression detection rate (true positive rate, or TPR) for the period $(t, s]$.

However, we also intend to balance for unnecessary biopsies (high false positive rate), or a low number of correct detections (high false negative rate) when the false positive rate is minimized. An approach to mitigate these issues is to maximize the TPR and positive predictive value (PPV) simultaneously. To this end, we utilize the F_1 score, which is a composite of both TPR and PPV (estimated as in Rizopoulos et al., 2017²²), and is defined as:

$$\begin{aligned} F_1(t, s, \kappa) &= 2 \frac{\text{TPR}(t, s, \kappa) \text{PPV}(t, s, \kappa)}{\text{TPR}(t, s, \kappa) + \text{PPV}(t, s, \kappa)}, \\ \text{TPR}(t, s, \kappa) &= \Pr\{R_j(s | t) > \kappa \mid t < T_j^* \leq s\}, \\ \text{PPV}(t, s, \kappa) &= \Pr\{t < T_j^* \leq s \mid R_j(s | t) > \kappa\}. \end{aligned} \quad (6)$$

The F_1 score ranges between 0 and 1, where a value of 1 signifies perfect TPR and PPV. Since a high F_1 score is desired, the visit-specific threshold $\kappa = \arg \max_{\kappa} F_1(t, s, \kappa)$. The criteria on which we evaluate the personalized schedules based on fixed risk threshold at all visits, as well as visit-specific risk threshold using F_1 score, is the total number of biopsies scheduled, and the delay in detection of cancer progression (details in [Results](#)).

Simulation Study

Although the personalized decision making approach is motivated by the PRIAS study, it is not possible to evaluate it on the PRIAS dataset. This is because the patients in PRIAS have already had their biopsies as per the PRIAS protocol. In addition, the true time of cancer progression is interval or right censored for all patients, making it impossible to correctly estimate the delay in detection of cancer progression due to a particular schedule. To this end, we conduct an extensive simulation study to find the utility of personalized, PRIAS, and fixed/heuristic schedules. For a realistic comparison, we simulate data from the joint model fitted to the PRIAS dataset. The simulated population has the same ten year follow-up period as the PRIAS study. In addition, the estimated relations between DRE and PSA measurements, and the risk of cancer progression are retained in the simulated population.

From this population, we first sample 500 datasets, each representing a hypothetical AS program with 1000 patients in it. We generate a true cancer progression time for each of the 500×1000 patients, and then

sample a set of DRE and PSA measurements at the same follow-up visit times as given in PRIAS protocol. We then split each dataset into a training (750 patients) and a test (250 patients) part, and generate a random and noninformative censoring time for the training patients. We next fit a joint model of the specification given in Equations (1), (2), and (3) to each of the 500 training datasets and obtain MCMC samples from the 500 sets of the posterior distribution of the parameters.

In each of the 500 hypothetical AS programs, we utilize the corresponding fitted joint models to develop cancer progression risk profiles for each of the 500×250 test patients. We make the decision of biopsies for patients at their pre-scheduled follow-up visits for DRE and PSA measurements (see **Study Population**), on the basis of their visit and patient-specific estimated cumulative risk of cancer progression. These decisions are made iteratively until a positive biopsy is observed. A recommended gap of one year between consecutive biopsies³ is also maintained. Subsequently, for each patient, an entire personalized schedule of biopsies is obtained.

We evaluate and compare both personalized and currently practiced schedules of biopsies in this simulation study. Comparison of the schedules is based on the number of biopsies scheduled and the corresponding delay in detection of cancer progression. We evaluate the following currently practiced fixed/heuristic schedules: biopsy annually, biopsy every one and a half years, biopsy every two years and biopsy every three years. We also evaluate the biopsy schedule of the PRIAS program (see **Introduction**). For the personalized biopsy schedules, we evaluate schedules based on three fixed risk thresholds: 5%, 10% and 15%, corresponding to a missed cancer progression being 19, 9, and 5.5 times more harmful than an unnecessary biopsy²¹, respectively. We also implement a personalized schedule where a follow-up visit-specific risk threshold is chosen using F_1 score.

Results

From the joint model fitted to the PRIAS dataset, we found that both $\log_2\{\text{PSA} + 1\}$ velocity, and \log odds of having $\text{DRE} > \text{T1c}$ were significantly associated with the hazard of cancer progression. For any patient, an increase in $\log_2\{\text{PSA} + 1\}$ velocity from -0.03 to 0.16 (first and third quartiles of the fitted velocities, respectively) corresponds to a 1.94

fold increase in the hazard of cancer progression. Whereas, an increase in odds of DRE > T1c from 0.001 to 0.01 (first and third quartiles of the fitted odds, respectively) corresponds to a 1.40 fold increase in the hazard of cancer progression. Detailed results pertaining to the fitted joint model are presented in Appendix B.

Comparison of Various Approaches for Biopsies

From the simulation study, we obtain the number of biopsies and the delay in detection of cancer progression for each of the 500×250 test patients using different schedules. The corresponding median number of biopsies and delay are shown in Figure 6. The personalized and PRIAS approaches fall in the region of better balance between the number of biopsies and the delay, than fixed/heuristic schedules. We next evaluate these schedules on the basis of both median and interquartile range (IQR) of the number of biopsies and delay. For brevity, only the most widely used annual and PRIAS schedules, the proposed personalized approach with fixed risk thresholds of 5% and 10%, and visit-specific threshold chosen using F_1 score are discussed next (see Appendix C for remaining).

Since patients have varying cancer progression speeds, the impact of each schedule also varies with it. In order to highlight these differences, we divide results for three types of patients, as per their time of cancer progression. They are *fast*, *intermediate*, and *slow progressing* patients. Although such a division may be imperfect and can only be done retrospectively in a simulation setting, we do it only for the purpose of illustration. Roughly 50% of the patients did not obtain cancer progression in the ten year follow-up period of the simulation study. We assume these patients to be *slow progressing* patients. We assume *fast progressing* patients are the ones with an initially misdiagnosed state of cancer²³, or high-risk patients who choose AS instead of immediate treatment upon diagnosis. These are roughly 30% of the population, having a cancer progression time less than 3.5 years. We label the remaining 20% patients as *intermediate progressing* patients.

The boxplots in Figure 7, show the variation in the number of biopsies, and the delay in detection of cancer progression, due to various biopsy schedules, for these three types of patients. For *fast progressing* patients (Panel A, Figure 7), we can see that the personalized schedules with a fixed 10% risk threshold and visit-specific threshold chosen using F_1 score,

reduce one biopsy for 50% of the patients, compared to PRIAS and annual schedule. Despite this, the delay due to personalized schedule with fixed 10% risk threshold (median: 0.7 years, IQR: 0.7 years), and the currently practiced annual (median: 0.6 years, IQR: 0.6 years) and PRIAS schedule (median: 0.7 years, IQR: 0.7 years), is similar.

For *intermediate progressing* patients (Panel A, Figure 7), we can see that the delay due to personalized schedule with fixed 5% risk threshold (median: 0.6 years, IQR: 0.6 years) is comparable to that of annual schedule (median 0.5 years, IQR: 0.5 years). However, it schedules fewer biopsies (median: 6, IQR: 2) than the annual schedule (median: 7, IQR: 3). The delay for PRIAS (median: 0.7 years, IQR: 1 year) and personalized schedule with fixed 10% risk (median: 0.7, IQR: 0.9) is similar, but the personalized approach schedules one less biopsy for 50% of the patients. Although the approach with visit-specific risk threshold chosen using F_1 score schedules fewer biopsies than the 10% fixed risk approach, but it also has a higher delay.

The patients who are at the most advantage with the personalized schedules are the *slow progressing* patients. These are a total of 50% patients who did not progress during the entire study. Hence, the delay is not available for these patients (Panel C of Figure 7). For all of these patients, annual schedule leads to 10 (unnecessary) biopsies. The PRIAS schedule, schedules a median of 6 biopsies (IQR: 4). In comparison, the biopsies scheduled by the personalized schedules using fixed 10% risk threshold (median: 4, IQR: 2) and visit-specific risk chosen using F_1 score (median: 2, IQR: 2), are much fewer.

Discussion

In this paper, we proposed a methodology for making personalized biopsy decisions for prostate cancer patients included in active surveillance (AS) programs. Our goal was to better balance the number of biopsies, and the delay in detection of cancer progression, than existing biopsy schedules. The proposed methodology combines a patient's observed DRE and PSA measurements, and the time of latest biopsy, into a cancer progression risk function. If the cumulative risk of cancer progression at a follow-up visit is above a certain threshold, then a biopsy is scheduled. We conducted a simulation study to compare the personalized approach for biopsies with the currently practiced fixed/heuristic biopsy schedules common for

all patients, and the PRIAS schedule (see [Introduction](#)). We found that the personalized schedule which uses 10% risk threshold at all follow-up visits is dominant over the PRIAS schedule, biennial schedule of biopsies, and biopsies every one and a half years. This personalized schedule not only schedules less number of biopsies than the aforementioned currently practiced schedules, but the delay in detection of cancer progression is also either equal or less. The personalized schedule which uses risk threshold chosen on the basis of classification accuracy (F_1 score) is dominant over the triennial schedule of biopsies. The personalized schedule which uses 5% risk threshold schedules less biopsies than the annual schedule, while the delay is only trivially more than the annual schedule. These results conclude that the personalized schedules better balance the number of biopsies, and the delay in detection of cancer progression than the currently practiced schedules.

The main reason for better performance of personalized schedules is that they account for the variation in cancer progression rate between patients, and also over time within the same patient. In contrast, the existing fixed/heuristic schedules ignore that roughly 50% patients never progress in the first ten years of follow-up (*slow progressing* patients) and do not require biopsies. While *fast progressing* patients require early detection, but existing methods of identifying these patients, such as the use of PSA doubling time in PRIAS, inappropriately assume that PSA evolves linearly over time. Thus, they may not correctly identify such patients. The personalized approach however, models the PSA profiles non-linearly. Furthermore, it appends information from PSA with information from DRE and previous biopsy results, and combines them into a cancer progression risk function. The risk function is a finer quantitative measure than individual data measurements observed for the patients. Risk is also easy to understand for both patients and doctors. Consequently, we expect that a more informed decision of biopsy can be made.

Existing work on mitigating the issue of non-compliance of biopsies and/or reducing the burden of biopsies in AS, primarily advocates less frequent schedule of biopsies. To our knowledge, risk based biopsy schedules have not been explored yet in AS⁹. However, the results that we estimated for fixed/heuristic schedules are comparable with corresponding results obtained in existing work⁶, even though the AS cohorts are not

same. Unless the cohorts differ significantly in patient population, the results for personalized schedules may also be valid in other cohorts.

A limitation of the personalized approach is that the choice of risk threshold is not straightforward, as different thresholds lead to different combinations of number of biopsies and delay in detection of cancer progression. An approach is to choose a risk threshold which leads to personalized schedule dominant over the currently practiced schedules, for a given delay. Although dominance may not guarantee compliance, it may be likely, because personalized schedules are less burdensome. A second limitation is that the results that we presented are valid only in a 10 year follow-up period, whereas prostate cancer is a slow progressing disease. Thus more detailed results, especially for *slow progressing* patients cannot be estimated. However, very few AS cohorts have a longer follow-period than PRIAS. Another limitation of risk based personalized schedules is that the biopsy decisions do not optimize for a reward/penalty function. While the risk itself can be interpreted as a ratio of immediate reward/penalty of biopsies²¹, but future rewards/penalties need to be considered as well.

There are multiple ways to extend the personalized decision making approach. One such way is to account for competing risks (see **Study Population**) in the joint model. Since biopsy Gleason grading is susceptible to inter-observer variation²⁴, accounting for it in our model will also be interesting to investigate further. To improve the decision making methodology, future consequences of a biopsy can be accounted in the model by combining Markov decision processes with joint models for time-to-event and longitudinal data. There is also potential for including diagnostic information from magnetic resonance imaging (MRI), and quality of life measures in our model. However, given the scarceness of such information in the dataset, including it in the current model may not be feasible. Lastly, we intend to further validate our results in a multi-center AS cohort, and subsequently develop a web-application to assist doctors in making personalized decisions for biopsies.

Acknowledgements

The first and last authors would like to acknowledge support by the Netherlands Organization for Scientific Research's VIDI grant nr. 016.146.301, and Erasmus MC funding. The authors also thank the Erasmus MC Cancer Computational Biology Center for giving access to their IT-infrastructure

and software that was used for the computations and data analysis in this study. Lastly, we thank Joost Van Rosmalen from the Department of Biostatistics, Erasmus University Medical Center, for feedback on the manuscript.

Supplemental material

Supplementary material for this article are available after references and figures in this document.

References

1. Torre LA, Bray F, Siegel RL et al. Global cancer statistics, 2012. *CA: A Cancer Journal for Clinicians* 2015; 65(2): 87–108.
2. Etzioni R, Penson DF, Legler JM et al. Overdiagnosis due to prostate-specific antigen screening: lessons from us prostate cancer incidence trends. *Journal of the National Cancer Institute* 2002; 94(13): 981–990.
3. Bokhorst LP, Alberts AR, Rannikko A et al. Compliance rates with the Prostate Cancer Research International Active Surveillance (PRIAS) protocol and disease reclassification in noncompliers. *European Urology* 2015; 68(5): 814–821.
4. Ehdaie B, Vertosick E, Spaliviero M et al. The impact of repeat biopsies on infectious complications in men with prostate cancer on active surveillance. *The Journal of urology* 2014; 191(3): 660–664.
5. Fujita K, Landis P, McNeil BK et al. Serial prostate biopsies are associated with an increased risk of erectile dysfunction in men with prostate cancer on active surveillance. *The Journal of urology* 2009; 182(6): 2664–2669.
6. Inoue LY, Lin DW, Newcomb LF et al. Comparative analysis of biopsy upgrading in four prostate cancer active surveillance cohorts. *Annals of internal medicine* 2018; 168(1): 1–9.
7. de Carvalho TM, Heijnsdijk EA and de Koning HJ. Estimating the risks and benefits of active surveillance protocols for prostate cancer: a microsimulation study. *BJU international* 2017; 119(4): 560–566.
8. Loeb S, Carter HB, Schwartz M et al. Heterogeneity in active surveillance protocols worldwide. *Reviews in urology* 2014; 16(4): 202.
9. Nieboer D, Tomer A, Rizopoulos D et al. Active surveillance: a review of risk-based, dynamic monitoring. *Translational andrology and urology* 2018; 7(1): 106.
10. Bokhorst LP, Valdagni R, Rannikko A et al. A decade of active surveillance in the PRIAS study: an update and evaluation of the criteria used to recommend a switch to active treatment. *European Urology* 2016; 70(6): 954–960.
11. Tsiatis AA and Davidian M. Joint modeling of longitudinal and time-to-event data: an overview. *Statistica Sinica* 2004; 14(3): 809–834.
12. Rizopoulos D. *Joint Models for Longitudinal and Time-to-Event Data: With Applications in R*. CRC Press, 2012. ISBN 9781439872864.
13. Turnbull BW. The empirical distribution function with arbitrarily grouped, censored and truncated data. *Journal of the Royal Statistical Society Series B (Methodological)* 1976; : 290–295.

14. Schröder F, Hermanek P, Denis L et al. The tnm classification of prostate cancer. *The Prostate* 1992; 21(S4): 129–138.
15. Pearson JD, Morrell CH, Landis PK et al. Mixed-effects regression models for studying the natural history of prostate disease. *Statistics in Medicine* 1994; 13(5-7): 587–601.
16. Lin H, McCulloch CE, Turnbull BW et al. A latent class mixed model for analysing biomarker trajectories with irregularly scheduled observations. *Statistics in Medicine* 2000; 19(10): 1303–1318.
17. De Boor C, De Boor C, Mathématicien EU et al. *A practical guide to splines*, volume 27. Springer-Verlag New York, 1978.
18. Eilers PH and Marx BD. Flexible smoothing with B-splines and penalties. *Statistical Science* 1996; 11(2): 89–121.
19. Rizopoulos D. The R package JMBayes for fitting joint models for longitudinal and time-to-event data using MCMC. *Journal of Statistical Software* 2016; 72(7): 1–46.
20. Rizopoulos D. Dynamic predictions and prospective accuracy in joint models for longitudinal and time-to-event data. *Biometrics* 2011; 67(3): 819–829.
21. Vickers AJ and Elkin EB. Decision curve analysis: a novel method for evaluating prediction models. *Medical Decision Making* 2006; 26(6): 565–574.
22. Rizopoulos D, Molenberghs G and Lesaffre EM. Dynamic predictions with time-dependent covariates in survival analysis using joint modeling and landmarking. *Biometrical Journal* 2017; 59(6): 1261–1276.
23. Cooperberg MR, Cowan JE, Hilton JF et al. Outcomes of active surveillance for men with intermediate-risk prostate cancer. *Journal of Clinical Oncology* 2011; 29(2): 228.
24. Carlson GD, Calvanese CB, Kahane H et al. Accuracy of biopsy Gleason scores from a large uropathology laboratory: use of a diagnostic protocol to minimize observer variability. *Urology* 1998; 51(4): 525–529.




Figure 5. Illustration of personalized decision of biopsy for patient j at two different follow-up visits. Biopsy is recommended if the personalized cumulative risk of cancer progression estimated from the joint model fitted to the observed data of the patient, is higher than the example risk threshold for biopsy ($\kappa = 10\%$). **Panel A:** biopsy is not recommended for the patient j at the follow-up visit time $s = 4$ years, because his estimated personalized cumulative risk of cancer progression (7.8%) is less than the threshold. **Panel B:** biopsy is recommended for the patient j at the follow-up visit time $s = 5.3$ years, because his estimated personalized cumulative risk of cancer progression (13.5%) is more than the threshold.




Figure 6. Simulation study results for burden-biopsy frontier: Estimated median number of biopsies, and median delay in detection of cancer progression, due to the currently practiced fixed/heuristic biopsy schedules (red squares) and PRIAS schedule (blue rhombus), and personalized schedules (green triangles), over a follow-up of ten years. **Types of personalized schedules:** Risk: 15%, Risk: 10% and Risk: 5% approaches, schedule biopsy if the cumulative risk of cancer progression at a visit is more than 15%, 10% and 5%, respectively. Risk: F1 works similar as previous, except that a visit-specific risk threshold is chosen by maximizing F₁ score (see [Methods](#)). The green shaded region depicts the region of better balance in number of biopsies and delay, than the currently practiced fixed/heuristic schedules. Estimation is based on results obtained from the simulation study we conducted.




Figure 7. Boxplot showing variation in the number of biopsies, and the delay in detection of cancer progression, in years (time of positive biopsy - true time of cancer progression) for various biopsy schedules. Biopsies are conducted until cancer progression is detected. **Panel A:** results for simulated patients who had a faster speed of cancer progression, with progression times between 0 and 3.5 years. **Panel B:** results for simulated patients who had an intermediate speed of cancer progression, with progression times between 3.5 and 10 years. **Panel C:** results for simulated patients who did not have cancer progression in the ten years of follow-up. **Types of personalized schedules:** Risk: 10% and Risk: 5% approaches, schedule biopsy if the cumulative risk of cancer progression at a visit is more than 10% and 5%, respectively. Risk: F1 works similar as previous, except that a visit-specific risk threshold is chosen by maximizing F_1 score (see [Methods](#)). Annual corresponds to a schedule of yearly biopsies and PRIAS corresponds to biopsies as per PRIAS protocol (see [Introduction](#)).