

HW1

Part A code:

```
% Zhiwei Wen
% EE102A HW1
% Problem 1
clear all; close all;
%% 1a
n = 0:10;
Omega0 = -pi/4;
xn = sin(Omega0*n);
figure;
stem (n,xn);
xlabel('Time n');
ylabel('x[n] = sin(\Omega_{0}n)');
title('Task 1, Part (a), \Omega_{0} = -\pi/4');
t = 0:.01:10;
xn_ct = sin(Omega0 * t);
figure;
hold on;
set(gca,'FontName','times','FontSize',16);
plot(t, xn_ct,'k--', 'LineWidth', 3);
stem(n, xn, 'b', 'LineWidth', 2);
xlabel('Time \itn');
ylabel('\itx\rm[\itn\rm] = sin(\Omega_{0}\itn\rm)');
title('Task 1, Part (a) with discrete and continuous n, \Omega_{0} = -\pi/4')
grid on;
%% 1b
Omega1 = 7*pi/4;
yn = sin(Omega1*n);
yn_ct = sin(Omega1 * t);
figure; hold on;
set(gca,'FontName','times','FontSize',16);
plot(t, xn_ct,'r--', 'LineWidth', 2.5, 'DisplayName', '\Omega = -\pi/4');
plot(t, yn_ct,'k--', 'LineWidth', 2.5, 'DisplayName', '\Omega = 7\pi/4');
stem(n, xn, 'r', 'LineWidth', 4, 'DisplayName', '\Omega = -\pi/4');
```

```

stem(n, yn, 'b', 'LineWidth', 4, 'DisplayName', '\Omega = 7\pi/4');
xlabel('Time \itn');
ylabel('\itx\rm(\itt\rm) = sin(\Omega\itn\rm)');
title('Task 1, Part (b) with discrete and continuous n')
legend('show');
grid on;
%% 1c
wn = xn.*xn;
figure;
set(gca, 'FontName', 'times', 'FontSize', 16);
stem(n, wn, 'LineWidth', 3);
xlabel('Time n');
ylabel('w[n]= x^{2}[n]');
grid on;
%periodic signal: T = 4.
%% 1d
t = 0:.005:5;
zt = exp(-t+1j*2*pi*t);
re_zt = real(zt);
im_zt = imag(zt);
envel1 = exp(-1*t);
envel2 = -exp(-1*t);
figure;
hold on;
set(gca, 'FontName', 'times', 'FontSize', 16);
plot(t, re_zt, 'r', 'LineWidth', 2.5, 'DisplayName', 'Real Part');
plot(t, im_zt, 'k', 'LineWidth', 2.5, 'DisplayName', 'Imaginary Part');
plot(t, envel1, 'b--', 'LineWidth', 1.5, 'DisplayName', 'Envelop e^{-t}');
plot(t, envel2, 'g--', 'LineWidth', 1.5, 'DisplayName', 'Envelop -e^{-t}');
xlabel('Time t');
title('Real and Imaginary parts of e^{-t+j2\pit}');
legend('show');
grid on;
% as can be seen in the figure, the real part and imaginary part oscillate
% within the envelop defined by +-e^{-t}. They are decaying sinusoids.
figure; hold on;


```

```


set(gca,'FontName','times','FontSize',16);
plot(re_zt, im_zt,'b', 'LineWidth', 4);
xlabel('Re\(\it{z}\rm(\it{t}\rm)'));
ylabel('Im\(\it{z}\rm(\it{t}\rm)'));
title('Task 1, Part (d), \(\it{z}\rm(\it{t}\rm) = \exp(-\it{t} + j2\pi\it{t})');
grid on;
% e^{-t+j*2*pi*t}=e^{-t}*e^{1j*2*pi*t};
% the real part and imaginary part oscillate
% within the envelop defined by +-e^{(-t)}.
% They are decaying sinusoids.

```


Part A figures:






Task 1, Part (a) with discrete and continuous n , $\Omega_0 = -\pi/4$



Task 1, Part (b) with discrete and continuous n







As can be seen in the figure of real and imaginary part, the real part and imaginary part oscillate within the envelop defined by $\pm e^{-t}$. They are decaying sinusoids. This is because in $e^{-t+1j*2*\pi*t} = e^{-t} * e^{1j*2*\pi*t}$, the coefficient of t in e^{-t+1} is -1 . This shows that the signal is decaying its module. the real part and imaginary part oscillate within the envelop defined by $\pm e^{-t}$. They are decaying sinusoids.

Part B codes:

```
% Zhiwei Wen
% EE102A HW1
% Problem 2
clear all; close all;
load guitar_note.mat;
t = (1:length(note))*dt;
figure; hold on;
plot(t(1:8:end),note(1:8:end),'DisplayName','Guitar Sound');
xlabel('time (s)');
ylabel('Amplitude');
title('Guitar note waveform');
alpha = 1.2;
```

```

sigma = -1;
t0 = 0.5;
% alpha is set to 1.2, sigma -1, t0 to 0.5.
envelope = alpha*exp(sigma*t).*double(t>=t0);
plot(t,envelope,'r','LineWidth',1.5,'DisplayName','envelop');
plot(t,-envelope,'r','LineWidth',1.5,'DisplayName','-envelop');
legend('show');grid on;
index_sample = 1*fs:round((1+1/16)*fs);
t_sample = t(index_sample);
note_sample = note(index_sample);
figure;
plot(t_sample,note_sample);
xlabel('sampled time(s)');
ylabel('Amplitude');
title('Sampled note waveform');
grid on;
% the cycle should be approximately 16.2 rounds. So the frequency
% should be 16.2 x 16 = 259.2 hz, which is very close to the real
% frequency of middle C : 261.62 hz.
f_note = 259.2;
simulated_note = envelope.*sin(2*pi*f_note*t);
figure;hold on;
plot(t(1:16:end),simulated_note(1:16:end),'DisplayName','Simulated Sound');
xlabel('time (s)');
ylabel('Amplitude');
title('Simulated note waveform');
envelope = alpha*exp(sigma*t).*double(t>=t0);
plot(t,envelope,'r','LineWidth',1.5,'DisplayName','envelop');
plot(t,-envelope,'r','LineWidth',1.5,'DisplayName','-envelop');
legend('show');grid on;
% By playing this sound with(simulated_note,fs) command in the command
% line, it is found that they are the same note with same fading rate.

```

Part B Figures:

