

Gateway Harmonized SHARE End of Life Documentation

Version G (2006-2022; using release 9.0.0), October 2024

Jenny Wilkens, Sarah Gao, Drystan Phillips,
& Jinkook Lee

Please cite use of dataset or documentation as:
Wilkens, Jenny, Sarah Gao, Drystan Phillips
& Jinkook Lee. Gateway Harmonized SHARE, Version G (using release 9.0.0).
Gateway to Global Aging Data, 2024.
<https://doi.org/10.25553/epgq-vv71>

We greatly appreciate support from the National Institute on Aging
(R01 AG030153, RC2 AG036619, R03 AG043052)

eol.g2aging.org

Preface

The Survey of Health, Ageing and Retirement in Europe (SHARE) is a longitudinal household survey for the study of retirement and health among the elderly in continental Europe. It is modeled on the Health and Retirement Study (HRS) and the English Longitudinal Study of Ageing (ELSA), which are similar longitudinal surveys in the United States and England. Part of the reason for the close connection is to allow cross-country comparisons using these data.

In order to make the data more accessible to researchers and to facilitate such comparisons, we, the USC Gateway to Global Aging team, created the Gateway Harmonized SHARE End of Life, a user-friendly version of a subset of data from the SHARE end of life interviews. The Gateway Harmonized SHARE End of Life is part of a larger initiative to provide Harmonized End of Life data for the HRS Family of Surveys. The Gateway creates and releases Harmonized End of Life datasets containing easily-merged and comparable research-ready variables. The variables are defined to be as comparable as possible to the core Harmonized dataset and across all Harmonized End of Life datasets. They use a similar intuitive naming convention as the core Harmonized dataset, with some modifications to avoid confusion with the core data. Harmonized End of Life datasets currently include the Gateway Harmonized SHARE End of Life, Harmonized HRS End of Life, Harmonized MHAS End of Life, Harmonized ELSA End of Life, Harmonized KLoSA End of Life, and Harmonized CHARLS End of Life, with more studies to be added in the near future. Further information about these Harmonized data files with questionnaires and other metadata is available on our searchable website, <https://eol.g2aging.org/>.

The Gateway Harmonized SHARE End of Life initiative is part of a larger set of projects. With funding and support from the National Institute on Aging, we have also created the Gateway Harmonized SHARE COVID and Gateway Harmonized SHARE Life History; Harmonized HRS, Harmonized HRS COVID, and Harmonized HRS-HCAP (United States); Harmonized MHAS and Harmonized Mex-Cog (Mexico); Harmonized ELSA, Harmonized ELSA COVID, Harmonized ELSA-HCAP, and Harmonized ELSA Life History (England); Harmonized CRELES (Costa Rica); Harmonized KLoSA (South Korea); Harmonized JSTAR (Japan); Harmonized TILDA (Ireland); Harmonized CHARLS and Harmonized CHARLS Life History (China); Harmonized ELSI (Brazil); Harmonized SPS (Chile); Harmonized LASI and Harmonized LASI-DAD (India); and Harmonized MARS (Malaysia) data. Further information about the Harmonized core, life history, and HCAP data files with questionnaires and other metadata is available on our searchable website, <https://g2aging.org/>. Further information about the Harmonized COVID data files with questionnaires and other metadata is available on our searchable COVID website, <https://covid.g2aging.org/>.

Note, however, that SHARE license agreements do not allow us to disseminate the data directly. Instead, SHARE distributes the Gateway Harmonized SHARE End of Life dataset in a separate download, which is made available on their website at <https://share-eric.eu/data/>. Additional information about SHARE can be obtained from SHARE's website at <https://share-eric.eu/>.

We are grateful for the continuing support of and funding from NIA. We also thank Axel Börsch-Supan, Giuseppe De Luca, Dimitris Christelis, Enrica Croda, Tim Birkenbach, Giacomo Pasini, and SHARE staff. We have greatly benefited from discussions with and the suggestions from our colleagues Jennifer Ailshire, Erik Meijer, Marco Angrisani, and Arie Kapteyn. We would also like to acknowledge our current and former colleagues at the Gateway to Global Aging Data: Sandy Chien, Astrid Ouyang, Kanghong Shao, and Yizhou Chen.

Requested Acknowledgment

We ask all users of the Gateway Harmonized SHARE End of Life to please inform our team of any written analysis using data from the Gateway Harmonized SHARE End of Life or information from the Gateway Harmonized SHARE End of Life Codebook by sending an email to papers@g2aging.org. We also ask users to include the following acknowledgement in their written work:

This analysis uses data or information from the Gateway Harmonized SHARE End of Life dataset and Codebook, Version G as of October 2024 (using release g.o.o) developed by the Gateway to Global Aging Data. The development of the Gateway Harmonized SHARE End of Life was funded by the National Institute on Aging (R01 AG030153, RC2 AG036619, R03 AG043052). For more information, please refer to <https://g2aging.org/>.

SHARE Version and Acknowledgment

This document uses data from SHARE Waves 1-9 release 9.0.0 as of March 2024. The SHARE data collection has been funded by the European Commission through the 5th framework programme (project QLK6-CT-2001-00360 in the thematic programme Quality of Life). Further support by the European Commission through the 6th framework programme (projects SHARE-I3, RII-CT-2006-062193, as an Integrated Infrastructure Initiative, COMPARE, CIT5-CT-2005-028857, as a project in Priority 7, Citizens and Governance in a Knowledge Based Society, and SHARE-LIFE (CIT4-CT-2006-028812)), through the 7th framework programme (SHARE-PREP (No 211909), SHARE-LEAP (No 227822), M4 (No 261982)), and DASISH (No 283646); through Horizon 2020 (SHAREDEV3 (No 676536), SERISS (No 654221), SSHOC (No 823782), SHARE-COHESION (No 870628), SHARE-COVID19 (No 101015924), Rltrain (No 654156) and ERIC Forum (No 823798)) and by DG Employment, Social Affairs & Inclusion through VS 2015/0195, VS 2016/0135, VS 2018/0285, VS2019/0332, VS2020/0313, SHARE-EUCOV: GA No 101052589 and EUCOVII: GA No 101102412 is gratefully acknowledged. Substantial co-funding for add-ons such as the intensive training and retention program for SHARE interviewers and the collection of HRS-harmonised biomarkers was granted by the US National Institute on Aging (U01 AG09740-13S2, P01 AG005842, P01 AG08291, P30 AG12815, R21 AG025169, Y1-AG-4553-01, IAG BSR06-11, OGHA 04-064, BSR12-04, and R01AG052527-02), further funding was granted for the development of a Harmonized Cognitive Assessment Protocol (HCAP) (R01 AG056329-02). Substantial funding for the central coordination of SHARE was received from the German Federal Ministry for Education and Research (Bundesministerium für Bildung und Forschung, BMBF) and the Max Planck Society for the Advancement of Science. See <https://share-eric.eu/infrastructure/funding> for a full list of funding institutions.

Contents

Preface	1
Requested Acknowledgment	1
SHARE Version and Acknowledgment	2
What's New in Version G of the Gateway Harmonized SHARE End of Life?	4
1. Introduction and Overview	5
1.1. Gateway to Global Aging Data	5
1.2. Units of Observation	6
1.3. Data File Structure	6
1.4. Variable Naming Convention	6
1.5. Missing Values, Nonresponse, and Imputations	7
1.6. Weighting and Accounting for Survey Design	7
1.7. Specifics to the End of Life Interview	7
1.8. Merging the End of Life Data with the Core Wave Data	7
2. Imputations	9
2.1. Currencies, Timing, and Exchange Rates	9
2.2. Imputations: Background	9
2.3. Imputation Process	10
3. Structure of Codebook	11
4. Distribution and Technical Notes	15
5. Data Codebook	16
A. Demographics	17
B. Health	50
C. Health Care Utilization and Insurance	81
D. Cognition	113
E. Financial and Housing Wealth	116
G. Family History	144
L. Assistance and Caregiving	146
O. End of Life Planning	200
6. References	214

What's New in Version G of the Gateway Harmonized SHARE End of Life?

Version G incorporates the latest released version of SHARE data, release 9.0.0, and adds observations from Wave 9 and from the Waves 8 and 9 COVID interviews. It contains 21,098 observations or rows. It is a Respondent-level file so each row represents a unique deceased Respondent. It also adds new variables and makes adjustments and corrections.

We have added the following variables to the file:

Demographics, Identifiers, and Weights:

- We added **isoa3**, which is a country-identifier.
- We added **raxiwmode**, which indicates the mode of the interview.
- We added **raxlang_s**, which indicates the language the interview took place in.
- We added harmonized gender and education variables from the Gateway Harmonized SHARE: **ragender**, **raeducel**, **raeduc1**, and **raedisced**.
- We added **inw8xtc** and **inw9xtc**, which indicate if the end of life interview took place as part of the core or COVID interview in Waves 8 and 9.
- We added **raxprxysp**, which indicates if the respondent's spouse or partner completed the end of life interview on the deceased respondent's behalf.
- We added **radlocs**, which is a simplified version of **radloc** to maximize comparability across Harmonized End of Life datasets.

Health:

- We added **ralhibpe**, which indicates whether the respondent had ever been diagnosed with high blood pressure as of their last core interview.

Healthcare Utilization and Insurance:

- We added **raxdoctor1y**, **raxdrugs1y**, **raxhomcar1y**, and **raxhsplc1m** for all available waves, which indicate having doctor visits, regularly taking medication, and receiving home care in the past year, and receiving hospice or palliative care in the last month, respectively.

We have made the following adjustments, improvements, and corrections to the data and documentation:

Demographics, Identifiers, and Weights:

- We adjusted **raxyyear** to assign special missing value (.i) if the reported death year was before the year of the respondent's last core interview.

Assistance and Caregiving:

- We replaced special missing value (.x) with 0 values for **raxracany1y**, **raxracaany1y**, **raxrafaany1y**, **raxricany1y_s**, **raxricaany1y_s**, **raxrifaany1y_s**, **raxrcany1y**, **raxrcany1y_s**, **raxrcaany1y**, **raxrcaany1y_s**, **raxrfaany1y**, and **raxrfaany1y_s** to maximize comparability with other Harmonized End of Life datasets.
- We fixed **raxrcaany1y_s** and **raxrfaany1y_s** in Wave 8, as a small percentage of respondents were mistakenly assigned special missing value (.m) instead of a 0 value for not receiving any help.

1. Introduction and Overview

This report documents the Gateway Harmonized SHARE End of Life data file, a streamlined collection of variables derived from the Survey of Health, Ageing, and Retirement in Europe (SHARE) end of life interviews. SHARE is a cross-national panel survey of individuals over age 50 and their spouses or partners in a growing number of countries in continental Europe. Its main goal is to provide panel data that enables research and analysis in support of policies on retirement, health insurance, saving, and economic well-being. The survey elicits information about demographics, income, assets, health, cognition, family structure and connections, health care use and costs, housing, job status and history, expectations, and insurance.

SHARE is sponsored by the European Commission, national funding agencies, and the (U.S.) National Institute of Aging (NIA), is administered by national survey agencies and universities, and is coordinated by the SHARE BERLIN Institute, and previously by the Munich Center for the Economics of Aging (MEA). The first wave was conducted in 2004-2005. This sampled individuals born in 1954 or earlier. In addition to respondents from eligible birth years, the survey interviewed respondents' spouses or partners regardless of age.

The data include any individual interviewed at least once in the core interview, and for whom a proxy completed an end of life interview. This includes individuals who were age-eligible (born in eligible years) at the time of their first interview, spouses who were not age-eligible at baseline, and spouses who married an age-eligible respondent between survey waves, and then subsequently passed away.

The most recent release of the SHARE data for Waves 1-9 is release 9.0.0 as of March 2024. Currently, Waves 1-2 and Waves 4-9 of the core interview are available for study. SHARE Wave 3 and a large sample in Wave 7 focused on the life histories of the respondents in the core interview, but conducted an end of life interview similar to other waves. Data and information from the end of life interviews in Waves 2-9 in addition to the core interview data from Waves 1-2, and 4-8 and both rounds of COVID interviews associated with Waves 8 and 9 are used and described in this document.

Documentation of the SHARE methodology can be found in Börsch-Supan et al. (2008) and Börsch-Supan and Jürges (2005). Documentation of the data can be found in SHARE (2024).

1.1. Gateway to Global Aging Data

The Health and Retirement Study (HRS) has achieved remarkable scientific success, as demonstrated by an impressive number of users, research studies, and publications using it. Its success has generated substantial interest in collecting similar data as population aging has progressed in every region of the world.

The result has been a number of surveys designed to be comparable with the HRS: the Mexican Health & Aging Survey (MHAS), the English Longitudinal Study of Ageing (ELSA), the Survey of Health, Ageing and Retirement in Europe (SHARE), the Costa Rican Longevity and Healthy Aging Study (CRELES), the Korean Longitudinal Study of Aging (KLoSA), the Japanese Study on Aging and Retirement (JSTAR), the Irish Longitudinal Study on Ageing (TILDA), the China Health and Retirement Longitudinal Study (CHARLS), the Brazilian Longitudinal Study of Ageing (ELSI), the Northern Ireland Cohort Longitudinal Study of Ageing (NICOLA), the Chilean Social Protection Survey (SPS), the Malaysia Ageing and Retirement Survey (MARS), and the Longitudinal Aging Study in India (LASI). The overview of this family of surveys, including their research designs, samples, and key domains can be found in Lee, Phillips, and Wilkens (2019).

As these surveys were designed with harmonization as a goal, they provide remarkable opportunities for cross-country studies. The value of comparative analyses, especially the opportunities they offer for learning lessons resulting from policies adopted elsewhere, is widely recognized. Yet there are only a limited number of empirical studies exploiting such opportunities. This is partly due to the difficulty associated with learning multiple surveys and the policies and institutions of each country.

Identifying comparable questions across surveys is the first step toward cross-country analyses. The Gateway to Global Aging Data (Gateway) helps users understand and use these large-scale population surveys on health and retirement. The Gateway includes several tools to facilitate cross-national health and retirement research. It includes a digital library of survey questions for all participating surveys. Its search engine enables users to find relevant survey questions. The Gateway also includes a concordance with information comparing measures within and across surveys over time. Using these

tools, researchers can identify all questions related to particular key words or within a domain. The Gateway also includes population and sub-population estimates for key harmonized variables and present them in graphs and tables that can be downloaded.

The Gateway to Global Aging End of Life Data can be accessed at <https://eol.g2aging.org>. The Gateway to Global Aging Data, with information on core interviews, can be accessed at <https://g2aging.org>. For more information about using the Gateway visit the Help page. For more information about obtaining the Gateway Harmonized SHARE End of Life using the Gateway see 'Chapter 4. Distribution and Technical Notes.'

1.2. Units of Observation

The Gateway Harmonized SHARE End of Life employs a single unit of observation, the individual. We do not distinguish between respondents and spouses, as is done in the Gateway Harmonized SHARE. Rather, all variables are respondent-level and can be merged with SHARE study data or the Gateway Harmonized SHARE using the id variable `mergeid`. We do not include non-respondents to the core interview or non-respondents to the end of life interview in our files.

1.3. Data File Structure

The Gateway Harmonized SHARE End of Life data are contained in a single file which includes the second through ninth waves of SHARE, from 2006 to 2022. The unit of observation is the individual. Each individual is uniquely identified by the unique identifier `mergeid`. This file may be merged with other SHARE data using `mergeid`.

1.4. Variable Naming Convention

Variable names in the Gateway Harmonized SHARE End of Life Data follow a consistent pattern. The first character indicates whether the variable refers to the reference person ('**r**'). The second character indicates that it pertains to 'all' ('**a**') waves, i.e., the variable is not specific to any single wave because the end of life interview is only given once for each deceased respondent. The remaining characters describe the concept that the variable captures. For example:

Variable `radmarrp` captures whether the respondent was married or partnered at the time of death. Because the respondent is deceased, the SHARE end of life interview is given exclusively to a proxy respondent. The relationship of the deceased respondent to the proxy is recorded in `raxprxy` and `raxprxysp`.

Variable labels also follow a consistent pattern. The first characters denote the name of the variable, followed by a colon. Then the remainder of the label describes the concept that the variable captures. For example, the variable label of `radmarrp` is:

radmarrp: r married/partnered at death

It may seem duplicative to include the name of the variable and the wave in the variable label. However, Stata often suppresses the variable name and instead uses its label in the presentation of results.

Variable names in the Gateway Harmonized SHARE End of Life are generally based on the variable name used in the Harmonized HRS End of Life for the same measure. Measures which are exactly or near-exactly comparable between the Gateway Harmonized SHARE End of Life and Harmonized HRS End of Life use the exact same name. For instance, `raxyear` is the variable name for the respondent's death year in both the Gateway Harmonized SHARE End of Life, as well as the Harmonized HRS End of Life. If the Gateway Harmonized SHARE End of Life measure is deemed only somewhat comparable with the Harmonized HRS End of Life version of that measure, the variable name in the Gateway Harmonized SHARE End of Life will often end in '`_s`'. This variable name suffix indicates some SHARE-specific difference with the Harmonized HRS End of Life version of this measure. Other reasons for Gateway Harmonized SHARE End of Life-specific variable names include: differences in survey routing, and whether both sets of variables use imputed values. Gateway Harmonized SHARE End of Life-specific variable names are used to notify the user that there are substantial differences between the Harmonized HRS End of Life and Gateway Harmonized SHARE End of Life measures and clean harmonization between these measures is not possible.

The Gateway Harmonized SHARE End of Life includes some variables without Gateway Harmonized SHARE End of Life-

specific variable names even though the Gateway Harmonized SHARE End of Life measure is significantly different from the Harmonized HRS End of Life measure of the same name. In particular, out-of-pocket expenditure and wealth measures in the Gateway Harmonized SHARE End of Life do not use Gateway Harmonized SHARE End of Life-specific variable names even though out-of-pocket expenditures and wealth measures in the Gateway Harmonized SHARE End of Life are expressed in 2010 Euros while these measures in the Harmonized HRS End of Life are always expressed in 2010 dollars. Users should always check the 'Differences with the Harmonized HRS End of Life' section of each measure before comparing any Gateway Harmonized SHARE End of Life measure to the Harmonized HRS End of Life version of the same measures or any other Harmonized Dataset version of the same measure.

1.5. Missing Values, Nonresponse, and Imputations

Variables may contain missing values for several reasons. Stata, SAS, and SPSS offer the capability to distinguish multiple types of missing values, and we have attempted to record as much information as possible. Generally, the codes adhere to the classification in Table 1.

Table 1. Missing Codes

Code	Reason for missing
.d	Don't know
.i	Invalid response
.m	Other missing
.n	Not assessed
.q	Question not asked in wave
.r	Refused
.x	Does not apply

The coding scheme varies across variables. Consult the Data Codebook for details on individual variables.

1.6. Weighting and Accounting for Survey Design

The Gateway Harmonized SHARE End of Life does not include weighting variables to account for non-response because none have been created and released by SHARE.

1.7. Specifics to the End of Life Interview

SHARE end of life interviews are conducted with a proxy informant for panel members who have died. The content of the end of life interviews is similar to the core interview, but end of life interviews are conducted to obtain information on the respondent's last year of life and the circumstances and cause of death. To the extent possible, proxy informants are selected to be in the close social network of the deceased respondent, and can be a family or household member, a neighbor, or any other person close to the respondent. About 84% of proxy responders are related to the respondent; 75% are a spouse, partner, or child of the respondent.

End of life interviews are attempted with a proxy in the first survey period following the death of the respondent and the majority of interviews occur within two years of death. End of life interviews began in the 2006 survey period (Wave 2) and have been conducted in every survey period since. As of the 2021-2022 survey year (Wave 9), end of life interviews have been completed for 21,098 SHARE respondents, which represent approximately 71% of all known decedents.

1.8. Merging the End of Life Data with the Core Wave Data

The Gateway Harmonized SHARE End of Life dataset can be easily merged with SHARE study data or the Gateway Harmonized SHARE using the unique SHARE id variable `mergeid`. Here we provide an example of Stata code to correctly merge the Gateway Harmonized SHARE dataset with the Gateway Harmonized SHARE End of Life dataset.

```
use "filepath/GH_SHARE_EOL_g.dta"
merge 1:1 mergeid using "filepath/GH_SHARE_g.dta"
```

Here we provide an example of Stata code to correctly merge select variables from the Gateway Harmonized SHARE dataset with select variables from the Gateway Harmonized SHARE End of Life dataset.

```
use variable1 variable2 variable3 using "filepath/GH_SHARE_EOL_g.dta"  
merge 1:1 mergeid using "filepath/GH_SHARE_g.dta", keepusing(variable4 variable5  
variable6)
```

In both cases, the same method can be used to merge the original SHARE study data if the name of the dataset is changed.

Because the time from death to the completion of the SHARE end of life interview varies by individual, care must be taken when merging wave-specific variables from the SHARE study data or Gateway Harmonized SHARE. In order to merge wave-specific variables from the last completed core wave, it is necessary to employ the use of **ralstcore**. Here we provide an example of Stata code to correctly assign whether the respondent had ever reported a diagnosis of cancer in their last completed core wave, assuming the SHARE end of life interview took place in Wave 9.

```
generate ralcancr = .  
foreach w in 1 2 4 5 6 7 8  
replace ralcancr = r`w'cancr if ralstcore==`w'
```

Here, we are generating a new variable called **ralcancr**, with the 'r' indicating the respondent, the 'l' indicating the last completed wave, and 'cancr' indicating ever receiving a cancer diagnosis. We employ the use of a foreach loop for Waves 1, 2, 4, 5, 6, 7, and 8 as it is possible for deceased respondents with an end of life interview in Wave 9 to have had their last core interview in any previous wave except Wave 3, which was a life history interview. Within the loops, we replace **ralcancr** to take the values of **rWcancr** from the Gateway Harmonized SHARE if the last completed core wave, **ralstcore**, is equal to 'w'.

2. Imputations

2.1. Currencies, Timing, and Exchange Rates

The SHARE survey uses several types of currency for monetary variables. In Wave 2 and forward proxy respondents in Euro countries had to report amounts in Euros. In countries that have not (or had not yet at the time of interview) adopted the Euro, proxy respondents gave amounts in their local currency. In the SHARE data files of the CAPI modules, these amounts have all been converted to Euros, using a fixed nominal exchange rate per country, i.e., an exchange rate not varying with the time of interview. This exchange rate is available in the SHARE imputations data file in the variable **nomxCAP1**. These currency conversions are performed using the exchange rate to Euros for the year the question referred to as provided by SHARE in the **nomxCAP1** variables.

2.2. Imputations: Background

Many SHARE end of life questions which ask about financial values follow a similar pattern. In the case of assets, proxies are first asked whether the deceased respondent had ownership of the asset, and if so, the proxy is asked the value of the asset. In the case of expenses, the proxy is asked the amount of the expense. If the proxy does not provide an exact value for the respondent's asset or expense, the SHARE end of life surveys the value using unfolding brackets, asking the proxy to identify ranges in which the value of the asset or expense lies. Starting in Wave 3, the SHARE end of life interview starts the unfolding bracket sequence by asking the proxy whether the value is less than, more than, or about equal to an amount randomly chosen from a predefined set of thresholds by country. Based on this response, the end of life interview asks up to twice more whether the value is less than, more than, or about equal to some of the other thresholds in an attempt to narrow the possible range of values. The result for some respondents is a closed bracket where the proxy identified a number lower than the value and one higher than the value. Another result for other respondents is an open bracket where the proxy only identified one number lower or higher than the value. One more result for other respondents is an approximate value where the proxy identified a single amount as about equal to the requested value. A final result for other respondents can be that the proxy was not able to identify any amount which was less than, more than, or about equal to the requested value.

As a result, the SHARE end of life data contains no-ownership/zero-value responses, exact amount reports, closed bracket responses, open bracket responses, cases where no bracket information was provided, and cases where ownership is unknown. A no-ownership/zero-value response results if the proxy reports the respondent not having/owning the expense/asset. An exact amount report results if the proxy reports the respondent having/owning the expense/asset and is able to identify the value. A closed bracket response results if the proxy reports the respondent having/owning the expense/asset and provides an approximate value or an upper and lower bound in which the value lies. An open bracket results if the proxy reports the respondent having/owning the expense/asset and provides some information but only a lower bound or upper bound is identified. Cases where we have no bracket information result if the proxy indicates the respondent having/owning the expense/asset, but neither a lower nor upper bound are identified. Finally, there may be proxies who refused to answer or did not know whether the respondent had/owned the expense/asset. These proxies were not asked to identify the value or taken into an unfolding bracket sequence, and their response is classified as "DK ownership" or "DK whether has expense".

In summary, the data contain valid responses (exact values and cases where we know the individual did not have this type of asset or expense) and several types of responses that require imputations. In decreasing order of the available information, those cases which require imputation:

Case 1: We may know an approximate value, this is considered similar to a closed bracket.

Case 2: We may know a number that the value is less than and a number that the value is more than, this is a closed bracket.

Case 3: We may know only one number that the value is more than or less than, this is an open bracket.

Case 4: We may know that the individual owned the asset or had the expense, but have no information on its value.

Case 5: We may not even know whether the individual owned an asset or had the expense.

For all variables which include imputed values, the Gateway Harmonized SHARE End of Life provides and flags imputed values in separate variables.

2.3. Imputation Process

In principle, imputations should use the conditional distribution of the variable to be imputed conditional on all observed variables. In practice, however, this is impractical, undesirable, and often impossible, especially if the data set has large numbers of variables, as does the SHARE, which would lead to overfitting and inability to estimate all the coefficients in the models. On the other hand, using a very simplified imputation procedure like a hot deck imputation which does not take into account any observed variables or only a few observed variables (in a conditional hot deck) can lead to a match bias, where the imputation method tends to impute too many values in the middle of the distribution (Hirsch & Schumacher 2004 and Bollinger & Hirsch 2006). Taking into consideration the advantages and disadvantages of different imputation procedures, the Gateway Harmonized SHARE End of Life provides imputed values using a predictive mean matching imputation method (PMM; Little 1988) with a small number of covariates.

For all of the five cases identified above, Gateway Harmonized SHARE End of Life imputes the value using the PMM method. The Gateway Harmonized SHARE End of Life PMM method uses all reported values from all waves, treating 'no ownership' as a zero value, and estimates a linear regression model for the inverse hyperbolic sine of this value with a small set of covariates by country including wave dummies. The imputed value is then the reported value of the individual with the closest predicted value, where the donor pool consists of the individuals who reported a value that is consistent with the reported information from the individual that needs imputation. For instance, if a proxy reported the respondent having a value between 10,000 and 25,000 in the unfolding bracket sequence, the donor pool would consist of individuals whose proxy reported a value between 10,000 and 25,000. If the donor pool consists of fewer than 50 individuals, a simple hotdeck imputation is performed instead. In these cases, the imputed value is taken from a randomly selected individual who reported a value that is consistent with the reported information from the individual that needs imputation. The selection of the individual is performed without consideration of the closeness of predicted values resulting from the linear regression model, that is, it is random.

There are also two cases under which missing values are not imputed. The first case occurs when the donor pool consists of fewer than 2 individuals. For instance, a proxy can report that an asset amount was above value X but there are no or only one record with a value above X, and this donor pool would be considered insufficient for imputation. The second case occurs when the individual with a missing value is also missing a value for one of the small set of covariates used by the imputation procedure to predict values. For instance, if age is used as a covariate and an individual with missing age is also missing the value of an asset, the imputation procedure is not able to predict which other individuals would have the closest value. In both these cases, the values are not imputed but instead are assigned a special missing (.m).

As part of the imputation process, imputation flag variables are also created, which allow data users to know whether the value was reported, left missing and for what reason, or imputed and, if imputed, what information was known (regarding ownership and bracket values). In addition to the value, ownership is also taken from the donor individual, so both are jointly imputed. (Note that SHARE end of life allows ownership with a zero value, so the two are not equivalent.) For more information about this PMM model refer to Lee, Meijer, and Phillips (2015).

When imputing individual-level values, the gender and age at death of the individual are used as covariates, and all imputations are conducted separately by country.

We have chosen to remove any reported values greater than 10 million Euro. We have also chosen to remove any reported values in which a grouping of 5 or more 9's are reported, such as 99999 or 999999, because we assume that these are don't know or refused values entered incorrectly. The values for these respondents in both cases are imputed with positive ownership and without any bracket information.

The Gateway Harmonized SHARE End of Life provides imputed values for individual medical expenditures, home value, estate value, and life insurance value. All of the financial variables in the Gateway Harmonized SHARE End of Life, whether representing assets or expenditures, have already been adjusted by the annual consumer price index (`cYYYYcpindex`) of the Euro based on the year of the respondent's death. All values in the SHARE end of life data had previously been converted to Euros by the SHARE team. For more information on the calculation of the consumer price index see <http://stats.oecd.org>.

3. Structure of Codebook

The Data Codebook contains the codebook documenting all variables in the Gateway Harmonized SHARE End of Life Data. This section explains how to interpret the codebook entries. The figure below shows a typical codebook page; the numbers in blue correspond to comments below.

1	Life Insurance Policies: Whether Had Life Insurance and Value			
	Variable	Waves	Label	Type
2	raxlifein	All	raxlifein: r any life insurance policies	Categ
	raxlifeinv	All	raxlifeinv: r life insurance: total value of policies	Cont
	raxlifeinvf	All	raxlifeinvf: r life insurance flag: total value of policies	Categ

3 How Constructed

raxlifein indicates whether the deceased respondent owned any life insurance policies. **raxlifein** is coded as 0 if the respondent did not own any life insurance policies. **raxlifein** is coded as 1 if the respondent did own a life insurance policy. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

raxlifeinv has been adjusted to 2010 Euros based on the consumer price index for the year and country of death.

raxlifeinv indicates the total value of all life insurance policies owned by the respondent and includes imputed values. If the respondent owned any life insurance policies, then the proxy is asked the approximate total value of all life insurance policies owned by the respondent. The value of the life insurance policies is elicited in Euros in all SHARE countries except in Denmark, Sweden, Switzerland, Poland, Czech Republic, and Israel where local currencies are used and which have been converted to Euros by the SHARE team; however, all reported or imputed values in **raxlifeinv** have been converted and adjusted to 2010 Euros based on the consumer price index for the year and country of death. **raxlifeinv** is assigned a value of 0 if the respondent did not own any life insurance policies. If the proxy answers don't know or refuses to answer the value of the life insurance policies, the proxy is not asked any additional questions to obtain threshold values; instead values are imputed with no bracket information. **raxlifeinv** is assigned special missing (.m) if the values are not able to be imputed due to a lack of reference observations in the donor pool or missing covariates necessary for the imputation procedure.

raxlifeinvf is a flag variable indicating the level of imputation for the total value of life insurance policies. A code of 1 indicates the proxy reported a continuous value and no imputation was necessary. A code of 5 indicates that the value was imputed without any bracket information. A code of 6 indicates the proxy reported the respondent not having any life insurance policies and the value is 0. A code of 7 indicates that the proxy was not sure whether the respondent had any life insurance policies. A code of -1 indicates that the value could not be imputed due to a lack of reference observations in the donor pool. A code of -2 indicates that the expense could not be imputed due to one or more missing values in the covariates used in the imputation procedure.

4 Cross Wave Differences in SHARE EOL

No differences known.

5 Differences with the Harmonized HRS End of Life

raxlifeins indicates whether anyone received a settlement from the deceased respondent's life insurance in the Harmonized HRS End of Life, whereas **raxlifein** indicates whether the respondent owned any life insurance policies in the Gateway Harmonized SHARE End of Life. While the two concepts are very similar, it is possible to own a life insurance policy and to ultimately not receive a settlement from it, as such we have provided different variable names for these two concepts.

In the HRS exit interview, the proxy is asked what amounts or percentages that each beneficiary received from the deceased's life insurance policy. Depending on the wave, proxies might also be asked the total amount of life insurance received altogether. In the SHARE end of life interview, on the other hand, proxies are asked a single direct question asking for the total of the deceased's owned life insurance policies.

In the HRS exit interview, if the proxy responds that they don't know or refuse to answer the value of the life insurance settlement, then a series of unfolding bracket questions is asked to obtain a minimum and maximum value of the settlement. However, in the SHARE end of life interview, no unfolding bracket questions are asked to the proxy if they respond that they don't know or refuse to obtain the approximate value of the life insurance policies.

The monetary values in the Harmonized HRS End of Life are provided in 2010 U.S. dollars, whereas monetary values in the Gateway Harmonized SHARE End of Life are provided in 2010 Euro. As such, adjustment must be made before any comparison of these variables.

Continuous Variable Distributions

6

Variable	N	Mean	Std.Dev	Minimum	Maximum	Total missing
raxlifeinv	21,026	2,576.06	39,686.41	0.00	2,042,900.88	72

Categorical Variable Frequencies

7

	raxlifein
0.no	17,368
1.yes	1,917
.d.dk	1,335
.m:missing	219
.r:refuse	259
Total	21,098
	raxlifeinvf
-2.not imputed, missing cov	67
-1.not imputed, missing nei	5
1.continuous value	1,235
5.no bracket info	677
6.no asset	17,368
7.ownership unknown	1,746
Total	21,098

Original SHARE Variables Used

8

Wave 2 End of Life:

xt033_	Deceased owned any life insurance policies
xt034e	Value of all life insurance policies

Wave 3 End of Life:

sl_xt033_	The deceased owned any life insurance policies
-----------	--

sl_xto34e	Value of all life insurance policies
Wave 4 End of Life:	
xto33_	Deceased owned any life insurance policies
xto34e	Value of all life insurance policies
Wave 5 End of Life:	
xto33_	Deceased owned any life insurance policies
xto34e	Value of all life insurance policies
Wave 6 End of Life:	
xto33_	Deceased owned any life insurance policies
xto34e	Value of all life insurance policies
Wave 7 End of Life:	
xto33_	Deceased owned any life insurance policies
xto34e	Value of all life insurance policies
Wave 8 End of Life:	
xto33_	Deceased owned any life insurance policies
xto34e	Value of all life insurance policies
Wave 9 End of Life:	
xto33_	Deceased owned any life insurance policies
xto34e	Value of all life insurance policies
Wave 9 COVID End of Life:	
xto33_	Deceased owned any life insurance policies
xto34e	Value of all life insurance policies

- 1 **Title:** The variables are documented in groups according to the concept that they measure. For example, there are three variables related to having life insurance policies, corresponding to all waves and the respondent.
- 2 **Variable Overview:** This entry overviews the provided variables in this group. The table includes the names of variables. The wave column displays "All" for variables which are not wave-specific (most variables), and otherwise displays the waves for which this variable is built (includes a "W" in place of individual wave numbers). Variable labels are also included which typically include the name of the variable and a short description. Lastly, the Type column indicates the variable type, which may be continuous (Cont), categorical (Categ), or character (Char).
- 3 **How Constructed:** This entry provides background on the manner in which variables were constructed.
- 4 **Cross-Wave Differences in SHARE EOL:** This entry briefly describes differences in question wording or contents between interview waves.
- 5 **Differences with the Harmonized HRS End of Life:** This entry describes any differences between the Harmonized HRS End of Life version of the variable and the Gateway Harmonized SHARE End of Life version of the variable. It is imperative these differences are understood when using harmonized measures.
- 6 **Continuous Variable Distributions:** This entry shows descriptive statistics of each continuous variable. They include the number of nonmissing values, the mean, standard deviation, minimum value, maximum value, and number of missing values in the wave.
- 7 **Categorical Variable Frequencies:** This entry shows the value label codes. These are only relevant for categorical variables. The first character(s) of the value labels indicate the value to which each label has been assigned. For example, value '1' is mapped into '1. Yes' (not just 'Yes'). The entry also indicates which labels are assigned to which variables, and shows frequency tabulations for all categorical variables.

8

Original SHARE Variables Used: This entry provides the names and labels of raw SHARE variables that were used to construct the new variables.

4. Distribution and Technical Notes

The Gateway Harmonized SHARE End of Life Data file is distributed through the SHARE website along with the original SHARE data. The Gateway Harmonized SHARE End of Life data file is made available free of charge but only to users who register with the SHARE and agree to the standard conditions. For more information on obtaining access to the SHARE data, visit: <https://share-eric.eu/data/data-access>.

The Gateway Harmonized SHARE End of Life is distributed in Stata, SPSS, and R dataset formats.

This is version **G** using release 9.0.0 of the Gateway Harmonized SHARE End of Life Data.

A copy of this Gateway Harmonized SHARE Codebook can be obtained on the Gateway to Global Aging Data End of Life website under the Harmonized End of Life Data and Get Data tabs (https://eol.g2aging.org/hrd/get_data).

5. Data Codebook

A. Demographics

Person Specific Identifier			
Variable	Waves	Label	Type
hhid	All	hhid: household id /6-char	Char
pn	All	pn: Person number within household (char)	Char
mergeid	All	Person identifier (fix across modules and waves)	Char

How Constructed

SHARE provides a unique identifier for each respondent across waves, **mergeid**. It consists of three separate parts:

A two character country/language survey code derived from the first part of **mergeid**. These two characters are different from **country** because three countries, Switzerland, Belgium, and Israel, use a different **cid** for different survey languages. The values are as follows: 11.Austria, 12.Germany, 13.Sweden, 14.Netherlands, 15.Spain, 16.Italy, 17.France, 18.Denmark, 19.Greece, 20.Switzerland (German), 21.Switzerland (French), 22.Switzerland (Italian), 23.Belgium (French), 24.Belgium (Flemish), 25.Israel (Hebrew), 26.Israel (Arabic), 27.Israel (Russian), 28.Czech Republic, 29.Poland, 30.Ireland, 31.Luxembourg, 32.Hungary, 33.Portugal, 34.Slovenia, 35.Estonia, 36.Croatia, 37.Lithuania, 38.Bulgaria, 39.Cyprus, 40.Finland, 41.Latvia, 42.Malta, 43.Romania, and 44.Slovakia.

hhid is a 6 character household identifier used in SHARE to indicate the household to which a sample participant belonged when entering the panel. Note that **hhid** is different from SHARE's wave-specific household identifier **hhidw**, which indicates the household where the respondent resides in a particular wave *w*.

pn is the 2 digit respondent identifier used by SHARE to indicate each participant within the household.

mergeid is SHARE's 12 character unique and non-changing identifier across waves. **mergeid** is formed as **cid-hhid-pn**. For example, if a respondent is from Austria and has a **cid** of 11, is in a household with **hhid** 123456, and has a **pn** of 02 inside the household, their **mergeid** would be "11-123456-02".

mergeid is used for all merges between waves since it is the only identifier that tracks respondents across waves and through the end of life interview. **mergeid** is a SHARE specific identifier and can be used for merging additional information from SHARE.

Cross Wave Differences in SHARE EOL

Starting in Wave 5, SHARE began specifying whether surveys were conducted in Spain/Girona in Catalan or Castilian, in Luxembourg in French or German, and in Estonia in Estonian or Russian.

Differences with the Harmonized HRS End of Life

No differences known.

Original SHARE Variables Used

Harmonized SHARE:

hhid	hhid: household id /6-char
pn	pn: Person number within household (char)
mergeid	Person identifier (fix across modules and waves)

Variable	Waves	Country	
		Label	Type
country	All	Country identifier	Categ
isocountry	All	UN numerical country code	Categ
isoa3	All	isoa3: Country indicator	Char

How Constructed

country is the SHARE country identifier. **country** is a 2-digit code ranging from 11 to 47 which represents the country in which the respondent had been previously surveyed. The codes are mapped as follows: 11.Austria, 12.Germany, 13.Sweden, 14.Netherlands, 15.Spain, 16.Italy, 17.France, 18.Denmark, 19.Greece, 20.Switzerland, 23.Belgium, 25.Israel, 28.Czech Republic, 29.Poland, 30.Ireland, 31.Luxembourg, 32.Hungary, 33.Portugal, 34.Slovenia, 35.Estonia, and 47.Croatia. **country** is helpful for a variety of purposes because the core survey and end of life questionnaires varied among the countries that participated in SHARE.

isocountry is a modified version of the SHARE variable **country**. Using ISO 3166-1 (numeric-3) codes, **isocountry** indicates the country in which surveying took place. The codes are mapped as follows: 40.Austria, 56.Belgium, 100.Bulgaria, 191.Croatia, 196.Cyprus, 203.Czech Republic, 208.Denmark, 233.Estonia, 246.Finland, 250.France, 276.Germany, 300.Greece, 348.Hungary, 372.Ireland, 376.Israel, 380.Italy, 428.Latvia, 440.Lithuania, 442.Luxembourg, 470.Malta, 528.Netherlands, 616.Poland, 620.Portugal, 642.Romania, 703.Slovakia, 705.Slovenia, 724.Spain, 752.Sweden, and 756.Switzerland. **isocountry** is helpful for matching respondents to country-level statistics where countries are often identified using ISO codes.

isoa3 are three-letter country codes which indicate the country in which surveying took place. The codes are mapped as follows: AUT-Austria, BEL-Belgium, BGR-Bulgaria, HRV-Croatia, CYP-Cyprus, CZE-Czech Republic, DNK-Denmark, EST-Estonia, FIN-Finland, FRA-France, DEU-Germany, GRC-Greece, HUN-Hungary, IRL-Ireland, ISR-Israel, ITA-Italy, LVA-Latvia, LTU-Lithuania, LUX-Luxembourg, MLT-Malta, NLD-Netherlands, POL-Poland, PRT-Portugal, ROU-Romania, SVK-Slovakia, SVN-Slovenia, ESP-Spain, SWE-Sweden, and CHE-Switzerland. **isoa3** is helpful for matching respondents to country-level statistics where countries are often identified using ISO codes.

Cross Wave Differences in SHARE EOL

No differences known.

Differences with the Harmonized HRS End of Life

No differences known.

Categorical Variable Frequencies

	country
Austria	1,034
Germany	699
Sweden	1,265
Netherlands	506
Spain	1,909
Italy	1,425
France	972
Denmark	1,003
Greece	1,286
Switzerland	491
Belgium	1,232

Section A. Demographics

Israel	896
Czech Republic	1,537
Poland	1,191
Ireland	36
Luxembourg	104
Hungary	706
Portugal	368
Slovenia	893
Estonia	1,959
Croatia	457
Lithuania	202
Bulgaria	180
Cyprus	81
Finland	53
Latvia	152
Malta	81
Romania	314
Slovakia	66
Total	21,098
isocountry	
040.Austria	1,034
056.Belgium	1,232
100.Bulgaria	180
191.Croatia	457
196.Cyprus	81
203.Czech Republic	1,537
208.Denmark	1,003
233.Estonia	1,959
246.Finland	53
250.France	972
276.Germany	699
300.Greece	1,286
348.Hungary	706
372.Ireland	36
376.Israel	896
380.Italy	1,425
428.Latvia	152
440.Lithuania	202
442.Luxembourg	104
470.Malta	81
528.Netherlands	506
616.Poland	1,191
620.Portugal	368
642.Romania	314
703.Slovakia	66
705.Slovenia	893
724.Spain	1,909
752.Sweden	1,265
756.Switzerland	491
Total	21,098

Original SHARE Variables Used

Section A. Demographics

Harmonized SHARE:

country	Country identifier
isocountry	UN numerical country code
isoa3	isoa3: Country indicator

Wave Status: End of Life Interview			
Variable	Waves	Label	Type
inxt	All	inxt: r in eol interview	Categ
raxt	All	raxt: r eol interview wave	Cont
inwWxt	2-9	inwWxt: r in wW eol interview	Categ
inwWxtc	8-9	inwWxtc: r type of eol interview in wW	Categ

How Constructed

inxt indicates whether a proxy completed the end of life interview for the deceased respondent in any wave. **inxt** is coded as 1 if the respondent is deceased and a proxy completed the end of life interview on their behalf in any available wave. All respondents in this dataset are coded as 1 because this dataset only includes respondents who are deceased and have had a completed end of life interview.

raxt indicates in which wave a proxy completed the first exit interview for the deceased respondent. **raxt** can take values from 2 through 9, which corresponds to a value of 1 for **inwWxt** in the given wave, i.e. if **inw2xt** is equal to 1, then **raxt** will have a value of 2.

inwWxt indicates whether a proxy completed the end of life interview for the deceased respondent in the specified wave. **inwWxt** is coded as 1 if a proxy completed the end of life interview on behalf of the deceased respondent in the specified wave. In Waves 8 and 9, **inwWxt** is coded as 1 if the end of life interview was completed either in the core or COVID interview attributed to that wave. **inwWxt** is coded as 0 if a proxy completed the end of life interview on behalf of the deceased respondent as part of another wave. Please note that this dataset only includes respondents who are deceased and have had a completed end of life interview. **inwWxt** does not indicate whether the respondent is alive or dead, but rather which wave the respondent's end of life interview took place in.

inwWxtc indicates whether the respondent's end of life interview was completed during the core interview or the COVID interview attributed to the particular wave, and only applies to Waves 8 and 9. **inw8xtc** is coded as 1 if the respondent's end of life interview was completed during the Wave 8 core interview and coded as 2 if the respondent's end of life interview was completed during the first round of COVID data collection in the summer of 2020. **inw9xtc** is coded as 1 if the respondent's end of life interview was completed during the Wave 9 core interview and coded as 2 if the respondent's end of life interview was completed during the second round of COVID data collection in the summer of 2021.

Cross Wave Differences in SHARE EOL

The end of life interviews for Waves 2 through 7 were collected during the fielding of the core interview of that wave. The end of life interviews for Wave 8 were collected during the fielding of the core interview of Wave 8 and the fielding of the first round of COVID interviews in 2020. The end of life interviews for Wave 9 were collected during the fielding of the core interview of Wave 9 and the fielding of the second round of COVID interviews in 2021.

Differences with the Harmonized HRS End of Life

The HRS conducts exit interviews, post-exit interviews, and post post-exit interviews. The Harmonized HRS End of Life has comparable variables to **inwWxt** indicating the wave in which the respondent's first exit interview took place. The Harmonized HRS End of Life has additional variables indicating the wave(s) in which the respondent's post-exit or post post-exit interviews took place. The SHARE only conducts a single end of life interview, which is used to assign values to **inwWxt** in the Gateway Harmonized SHARE End of Life. The Gateway Harmonized SHARE End of Life includes **inwWxtc** to indicate whether the end of life interview was completed as part of the core or COVID interview for that wave, which is not a needed variable for the HRS.

Continuous Variable Distributions

Variable	N	Mean	Std.Dev	Minimum	Maximum	Total missing
raxt	21,098	6.71	1.96	2.00	9.00	0

Categorical Variable Frequencies

	inxt			
1.resp,deceased	21,098			
Total	21,098			
	inw2xt	inw3xt	inw4xt	inw5xt
0.nonresp	20,372	19,891	19,964	18,911
1.resp,deceased	726	1,207	1,134	2,187
Total	21,098	21,098	21,098	21,098
	inw6xt	inw7xt	inw8xt	inw9xt
0.nonresp	17,752	17,441	16,931	16,424
1.resp,deceased	3,346	3,657	4,167	4,674
Total	21,098	21,098	21,098	21,098
	inw8xtc	inw9xtc		
0.nonresp	16,931	16,424		
1.core iww	3,048	3,472		
2.covid iww	1,119	1,202		
Total	21,098	21,098		

Original SHARE Variables Used

Tracker:

interview_w2	Interview done (wave 2)
interview_w3	Interview done (wave 3)
interview_w4	Interview done (wave 4)
interview_w5	Interview done (wave 5)
interview_w6	Interview done (wave 6)
interview_w7	Interview done (wave 7)
interview_w8	Interview done (wave 8)
interview_w8ca	Interview done (wave 8: COVID-19 Survey 1)
interview_w9	Interview done (wave 9)
interview_w9ca	Interview done (wave 9: COVID-19 Survey 2)

End of Life Interview Date: Month and Year

Variable	Waves	Label	Type
raxtiwm	All	raxtiwm: r eol interview month	Cont
raxtiwy	All	raxtiwy: r eol interview year	Cont

How Constructed

raxtiwm and **raxtiwy** indicate the month and year, respectively, in which the respondent's end of life interview took place. Data are from the Cross-Waves Tracker file. End of life interviews have been conducted since SHARE Wave 2 for respondents who had participated in at least one previous wave of the SHARE. Don't know, refuse, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Cross Wave Differences in SHARE EOL

No differences known.

Differences with the Harmonized HRS End of Life

No differences known.

Continuous Variable Distributions

Variable	N	Mean	Std.Dev	Minimum	Maximum	Total missing
raxtiwm	21,067	5.85	3.08	1.00	12.00	31
raxtiwy	21,067	2,016.74	4.17	2,006.00	2,022.00	31

Original SHARE Variables Used

Tracker:

int_month_w2	Interview month (wave 2)
int_year_w2	Interview year (wave 2)
int_month_w3	Interview month (wave 3)
int_year_w3	Interview year (wave 3)
int_month_w4	Interview month (wave 4)
int_year_w4	Interview year (wave 4)
int_month_w5	Interview month (wave 5)
int_year_w5	Interview year (wave 5)
int_month_w6	Interview month (wave 6)
int_year_w6	Interview year (wave 6)
int_month_w7	Interview month (wave 7)
int_year_w7	Interview year (wave 7)
int_month_w8	Interview month (wave 8)
int_year_w8	Interview year (wave 8)
int_month_w8ca	Interview month (wave 8: COVID-19 Survey 1)
int_year_w8ca	Interview year (wave 8: COVID-19 Survey 1)
int_month_w9	Interview month (wave 9)
int_year_w9	Interview year (wave 9)
int_month_w9ca	Interview month (wave 9: COVID-19 Survey 2)
int_year_w9ca	Interview year (wave 9: COVID-19 Survey 2)

Variable	Waves	Last Completed Core Interview	
		Label	Type
ralstcore	All	ralstcore: r last completed core interview wave	Cont
ralstcorey	All	ralstcorey: r last completed core interview year	Cont

How Constructed

ralstcore indicates the most recent core wave that the now deceased respondent completed while alive, including if the last completed wave was completed by a proxy. This variable is determined based on the **inwW** variables in the Gateway Harmonized SHARE. Because SHARE Wave 3 was a life history interview and not a core interview, **ralstcore** cannot take the value of 3. **ralstcore** will take a value of 7 if the last interview the respondent completed was either the full questionnaire or the combined condensed questionnaire and life history questionnaire in Wave 7.

ralstcorey indicates the year of the most recent core wave interview that the now deceased respondent completed while alive, including if the last completed wave was completed by a proxy. This variable is determined based on the **inwW** and **rWiwy** variables in the Gateway Harmonized SHARE.

Cross Wave Differences in SHARE EOL

Because SHARE Wave 3 was a life history interview and not a core interview, **ralstcore** cannot be equal to 3.

Differences with the Harmonized HRS End of Life

No differences known.

Continuous Variable Distributions

Variable	N	Mean	Std.Dev	Minimum	Maximum	Total missing
ralstcore	21,098	5.06	2.17	1.00	8.00	0
ralstcorey	21,098	2,013.13	4.54	2,004.00	2,020.00	0

Original SHARE Variables Used

Harmonized SHARE:

inw1	inw1: =1 if respondent w1
inw2	inw2: =1 if respondent w2
inw4	inw4: =1 if respondent w4
inw5	inw5: =1 if respondent w5
inw6	inw6: =1 if respondent w6
inw7	inw7: =1 if respondent w7
inw7c	inw7c: =1 if respondent w7 condensed version
inw8	inw8: =1 if respondent w8
inw9	inw9: =1 if respondent w9
r1iwy	r1iwy:w1 r interview year
r2iwy	r2iwy:w2 r interview year
r4iwy	r4iwy:w4 r interview year
r5iwy	r5iwy:w5 r interview year
r6iwy	r6iwy:w6 r interview year
r7iwy	r7iwy:w7 r interview year
r8iwy	r8iwy:w8 r interview year
r9iwy	r9iwy:w9 r interview year

Relationship of Proxy to Respondent

Variable	Waves	Label	Type
raxprxy	All	raxprxy: proxy relationship to r: eol iww	Categ
raxprxysp	All	raxprxysp: spouse is r's proxy: eol iww	Categ

How Constructed

raxprxy indicates the relationship of the proxy who completed the end of life interview to the deceased respondent, as reported by the proxy. **raxprxy** is coded as follows: 1.spouse, 2.child, 3.other relative, 4.other non-relative. **raxprxy** is coded as 1 if the proxy is the deceased respondent's spouse or partner. **raxprxy** is coded as 2 if the proxy is the deceased respondent's son or daughter, son- or daughter-in-law, or son or daughter of husband/wife/partner. **raxprxy** is coded as 3 if the proxy is the deceased respondent's grandchild, sibling, or other relative. **raxprxy** is coded as 4 if the proxy is another non-relative. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

raxprxysp indicates if the proxy who completed the end of life interview is the deceased respondent's spouse. **raxprxysp** is coded as 0 if the proxy is someone other than the deceased respondent's spouse and coded as 1 if the proxy is the deceased respondent's spouse. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Cross Wave Differences in SHARE EOL

No differences known.

Differences with the Harmonized HRS End of Life

In the HRS exit interview, the relationship of the proxy to the deceased respondent is reported by the interviewer. In the SHARE end of life interview, the relationship of the proxy to the deceased respondent is reported by the proxy. Additionally, the HRS exit interview and SHARE end of life interview have different answer categories for indicating how the proxy is related to the deceased respondent, though this does not impact the harmonized categories for **raxprxy** or **raxprxysp**.

Categorical Variable Frequencies

	raxprxy
1.spouse	8,652
2.child	7,102
3.other relative	2,037
4.other non-relative	3,259
.d:dk	2
.m:missing	44
.r:refuse	2
Total	21,098
	raxprxysp
0.non-spouse	12,398
1.spouse	8,652
.d:dk	2
.m:missing	44
.r:refuse	2
Total	21,098

Original SHARE Variables Used

Wave 2 End of Life: xt002_	Relationship to the deceased
Wave 3 End of Life: sl_xt002_	Relationship to the deceased
Wave 4 End of Life: xt002_	Relationship to the deceased
Wave 5 End of Life: xt002_	Relationship to the deceased
Wave 6 End of Life: xt002_	Relationship to the deceased
Wave 7 End of Life: xt002_	Relationship to the deceased
Wave 8 End of Life: xt002_	Relationship to the deceased
Wave 9 End of Life: xt002_	Relationship to the deceased
Wave 9 COVID End of Life: xt002_	Relationship to the deceased

Interview Mode			
Variable	Waves	Label	Type
raxiwmode	All	raxiwmode: r interview mode	Categ

How Constructed

raxiwmode indicates the mode of the end of life interview. **raxiwmode** is coded as follows: 1.face to face, 2.telephone. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Cross Wave Differences in SHARE EOL

No differences known.

Differences with the Harmonized HRS End of Life

A comparable variable will be available in the next version of the Harmonized HRS End of Life. The HRS also conducts web interviews in some recent waves.

Categorical Variable Frequencies

	raxiwmode
1.face to face	14,738
2.telephone	6,226
.d:dk	7
.m:missing	125
.r:refuse	2
Total	21,098

Original SHARE Variables Used

Wave 2 End of Life:	
xto43_	Interview mode
Wave 3 End of Life:	
sl_xto43_	Interview mode
Wave 4 End of Life:	
xto43_	Interview mode
Wave 5 End of Life:	
xto43_	Interview mode
Wave 6 End of Life:	
xto43_	Interview mode
Wave 7 End of Life:	
xto43_	Interview mode
Wave 8 End of Life:	
xto43_	Interview mode
Wave 9 End of Life:	
xto43_	Interview mode
Wave 9 COVID End of Life:	
xto43_	Interview mode

Variable	Waves	Interview Language		Type
		Label		
raxlang_s	All	raxlang_s: r interview language		Categ

How Constructed

raxlang_s indicates the language used for the interview. **raxlang_s** is coded as follows: 1.German, 2.Swedish, 3.Dutch, 4.Spanish, 5.Italian, 6.French, 7.Danish, 8.Greek, 9.Flemish, 10.Hebrew, 11.Arabic, 12.Russian, 13.Czech, 14.Polish, 15.English, 16.Hungarian, 17.Portuguese, 18.Slovenian, 19.Estonian, 20.Catalan, 21.Croatian, 22.Luxembourgian, French, German, or Portuguese, 23.Estonian or Russian, 24.Spanish or Catalan, 25.French or German, 26.French, German, or Portuguese, 27.Lithuanian, 28. Bulgarian, 29. Finnish, 30. Latvian, 31. Maltese, 32. Romanian, 33. Slovakian.

Cross Wave Differences in SHARE EOL

No differences known.

Differences with the Harmonized HRS End of Life

A comparable variable will be available in the next version of the Harmonized HRS End of Life. The HRS allows interviews in English and Spanish.

Categorical Variable Frequencies

	raxlang_s
1.German	2,135
2.Swedish	1,269
3.Dutch	506
4.Spanish	1,659
5.Italian	1,451
6.French	1,715
7.Danish	1,003
8.Greek	1,367
9.Flemish	638
10.Hebrew	659
11.Arabic	116
12.Russian	387
13.Czech	1,537
14.Polish	1,191
15.English	44
16.Hungarian	706
17.Portuguese	368
18.Slovenian	893
19.Estonian	1,003
20.Catalan	116
21.Croatian	457
22.Luxembourgian,French,Ger	18
23.Estonian or Russian	699
24.Spanish or Catalan	134
27.Lithuanian	202
28.Bulgarian	180
29.Finnish	49
30.Latvian	143
31.Maltese	73

32.Romanian	314
33.Slovakian	66
Total	21,098

Original SHARE Variables Used

Wave 2 End of Life:	
language_xt	Language: End of life interview
Wave 3 End of Life:	
language_xt	Language: End of life interview
Wave 4 End of Life:	
language_xt	Language: End of life interview
Wave 5 End of Life:	
language_xt	Language: End of life interview
Wave 6 End of Life:	
language_xt	Language: End of life interview
Wave 7 End of Life:	
language_xt	Language: End of life interview
Wave 8 End of Life:	
language_xt	Language: End of life interview
Wave 9 End of Life:	
language_xt	Language: End of life interview
Wave 9 COVID End of Life:	
language_xt	Language: End of life interview

Gender			
Variable	Waves	Label	Type
ragender	All	ragender: r gender	Categ

How Constructed

ragender is derived using the gender variable in the SHARE tracker file. If this is not available, then **ragender** takes the gender reported in the Demographics module. **ragender** is coded as 1.man and 2.woman.

Cross Wave Differences in SHARE

No differences known.

Differences with the Harmonized HRS End of Life

No differences known.

Categorical Variable Frequencies

	ragender
1.man	11,235
2.woman	9,863
Total	21,098

Original SHARE Variables Used

Tracker:
gender Male or female

Education			
Variable	Waves	Label	Type
raeducel	All	raeducel: r harmonized early education level	Categ
raeduc1	All	raeduc1: r harmonized education level	Categ
raedisced	All	raedisced: r education by isced code	Categ

How Constructed

raeducel and **raeduc1** identify the level of education completed according to five-tier and three-tier harmonized scales which we developed to compare education levels across countries. These Harmonized education scales are simplified versions of 1997 International Standard Classification of Education (ISCED-97) codes. For more information on ISCED codes, see www.uis.unesco.org and the OECD document entitled "Classifying Educational Programmes: Manual for ISCED-97 Implementation in OECD Countries, 1999 Edition". **raeducel** is coded as follows: 1.Less than primary education, 2.Primary education, 3.Lower secondary education, 4.Upper secondary & vocational training, and 5.Tertiary education. **raeduc1** is coded as follows: 1.Less than upper secondary education, 2.Upper secondary & vocational training, and 3.Tertiary education. These variables are built using **raedisced**. Less than primary education is equivalent to no education. Less than upper secondary education is equivalent to no education, primary education, and lower secondary education. Upper secondary and vocational training is equivalent to upper secondary and post-secondary non-tertiary education. Tertiary education is equivalent to the first and second stage of tertiary education. Don't know, refused, or otherwise missing responses are assigned special missing values (.d), (.r), (.m), respectively.

Additionally, in order to provide a harmonized measure of education across all countries included in SHARE, the survey categorized education achievements using the 1997 International Standard Classification of Education ISCED-97 codes. Respondents were asked to report their highest school degree and degrees of further education or vocational training as in their country's educational system. SHARE then asked educational experts in each country to map local educational achievements to ISCED codes. For more information on ISCED codes see www.uis.unesco.org.

raedisced is constructed by looking at reports from all waves of data. The first non-missing report of education level by ISCED code is always used. In all waves SHARE provides imputations for ISCED codes. For more information on the imputations see **implicat**. **raedisced** is coded as follows: 0.None, 1.Primary education, 2.Lower secondary education, 3.Upper secondary education, 4.Post-secondary non-tertiary education, 5.First stage of tertiary education, 6.Second stage of tertiary education. Don't know, refused, or otherwise missing responses are assigned special missing value (.d), (.r), (.m), respectively. **raedisced** is set to plain missing (.) for respondents who did not respond to any waves.

Cross Wave Differences in SHARE

After Wave 1, educational degree questions are asked of new respondents only.

ISCED code 6 (second stage of tertiary education) was not asked in all countries.

Differences with the Harmonized HRS End of Life

raeduc1 is included in the Harmonized HRS, and **raeducel** will be included in its next release.

The HRS does not include an education variable coded according to the ISCED categories.

Categorical Variable Frequencies

	raeducel
1.less than upper secondary	2,043
2.primary education	6,338
3.lower secondary education	4,008
4.upper secondary & vocatio	6,157

Section A. Demographics

5.tertiary education	2,552
Total	21,098
raeduc1	
1.less than upper secondary	12,389
2.upper secondary and vocat	6,157
3.tertiary education	2,552
Total	21,098
raedisced	
0.None	2,043
1.Primary education	6,338
2.Lower secondary education	4,008
3.Upper secondary education	5,409
4.Post-secondary non tertia	748
5.First stage of tertiary e	2,480
6.Second stage of tertiary	72
Total	21,098

Original SHARE Variables Used

Harmonized SHARE:

raeducel	raeducel: r harmonized early education level
raeduc1	raeduc1: r harmonized education level
raedisced	raedisced: r education by isced code

Date of Death: Month and Year			
Variable	Waves	Label	Type
raxmonth	All	raxmonth: r death month in eol ivw	Cont
raxyear	All	raxyear: r death year in eol ivw	Cont

How Constructed

raxmonth and **raxyear** indicate the deceased respondent's month and year of death, respectively, as reported by the proxy in the end of life interview. **raxyear** is assigned special missing value (.i) if the proxy reported a year of death beyond the end of life interview date or before the respondent's last core interview. **raxmonth** is assigned special missing (.i) if the proxy reported a month of death beyond the end of life interview date when the death and interview occurred in the same year. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Cross Wave Differences in SHARE EOL

No differences known.

Differences with the Harmonized HRS End of Life

No differences known.

Continuous Variable Distributions

Variable	N	Mean	Std.Dev	Minimum	Maximum	Total missing
raxmonth	20,666	6.34	3.57	1.00	12.00	432
raxyear	20,753	2,015.26	4.36	2,004.00	2,022.00	345

Original SHARE Variables Used

Wave 2 End of Life:

xt008_ Month of decease
xt009_ Year of decease

Wave 3 End of Life:

sl_xt008_ Month of decease
sl_xt009_ Year of decease

Wave 4 End of Life:

xt008_ Month of decease
xt009_ Year of decease

Wave 5 End of Life:

xt008_ Month of decease
xt009_ Year of decease

Wave 6 End of Life:

xt008_ Month of decease
xt009_ Year of decease

Wave 7 End of Life:

xt008_ Month of decease
xt009_ Year of decease

Wave 8 End of Life:

xt008_ Month of decease

Section A. Demographics

xto09_	Year of decease
Wave 9 End of Life:	
xto08_	Month of decease
xto09_	Year of decease
Wave 9 COVID End of Life:	
xto08_	Month of decease
xto09_	Year of decease

Age at Death

Variable	Waves	Label	Type
radage	All	radage: r age at death	Cont
radagef	All	radagef: r flag age at death	Categ

How Constructed

radage indicates the deceased respondent's age at death. **radage** takes the respondent's age as reported by the proxy where possible. If this value is missing, then **radage** is assigned the age at death as calculated in the SHARE All Waves Coverscreen (Tracker) file for Waves 2 through 9. If both these values are missing, then **radage** is calculated using the year and month of birth and the year and month of death. If the birth or death month is missing, then the birth and death years are used to calculate age at death. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

radagef is a flag variable indicating whether an age was reported, or whether year and month or only years are used in the calculation of **radage**. **radagef** is coded as follows: 0.reported age used, 1.month and year used, 2.only year used. **radagef** is assigned special missing (.m) if **radage** has a missing value.

Cross Wave Differences in SHARE EOL

No differences known.

Differences with the Harmonized HRS End of Life

If the respondent's age at death is not provided by the survey, the Harmonized HRS End of Life calculates age at death by incorporating the years of birth and death in all cases, and the months of birth and death where possible. In the Gateway Harmonized SHARE End of Life, if the respondent's age at death is not provided by the proxy, **radage** then takes the value of death age provided in the SHARE All Waves Coverscreen (Tracker) file when available, then calculates age at death by incorporating the years of birth and death in all cases, and the months of birth and death where possible.

Continuous Variable Distributions

Variable	N	Mean	Std.Dev	Minimum	Maximum	Total missing
radage	21,020	79.43	10.23	20.00	107.00	78

Categorical Variable Frequencies

	radagef
0.reported age used	20,765
1.month & year used	204
2.only year used	51
.m:missing	78
Total	21,098

Original SHARE Variables Used

Tracker:

deceased_age Age at decease

Wave 2 End of Life:

xt008_ Month of decease

Section A. Demographics

xt009_	Year of decease
xt010_	Age at the moment of decease
Wave 3 End of Life:	
sl_xt008_	Month of decease
sl_xt009_	Year of decease
sl_xt010_	Age at the moment of decease
Wave 4 End of Life:	
xt008_	Month of decease
xt009_	Year of decease
xt010_	Age at the moment of decease
Wave 5 End of Life:	
xt008_	Month of decease
xt009_	Year of decease
xt010_	Age at the moment of decease
Wave 6 End of Life:	
xt008_	Month of decease
xt009_	Year of decease
xt010_	Age at the moment of decease
Wave 7 End of Life:	
xt008_	Month of decease
xt009_	Year of decease
xt010_	Age at the moment of decease
Wave 8 End of Life:	
xt008_	Month of decease
xt009_	Year of decease
xt010_	Age at the moment of decease
Wave 9 End of Life:	
xt008_	Month of decease
xt009_	Year of decease
xt010_	Age at the moment of decease
Wave 9 COVID End of Life:	
xt008_	Month of decease
xt009_	Year of decease
xt010_	Age at the moment of decease

Time from Death to Interview: Months and Years			
Variable	Waves	Label	Type
radtoivwm	All	radtoivwm: r time between death and eol iww, months	Cont
radtoivwy	All	radtoivwy: r time between death and eol iww, years	Cont
radtoivwf	All	radtoivwf: r flag time between death and eol iww	Categ

How Constructed

radtoivwm and **radtoivwy** indicate the calculated months and years, respectively, from the respondent's death to the end of life interview. **radtoivwm** is calculated using the year and month of death and the year and month of the end of life interview. If end of life interview month or death month is missing, then the calculation only considers the end of life interview and death year values. **radtoivwy** is calculated by dividing **radtoivwm** by 12. If these calculations create a negative value or if death year is considered an invalid value, then **radtoivwm** and **radtoivwy** are assigned special missing (.i). Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

radtoivwf is a flag variable indicating whether year and month or only years are used in the calculation of **radtoivwm** and **radtoivwy**. **radtoivwf** is coded as follows: 1.month and year used, 2.only year used. **radtoivwf** is assigned special missing (.m) if **radtoivwm** or **radtoivwy** have a missing value.

Cross Wave Differences in SHARE EOL

No differences known.

Differences with the Harmonized HRS End of Life

No differences known.

Continuous Variable Distributions

Variable	N	Mean	Std.Dev	Minimum	Maximum	Total missing
radtoivwm	20,722	17.11	15.25	0.00	146.00	376
radtoivwy	20,722	1.43	1.27	0.00	12.17	376

Categorical Variable Frequencies

	radtoivwf
1.month & year used	20,424
2.only year used	298
.m:missing	376
Total	21,098

Original SHARE Variables Used

Tracker:

int_month_w2	Interview month (wave 2)
int_year_w2	Interview year (wave 2)
int_month_w3	Interview month (wave 3)
int_year_w3	Interview year (wave 3)
int_month_w4	Interview month (wave 4)

Section A. Demographics

int_year_w4	Interview year (wave 4)
int_month_w5	Interview month (wave 5)
int_year_w5	Interview year (wave 5)
int_month_w6	Interview month (wave 6)
int_year_w6	Interview year (wave 6)
int_month_w7	Interview month (wave 7)
int_year_w7	Interview year (wave 7)
int_month_w8	Interview month (wave 8)
int_year_w8	Interview year (wave 8)
int_month_w9	Interview month (wave 9)
int_year_w9	Interview year (wave 9)
int_month_w9ca	Interview month (wave 9: COVID-19 Survey 2)
int_year_w9ca	Interview year (wave 9: COVID-19 Survey 2)
Wave 2 End of Life:	
xt008_	Month of decease
xt009_	Year of decease
Wave 3 End of Life:	
sl_xt008_	Month of decease
sl_xt009_	Year of decease
Wave 4 End of Life:	
xt008_	Month of decease
xt009_	Year of decease
Wave 5 End of Life:	
xt008_	Month of decease
xt009_	Year of decease
Wave 6 End of Life:	
xt008_	Month of decease
xt009_	Year of decease
Wave 7 End of Life:	
xt008_	Month of decease
xt009_	Year of decease
Wave 8 End of Life:	
xt008_	Month of decease
xt009_	Year of decease
Wave 9 End of Life:	
xt008_	Month of decease
xt009_	Year of decease
Wave 9 COVID End of Life:	
xt008_	Month of decease
xt009_	Year of decease

Location of Death			
Variable	Waves	Label	Type
radloc_s	All	radloc_s: r specific death location	Categ
radloc	All	radloc: r death location	Categ
radlocs	All	radlocs: r death location, simplified	Categ

How Constructed

radloc_s is a more specific version of **radloc**, and indicates the respondent's location at the time of their death, as reported by the proxy. **radloc_s** is coded as follows: 1.own home, 2.another person's home, 3.hospital, 4.nursing home, 5.hospice, 6.residential home/sheltered housing, 7.in transit/other. **radloc_s** is coded as 1 if the proxy reports that the respondent died in their own home. **radloc_s** is coded as 2 if the proxy reports that the respondent died at another person's home. **radloc_s** is coded as 3 if the proxy reports that the respondent died at the hospital. **radloc_s** is coded as 4 if the proxy reports that the respondent died in a nursing home. **radloc_s** is coded as 5 if the proxy reports that the respondent died in a hospice. **radloc_s** is coded as 6 if the proxy reports that the respondent died in a residential home or sheltered housing. In Waves 3 to 6, **radloc_s** is coded as 7 if the proxy reports that the respondent died in some other place. Starting in Wave 7, **radloc_s** is coded as 7 if the proxy reports that the respondent died in transit to a medical facility or in some other place. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

radloc indicates the respondent's location at the time of their death, as reported by the proxy. **radloc** is coded as follows: 1.private home, 2.hospital, 3.nursing home, 4.hospice, 5.other. **radloc** is coded as 1 if the proxy reports that the respondent died in their own home or at another person's home. **radloc** is coded as 2 if the proxy reports that the respondent died at the hospital. **radloc** is coded as 3 if the proxy reports that the respondent died in a nursing home, residential home, or sheltered housing. **radloc** is coded as 4 if the proxy reports that the respondent died in a hospice. In Waves 3 to 6, **radloc** is coded as 5 if the proxy reports that the respondent died in some other place. Starting in Wave 7, **radloc** is coded as 5 if the proxy reports that the respondent died in transit to a medical facility or in some other place. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

radlocs is a simplified version of **radloc** and indicates the respondent's location at the time of their death. **radlocs** is coded as follows: 1.private home, 2.hospital or institution, 3.other. **radlocs** is coded as 1 if the proxy reports that the respondent died in their own home or at another person's home. **radlocs** is coded as 2 if the proxy reports that the respondent died at the hospital, in a nursing home, residential home, sheltered housing, or in a hospice. In Waves 3 to 6, **radlocs** is coded as 3 if the proxy reports that the respondent died in some other place. Starting in Wave 7, **radlocs** is coded as 3 if the proxy reports that the respondent died in transit to a medical facility or in some other place. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Cross Wave Differences in SHARE EOL

The option of "some other place" is available beginning in Wave 3. The option of "in transit to a medical facility" is available beginning in Wave 7.

Differences with the Harmonized HRS End of Life

The HRS exit interview and SHARE end of life interview have slightly different answer categories for indicating where the respondent died. As such, **radloc** and **radlocs** in the Gateway Harmonized SHARE End of Life are coded as 1.private home if the respondent died at their own home or at another person's home, whereas these variables in the Harmonized HRS End of Life are coded as 1 if the respondent died at home, presumably their own home. **radloc** in the Gateway Harmonized SHARE End of Life is coded as 3.nursing home if the respondent died at a nursing home, residential home, or sheltered housing, whereas **radloc** in the Harmonized HRS End of Life is coded as 3 if the respondent died at a nursing home, assisted living facility, rest home, retirement home, or senior care home. **radloc** in the Gateway Harmonized SHARE End of Life is coded as 5 if the respondent died in another location or while in transit to a medical facility, whereas **radloc** in the Harmonized

HRS End of Life is coded as 5 if the respondent died in another location. These same differences between nursing homes and other locations also apply to **radlocs** in the Gateway Harmonized SHARE End of Life and Harmonized HRS End of Life.

Categorical Variable Frequencies

	radloc_s
1.own home	6,741
2.another person's home	263
3.hospital	9,991
4.nursing home	2,036
5.hospice	521
6.residential home/sheltere	561
7.in transit/other	417
.d:dk	110
.m:missing	426
.r:refuse	32
Total	21,098

	radloc
1.private home	7,004
2.hospital	9,991
3.nursing home	2,594
4.hospice	524
5.other	417
.d:dk	110
.m:missing	426
.r:refuse	32
Total	21,098

	radlocs
1.private home	7,004
2.hospital or institution	13,109
3.other	417
.d:dk	110
.m:missing	426
.r:refuse	32
Total	21,098

Original SHARE Variables Used

Wave 2 End of Life:	
xt014_	Place of dying
Wave 3 End of Life:	
sl_xt014_	Place of dying
Wave 4 End of Life:	
xt014_	Place of dying
Wave 5 End of Life:	
xt014_	Place of dying
Wave 6 End of Life:	
xt014_	Place of dying
Wave 7 End of Life:	

Section A. Demographics

xt014_	Place of dying
Wave 8 End of Life:	
xt014_	Place of dying
Wave 9 End of Life:	
xt014_	Place of dying
Wave 9 COVID End of Life:	
xt014_	Place of dying

Main Cause of Death			
Variable	Waves	Label	Type
racod_s	All	racod_s: disease that caused r's death	Categ
ragcod	All	ragcod: grouped disease that caused r's death	Categ

How Constructed

racod_s indicates the respondent's specific main cause of death. **racod_s** is coded as follows: 1.cancer, 2.heart attack, 3.stroke, 4.other cardiovascular related illness, such as heart failure or arrhythmia, 5.respiratory disease, 6.disease of the digestive system, such as gastrointestinal ulcer or inflammatory bowel disease, 7.severe infectious disease, such as pneumonia, septicemia or flu, 8.accident, 9.parasitic diseases, 10.endocrine, nutritional and metabolic diseases, 11.mental and behavioral disorders, 12.diseases of the nervous system, 13.diseases of the skin and subcutaneous tissue, 14.diseases of the musculoskeletal system and connective tissue, 15.diseases of the genitourinary system, 16.suicide/self-harm, 17.decrepitude/dotage/senility, 18.autoimmune disorders/diseases, 19.multiple morbidities, 20.multiple organ failure/failure of vital parts, 21.murder/killed in combat, 22.consequences of hip fracture, 23.died during hospitalization, 24.COVID-19, 25.other. Please note that codes 9 through 25 are not answer choices in the SHARE end of life interview, rather when the proxy responds "other" when asked the respondent's main cause of death, the proxy is asked to explain the main cause of death. The answers from this question have been grouped by the SHARE team and coded in **racod_s**. **racod_s** is assigned a code of 25 if the proxy responded "other" when asked for the respondent's main cause of death, and the explanation given was not codable or there was no explanation given. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

ragcod indicates the respondent's grouped main cause of death. **ragcod** is coded as follows: 1.cancer, 2.cardiovascular, 3.other. **ragcod** is coded as 1 if the proxy reports that the respondent died as a result of cancer. **ragcod** is coded as 2 if the proxy reports that the respondent died as a result of a heart attack, stroke, or other cardiovascular related illness such as heart failure or arrhythmia. **ragcod** is coded as 3 if the proxy reports that the respondent died as a result of a respiratory disease, disease of the digestive system such as gastrointestinal ulcer or inflammatory bowel disease, severe infectious disease such as pneumonia, septicemia or flu, accident, parasitic disease, endocrine, nutritional and metabolic disease, mental and behavioral disorder, disease of the nervous system, disease of the skin and subcutaneous tissue, disease of the musculoskeletal system and connective tissue, disease of the genitourinary system, suicide/self-harm, decrepitude/dotage/senility, autoimmune disorder/disease, multiple morbidities, multiple organ failure/failure of vital parts, murder/killed in combat, consequences of hip fracture, died during hospitalization, COVID-19, or other. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Cross Wave Differences in SHARE EOL

Starting in Wave 8, COVID-19 is provided as an option for the cause of death by SHARE.

Differences with the Harmonized HRS End of Life

The HRS exit interview has more answer categories for main cause of death than the SHARE end of life interview in their respective questionnaires. The original answer categories, as released in the studies' respective datasets, are presented in **racod_h** and **racod_s** for the Harmonized HRS End of Life and Gateway Harmonized SHARE End of Life, respectively. These answer categories have been grouped to provide comparable measures of cause of death in **ragcod**. Furthermore, the HRS exit interview allows the proxy to list up to 2 main causes of death, though only the first one is considered in **ragcod** and **racod_h**, while the SHARE end of life interview asks the proxy to report only 1 main cause of death.

Categorical Variable Frequencies

	racod_s
1.cancer	5,189
2.heart attack	2,729

Section A. Demographics

3.stroke	2,296
4.other cardiovascular rela	3,046
5.respiratory disease	1,093
6.disease of the digestive	544
7.severe infectious disease	1,108
8.accident	502
9.parasitic diseases	11
10.endocrine, nutritional,	149
11.mental and behavioral di	395
12.dz of nervous system	271
13.dz of skin, subcutaneous	7
14.dz of musculoskeletal sy	32
15.dz of genitourinary syst	202
16.suicide/self-harm	97
17.decrepitude/dotage/senil	1,442
18.autoimmune disorder/dz	13
19.multiple morbidities	214
20.multiple organ failure/f	189
21.murder/killed in combat	8
22.consequences of hip frac	58
23.died during hospitalisat	90
24.COVID-19	541
25.other	250
.d:dk	437
.m:missing	123
.r:refuse	62
Total	21,098
ragcod	
1.cancer	5,189
2.cardiovascular	8,071
3.other	7,216
.d:dk	437
.m:missing	123
.r:refuse	62
Total	21,098

Original SHARE Variables Used

Wave 2 End of Life:

xt011_	Main cause of death
xt012c	Main cause of death: xt012 coded

Wave 3 End of Life:

sl_xt011_	Main cause of death
sl_xt012c	Main cause of death: xt012 coded

Wave 4 End of Life:

xt011_	Main cause of death
xt012c	Main cause of death: xt012 coded

Wave 5 End of Life:

xt011_	Main cause of death
xt012c	Main cause of death: xt012 coded

Wave 6 End of Life:

Section A. Demographics

xt011_	Main cause of death
xt012c	Main cause of death: xt012 coded
Wave 7 End of Life:	
xt011_	Main cause of death
xt012c	Main cause of death: xt012 coded
Wave 8 End of Life:	
xt011_	Main cause of death
xt012c	Main cause of death: xt012 coded
Wave 9 End of Life:	
xt011_	Main cause of death
xt012c	Main cause of death: xt012 coded
Wave 9 COVID End of Life:	
xt011_	Main cause of death
xt012c	Main cause of death: xt012 coded

Duration of Final Illness			
Variable	Waves	Label	Type
raddur	All	raddur: r duration final illness/death	Categ

How Constructed

raddur indicates the duration of the respondent's final illness. The proxy is asked, "How long had he/she been ill before he/she died?" **raddur** is coded as follows: 1.no warning, 1-2 hrs; 4.less than 1 month; 5.less than 1 year; 6.more than 1 year. This question in the SHARE end of life interview does not have fine enough categories to assign values of 2.less than one day, and 3.less than one week, in **raddur**. In all waves, **raddur** takes a value of 4 if the proxy reports that the respondent was ill for less than one month. **raddur** takes a value of 5 if the proxy reports that the respondent was ill for one or more months but less than 6 months, or if they were ill for 6 or more months but less than a year. **raddur** takes a value of 6 if the proxy reports that the respondent was ill for one year or more. Additionally, starting in Wave 8, **raddur** takes a value of 1 if the proxy reports that the respondent was not ill before death (spontaneous). Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Cross Wave Differences in SHARE EOL

The proxy can report that the respondent was not ill before death (spontaneous) starting in Wave 8.

Differences with the Harmonized HRS End of Life

The HRS exit interview has more categories than the SHARE end of life interview for this question. The SHARE answer categories for this question are not fine enough to assign value 1 in most waves and values 2 through 3 in all waves for **raddur**, while other answer categories have been combined where appropriate to be comparable to the HRS answer categories for the creation of **raddur** in the Harmonized End of Life datasets.

Categorical Variable Frequencies

	raddur
1.no warning, 1-2 hrs	873
4.less than 1 month	5,095
5.less than 1 year	6,000
6.1 year or more	8,435
.d:dk	208
.m:missing	444
.r:refuse	43
Total	21,098

Original SHARE Variables Used

Wave 3 End of Life:	
sl_xt013_	How long been ill before decease
Wave 4 End of Life:	
xt013_	How long been ill before decease
Wave 5 End of Life:	
xt013_	How long been ill before decease
Wave 6 End of Life:	
xt013_	How long been ill before decease
Wave 7 End of Life:	

Section A. Demographics

xt013_	How long been ill before decease
Wave 8 End of Life:	
xt013_	How long been ill before decease
Wave 9 End of Life:	
xt013_	How long been ill before decease
Wave 9 COVID End of Life:	
xt013_	How long been ill before decease

Relationship Status at Death			
Variable	Waves	Label	Type
radmarr	All	radmarr: r married at death	Categ
radmarrp	All	radmarrp: r married/partnered at death	Categ

How Constructed

radmarr indicates whether the respondent was married at the time of their death. Starting in Wave 3, **radmarr** is assigned based on a direct question in the end of life interview asking if the deceased respondent was married at the time of their death. **radmarr** is coded as 0 if the respondent was not married at the time of their death. **radmarr** is coded as 1 if the respondent was married at the time of their death. **radmarr** is assigned special missing (.q) in Wave 2 because there was no question asking about the respondent's marital status at the time of death. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

radmarrp indicates whether the respondent was married or partnered at the time of their death. Starting in Wave 3, the relationship of the proxy for the end of life interview, which has an option for husband/wife/partner, is combined with whether the proxy reports that the respondent was married at the time of their death. **radmarrp** is coded as 0 if the respondent was not married at the time of their death and the proxy was not the respondent's husband/wife/partner. **radmarrp** is coded as 1 if the respondent was married at the time of their death or the proxy was the respondent's husband/wife/partner. **radmarrp** is assigned special missing (.q) in Wave 2 because there was no question asking about the respondent's marital status at the time of death. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Cross Wave Differences in SHARE EOL

In Wave 2, the proxy is not asked about the respondent's marital status at death. Starting in Wave 3, the proxy is given a direct question asking whether the respondent was married at death.

Differences with the Harmonized HRS End of Life

The SHARE end of life interview inquires whether the respondent was married at death in a single question starting in Wave 3, but does not ask directly about being partnered at death. Rather, the relationship of the proxy for the end of life interview is used to supplement the direct question and determine whether the respondent was married or partnered at the time of their death. The HRS exit interview inquires about the respondent's relationship status at death in several different questions which are used to determine whether the respondent was married or partnered at the time of their death.

Categorical Variable Frequencies

radmarr	
0.no	9,443
1.yes	10,815
.d:dk	28
.m:missing	77
.q:not asked this wave	726
.r:refuse	9
Total	21,098
radmarrp	
0.no	9,052
1.yes	11,215
.d:dk	29
.m:missing	69
.q:not asked this wave	726

.r:refuse	7
Total	21,098

Original SHARE Variables Used

Wave 3 End of Life:

sl_xt002_	Relationship to the deceased
sl_xt109_	Deceased married at time of death

Wave 4 End of Life:

xt002_	Relationship to the deceased
xt109_	Deceased married at time of death

Wave 5 End of Life:

xt002_	Relationship to the deceased
xt109_	Deceased married at time of death

Wave 6 End of Life:

xt002_	Relationship to the deceased
xt109_	Deceased married at time of death

Wave 7 End of Life:

xt002_	Relationship to the deceased
xt109_	Deceased married at time of death

Wave 8 End of Life:

xt002_	Relationship to the deceased
xt109_	Deceased married at time of death

Wave 9 End of Life:

xt002_	Relationship to the deceased
xt109_	Deceased married at time of death

Wave 9 COVID End of Life:

xt002_	Relationship to the deceased
xt109_	Deceased married at time of death

B. Health

Activities of Daily Living: Whether Had Any Difficulty with ADLs in Final Year			
Variable	Waves	Label	Type
raxdressa1y	All	raxdressa1y: r had difficulty dressing final 12 months	Categ
raxwalkra1y	All	raxwalkra1y: r had difficulty walking across a room final 12 months	Categ
raxbatha1y	All	raxbatha1y: r had difficulty bathing final 12 months	Categ
raxeata1y	All	raxeata1y: r had difficulty eating final 12 months	Categ
raxbeda1y	All	raxbeda1y: r had difficulty getting in and out of bed final 12 months	Categ
raxtoilta1y	All	raxtoilta1y: r had difficulty using the toilet final 12 months	Categ
raxurina1y	All	raxurina1y: r had difficulty with continence final 12 months	Categ

How Constructed

raxdressa1y, **raxwalkra1y**, **raxbatha1y**, **raxeata1y**, **raxbeda1y**, **raxtoilta1y**, and **raxurina1y** indicate whether the deceased respondent had any difficulty with a specific activity of daily living in the last twelve months of life because of a physical, mental, emotional or memory problem, as reported by the proxy. The proxy was only asked to name difficulties that lasted at least three months. **raxdressa1y** indicates whether the respondent had difficulty with dressing, including putting on shoes and socks in the last twelve months of life. **raxwalkra1y** indicates whether the respondent had difficulty walking across a room in the last twelve months of life. **raxbatha1y** indicates whether the respondent had difficulty with bathing or showering in the last twelve months of life. **raxeata1y** indicates whether the respondent had difficulty with eating, such as cutting up food, in the last twelve months of life. **raxbeda1y** indicates whether the respondent had difficulty getting in or out of bed in the last twelve months of life. **raxtoilta1y** indicates whether the respondent had difficulty using the toilet, including getting up and down, in the last twelve months of life. **raxurina1y** indicates whether the respondent had difficulty with continence over urination or defecation in the last twelve months of life. **raxdressa1y**, **raxwalkra1y**, **raxbatha1y**, **raxeata1y**, **raxbeda1y**, **raxtoilta1y**, and **raxurina1y** are coded as 0 if the respondent had no difficulty with the specific activity in the last twelve months of life. **raxdressa1y**, **raxwalkra1y**, **raxbatha1y**, **raxeata1y**, **raxbeda1y**, **raxtoilta1y**, and **raxurina1y** are coded as 1 if the respondent had some difficulty with the specific activity in the last twelve months of life. Questions on difficulty with continence over urination or defecation are not asked until Wave 6, as such **raxurina1y** is assigned special missing (.q) in Waves 2 through 5. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Cross Wave Differences in SHARE EOL

Please note that difficulty with six activities of daily living (dressing, walking across a room, bathing, eating, getting in and out of bed, and using the toilet) are asked in the same question as four instrumental activities of daily living (preparing hot meals, shopping for groceries, using the phone, and taking medication) in Waves 2 through 4, whereas difficulty with the same six ADLs are asked by themselves starting in Wave 5.

Starting in Wave 6, difficulty with continence over urination or defecation is asked about with the instrumental activities of daily living.

Differences with the Harmonized HRS End of Life

The SHARE end of life interview asks about the respondent's difficulty with ADLs that lasted at least three months during the last twelve months of life. The HRS exit interview does not ask whether the respondent had difficulty with ADLs, but rather asks only if someone helped the respondent with the ADLs in the last three months of life. As such, the ADL difficulty variables in the Gateway Harmonized SHARE End of Life capture a different concept than the help received with ADL variables in the Harmonized HRS End of Life, and so are given different variable names.

SHARE asks about difficulty with continence that lasted at least three months during the last twelve months of life starting in Wave 6, while the HRS exit interview inquires about difficulty with continence that lasted at least one month during the last year of life. Please note that this variable has not yet been incorporated into the Harmonized HRS End of Life.

Categorical Variable Frequencies

	raxdressa1y
0.no	10,342
1.yes	10,078
.d.dk	432
.m.missing	145
.r.refuse	101
Total	21,098
	raxwalkra1y
0.no	12,716
1.yes	7,704
.d.dk	432
.m.missing	145
.r.refuse	101
Total	21,098
	raxbatha1y
0.no	9,337
1.yes	11,083
.d.dk	432
.m.missing	145
.r.refuse	101
Total	21,098
	raxeata1y
0.no	13,950
1.yes	6,470
.d.dk	432
.m.missing	145
.r.refuse	101
Total	21,098
	raxbeda1y
0.no	12,247
1.yes	8,173
.d.dk	432
.m.missing	145
.r.refuse	101
Total	21,098
	raxtoilta1y
0.no	12,560
1.yes	7,860
.d.dk	432
.m.missing	145
.r.refuse	101

Total	21,098
	raxurina1y
o.no	10,050
1.yes	5,349
.d:dk	307
.m:missing	64
.q:not asked this wave	5,254
.r:refuse	74
Total	21,098

Original SHARE Variables Used

Wave 2 End of Life:

xto2od1	Difficulties doing activities: dressing, including putting on shoes and socks
xto2od2	Difficulties doing activities: walking across a room
xto2od3	Difficulties doing activities: bathing or showering
xto2od4	Difficulties doing activities: eating, such as cutting up your food
xto2od5	Difficulties doing activities: getting in or out of bed
xto2od6	Difficulties doing activities: using the toilet, including getting up or down
xto2odno	Difficulties doing activities: none of these

Wave 3 End of Life:

sl_xto2od1	Difficulties: dressing, including putting on shoes and socks
sl_xto2od2	Difficulties: walking across a room
sl_xto2od3	Difficulties: bathing or showering
sl_xto2od4	Difficulties: eating, such as cutting up your food
sl_xto2od5	Difficulties: getting in or out of bed
sl_xto2od6	Difficulties: using the toilet, including getting up or down
sl_xto2odno	Difficulties: none of these

Wave 4 End of Life:

xto2od1	Difficulties doing activities: dressing, including putting on shoes and socks
xto2od2	Difficulties doing activities: walking across a room
xto2od3	Difficulties doing activities: bathing or showering
xto2od4	Difficulties doing activities: eating, such as cutting up your food
xto2od5	Difficulties doing activities: getting in or out of bed
xto2od6	Difficulties doing activities: using the toilet, including getting up or down
xto2odno	Difficulties doing activities: none of these

Wave 5 End of Life:

xto2od1	Difficulties doing activities: dressing, including putting on shoes and socks
xto2od2	Difficulties doing activities: walking across a room
xto2od3	Difficulties doing activities: bathing or showering
xto2od4	Difficulties doing activities: eating, such as cutting up your food
xto2od5	Difficulties doing activities: getting in or out of bed
xto2od6	Difficulties doing activities: using the toilet, including getting up or down
xto2odno	Difficulties doing activities: none of these

Wave 6 End of Life:

xto2od1	Difficulties doing activities: dressing, including putting on shoes and socks
xto2od2	Difficulties doing activities: walking across a room
xto2od3	Difficulties doing activities: bathing or showering
xto2od4	Difficulties doing activities: eating, such as cutting up your food
xto2od5	Difficulties doing activities: getting in or out of bed
xto2od6	Difficulties doing activities: using the toilet, including getting up or down

xt02odno	Difficulties doing activities: none of these
xt62od1o	Difficulties doing activities II: continence over urination/defecation
xt62odno	Difficulties doing activities II: none of these
Wave 7 End of Life:	
xt02od1	Difficulties doing activities: dressing, including putting on shoes and socks
xt02od2	Difficulties doing activities: walking across a room
xt02od3	Difficulties doing activities: bathing or showering
xt02od4	Difficulties doing activities: eating, such as cutting up your food
xt02od5	Difficulties doing activities: getting in or out of bed
xt02od6	Difficulties doing activities: using the toilet, including getting up or down
xt02odno	Difficulties doing activities: none of these
xt62od1o	Difficulties doing activities II: continence over urination/defecation
xt62odno	Difficulties doing activities II: none of these
Wave 8 End of Life:	
xt02od1	Difficulties doing activities: dressing, including putting on shoes and socks
xt02od2	Difficulties doing activities: walking across a room
xt02od3	Difficulties doing activities: bathing or showering
xt02od4	Difficulties doing activities: eating, such as cutting up your food
xt02od5	Difficulties doing activities: getting in or out of bed
xt02od6	Difficulties doing activities: using the toilet, including getting up or down
xt02odno	Difficulties doing activities: none of these
xt62od1o	Difficulties doing activities II: continence over urination/defecation
xt62odno	Difficulties doing activities II: none of these
Wave 9 End of Life:	
xt02od1	Difficulties doing activities: dressing, including putting on shoes and socks
xt02od2	Difficulties doing activities: walking across a room
xt02od3	Difficulties doing activities: bathing or showering
xt02od4	Difficulties doing activities: eating, such as cutting up your food
xt02od5	Difficulties doing activities: getting in or out of bed
xt02od6	Difficulties doing activities: using the toilet, including getting up or down
xt02odno	Difficulties doing activities: none of these
xt62od1o	Difficulties doing activities II: continence over urination/defecation
xt62odno	Difficulties doing activities II: none of these
Wave 9 COVID End of Life:	
xt02od1	Difficulties doing activities: dressing, including putting on shoes and socks
xt02od2	Difficulties doing activities: walking across a room
xt02od3	Difficulties doing activities: bathing or showering
xt02od4	Difficulties doing activities: eating, such as cutting up your food
xt02od5	Difficulties doing activities: getting in or out of bed
xt02od6	Difficulties doing activities: using the toilet, including getting up or down
xt02odno	Difficulties doing activities: none of these
xt62od1o	Difficulties doing activities II: continence over urination/defecation
xt62odno	Difficulties doing activities II: none of these

Activities of Daily Living: Summary Measures

Variable	Waves	Label	Type
raxadlwa	All	raxadlwa: r some diff 3 item adl wallace summary	Cont
raxadlwam	All	raxadlwam: r missings in 3 item adl wallace summary	Cont
raxadlwaa	All	raxadlwaa: r any diff 3 item adl wallace summary	Categ
raxadla	All	raxadla: r some diff 5 item adl wallace summary	Cont
raxadlam	All	raxadlam: r missings in 5 item adl wallace summary	Cont
raxadlaa	All	raxadlaa: r any diff 5 item adl wallace summary	Categ
raxadlfive	All	raxadlfive: r some diff 5 item adl summary	Cont
raxadlfivem	All	raxadlfivem: r missings in 5 item adl summary	Cont
raxadlfivea	All	raxadlfivea: r any diff 5 item adl summary	Categ
raxadltot1_s	All	raxadltot1_s: r some diff 6 item total adl summary	Cont
raxadltot1m_s	All	raxadltot1m_s: r missings in 6 item total adl summary	Cont
raxadltot1a_s	All	raxadltot1a_s: r any diff 6 item total adl summary	Categ
raxadltot2_s	All	raxadltot2_s: r some diff 7 item total adl summary	Cont
raxadltot2m_s	All	raxadltot2m_s: r missings in 7 item total adl summary	Cont
raxadltot2a_s	All	raxadltot2a_s: r any diff 7 item total adl summary	Categ

How Constructed

raxadlwa is a 3-item summary of activities of daily living (ADLs), as proposed by Wallace and Herzog in the paper (Wallace and Herzog, 1995). The ADLs include bathing or showering (**raxbatha1y**), dressing (**raxdressa1y**), and eating (**raxeata1y**). Each limitation adds one to the summary measure, that is: **raxadlwa** = **raxbatha1y** + **raxdressa1y** + **raxeata1y**. **raxadlwam** is the number of ADL questions with missing values, ranging from 0 to 3. **raxadlwa** is calculated for all respondents who answered at least one of the ADL component questions, that is respondents with **raxadlwam** values of less than three. **raxadlwaa** indicates whether the respondent had any difficulty with any of the component variables. **raxadlwaa** is assigned a 0 if the respondent had no difficulty with any of the component variables, and is assigned a 1 if the respondent had difficulty with at least one of the component variables. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

raxadla is a 5-item summary of activities of daily living, as proposed by Wallace and Herzog in the paper (Wallace and Herzog, 1995). The ADLs include bathing or showering (**raxbatha1y**), dressing (**raxdressa1y**), eating (**raxeata1y**), getting in and out of bed (**raxbeda1y**), and walking across a room (**raxwalkra1y**). Each limitation adds one to the summary measure, that is: **raxadla** = **raxbatha1y** + **raxdressa1y** + **raxeata1y** + **raxbeda1y** + **raxwalkra1y**. **raxadlam** is the number of ADL questions with missing values, ranging from 0 to 5. **raxadla** is calculated for all respondents who answered at least one of the ADL component questions, that is respondents with **raxadlam** values of less than five. **raxadlaa** indicates whether the respondent had any difficulty with any of the component variables. **raxadlaa** is assigned a 0 if the respondent had no difficulty with any of the component variables, and is assigned a 1 if the respondent had difficulty with at least one of the component variables. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

raxadlfive is an alternative 5-item summary of activities of daily living. The ADLs include bathing or showering (**raxbatha1y**), dressing (**raxdressa1y**), eating (**raxeata1y**), getting in and out of bed (**raxbeda1y**), and using the toilet (**raxtoilta1y**). Each limitation adds one to the summary measure, that is: **raxadlfive** = **raxbatha1y** + **raxdressa1y** + **raxeata1y** + **raxbeda1y** + **raxtoilta1y**. **raxadlfivem** is the number of ADL questions with missing values, ranging from 0 to 5. **raxadlfive** is calculated for all respondents who answered at least one of the ADL component questions, that is respondents with **raxadlfivem** values of less than five. **raxadlfivea** indicates whether

the respondent had any difficulty with any of the component variables. **raxadlfivea** is assigned a 0 if the respondent had no difficulty with any of the component variables, and is assigned a 1 if the respondent had difficulty with at least one of the component variables. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

raxadltot1_s is a 6-item summary of the most commonly provided activities of daily living. The ADLs include walking across a room (**raxwalkra1y**), bathing or showering (**raxbatha1y**), dressing (**raxdressa1y**), eating (**raxeata1y**), getting in and out of bed (**raxbeda1y**), and using the toilet (**raxtoilta1y**). Each limitation adds one to the summary measure, that is: **raxadltot1_s** = **raxwalkra1y** + **raxbatha1y** + **raxdressa1y** + **raxeata1y** + **raxbeda1y** + **raxtoilta1y**. **raxadltot1m_s** is the number of ADL questions with missing values, ranging from 0 to 6. **raxadltot1_s** is calculated for all respondents who answered at least one of the ADL component questions, that is respondents with **raxadltot1m_s** values of less than six. **raxadltot1a_s** indicates whether the respondent had any difficulty with any of the component variables. **raxadltot1a_s** is assigned a 0 if the respondent had no difficulty with any of the component variables, and is assigned a 1 if the respondent had difficulty with at least one of the component variables. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

raxadltot2_s is a 7-item summary of all provided activities of daily living. The ADLs include walking across a room (**raxwalkra1y**), bathing or showering (**raxbatha1y**), dressing (**raxdressa1y**), eating (**raxeata1y**), getting in and out of bed (**raxbeda1y**), using the toilet (**raxtoilta1y**), and continence over urination or defecation (**raxurina1y**). Each limitation adds one to the summary measure, that is: **raxadltot2_s** = **raxwalkra1y** + **raxbatha1y** + **raxdressa1y** + **raxeata1y** + **raxbeda1y** + **raxtoilta1y** + **raxurina1y**. **raxadltot2m_s** is the number of ADL questions with missing values, ranging from 0 to 7. **raxadltot2_s** is calculated for all respondents who answered at least one of the ADL component questions, that is respondents with **raxadltot2m_s** values of less than seven. **raxadltot2a_s** indicates whether the respondent had any difficulty with any of the component variables. **raxadltot2a_s** is assigned a 0 if the respondent had no difficulty with any of the component variables, and is assigned a 1 if the respondent had difficulty with at least one of the component variables. Questions about difficulty with continence over urination or defecation are asked starting in Wave 6, as such **raxadltot2_s**, **raxadltot2m_s**, and **raxadltot2a_s** are assigned special missing (.q) in Waves 2 through 5. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Cross Wave Differences in SHARE EOL

Please note that difficulty with six activities of daily living (dressing, walking across a room, bathing, eating, getting in and out of bed, and using the toilet) are asked in the same question as four instrumental activities of daily living (preparing hot meals, shopping for groceries, using the phone, and taking medication) in Waves 2 through 4, whereas difficulty with the same six ADLs are asked by themselves starting in Wave 5.

Starting in Wave 6, difficulty with continence over urination or defecation is asked about with the instrumental activities of daily living.

Differences with the Harmonized HRS End of Life

The SHARE end of life interview asks about the respondent's difficulty with ADLs that lasted at least three months during the last twelve months of life. The HRS exit interview does not ask whether the respondent had difficulty with ADLs, but rather asks only if someone helped the respondent with the ADLs in the last three months of life. As such, the ADL difficulty variables in the Gateway Harmonized SHARE End of Life capture a different concept than the help received with ADL variables in the Harmonized HRS End of Life, and so are given different variable names.

SHARE asks about difficulty with continence that lasted at least three months during the last twelve months of life starting in Wave 6, while the HRS exit interview inquires about difficulty with continence that lasted at least one month during the last year of life. Please note that this variable has not yet been incorporated into the Harmonized HRS End of Life.

Continuous Variable Distributions

Variable	N	Mean	Std.Dev	Minimum	Maximum	Total missing
----------	---	------	---------	---------	---------	---------------

Section B. Health

raxadlwa	20,420	1.35	1.28	0.00	3.00	678
raxadlwam	21,098	0.10	0.53	0.00	3.00	0
raxadla	20,420	2.13	2.08	0.00	5.00	678
raxadlam	21,098	0.16	0.88	0.00	5.00	0
raxadlfive	20,420	2.14	2.10	0.00	5.00	678
raxadlfivem	21,098	0.16	0.88	0.00	5.00	0
raxadltot1_s	20,420	2.52	2.50	0.00	6.00	678
raxadltot1m_s	21,098	0.19	1.06	0.00	6.00	0
raxadltot2_s	15,504	2.86	2.84	0.00	7.00	5,594
raxadltot2m_s	15,844	0.23	1.20	0.00	7.00	5,254

Categorical Variable Frequencies

	raxadlwaa
0.no	8,443
1.yes	11,977
.d.dk	432
.m.missing	145
.r.refuse	101
Total	21,098

	raxadlaa
0.no	8,098
1.yes	12,322
.d.dk	432
.m.missing	145
.r.refuse	101
Total	21,098

	raxadlfivea
0.no	8,174
1.yes	12,246
.d.dk	432
.m.missing	145
.r.refuse	101
Total	21,098

	raxadltot1a_s
0.no	8,021
1.yes	12,399
.d.dk	432
.m.missing	145
.r.refuse	101
Total	21,098

	raxadltot2a_s
0.no	5,860

1.yes	9,644
.d:dk	225
.m:missing	64
.q:not asked this wave	5,254
.r:refuse	51
Total	21,098

Original SHARE Variables Used

Wave 2 End of Life:

xto2od1	Difficulties doing activities: dressing, including putting on shoes and socks
xto2od2	Difficulties doing activities: walking across a room
xto2od3	Difficulties doing activities: bathing or showering
xto2od4	Difficulties doing activities: eating, such as cutting up your food
xto2od5	Difficulties doing activities: getting in or out of bed
xto2od6	Difficulties doing activities: using the toilet, including getting up or down
xto2od7	Difficulties doing activities: preparing a hot meal
xto2od8	Difficulties doing activities: shopping for groceries
xto2od9	Difficulties doing activities: making telephone calls
xto2od10	Difficulties doing activities: taking medication
xto2odno	Difficulties doing activities: none of these

Wave 3 End of Life:

sl_xto2od1	Difficulties: dressing, including putting on shoes and socks
sl_xto2od2	Difficulties: walking across a room
sl_xto2od3	Difficulties: bathing or showering
sl_xto2od4	Difficulties: eating, such as cutting up your food
sl_xto2od5	Difficulties: getting in or out of bed
sl_xto2od6	Difficulties: using the toilet, including getting up or down
sl_xto2od7	Difficulties: preparing a hot meal
sl_xto2od8	Difficulties: shopping for groceries
sl_xto2od9	Difficulties: making telephone calls
sl_xto2od10	Difficulties: taking medication
sl_xto2odno	Difficulties: none of these

Wave 4 End of Life:

xto2od1	Difficulties doing activities: dressing, including putting on shoes and socks
xto2od2	Difficulties doing activities: walking across a room
xto2od3	Difficulties doing activities: bathing or showering
xto2od4	Difficulties doing activities: eating, such as cutting up your food
xto2od5	Difficulties doing activities: getting in or out of bed
xto2od6	Difficulties doing activities: using the toilet, including getting up or down
xto2od7	Difficulties doing activities: preparing a hot meal
xto2od8	Difficulties doing activities: shopping for groceries
xto2od9	Difficulties doing activities: making telephone calls
xto2od10	Difficulties doing activities: taking medication
xto2odno	Difficulties doing activities: none of these

Wave 5 End of Life:

xto2od1	Difficulties doing activities: dressing, including putting on shoes and socks
xto2od2	Difficulties doing activities: walking across a room
xto2od3	Difficulties doing activities: bathing or showering
xto2od4	Difficulties doing activities: eating, such as cutting up your food
xto2od5	Difficulties doing activities: getting in or out of bed
xto2od6	Difficulties doing activities: using the toilet, including getting up or down

xt02odno	Difficulties doing activities: none of these
Wave 6 End of Life:	
xt02od1	Difficulties doing activities: dressing, including putting on shoes and socks
xt02od2	Difficulties doing activities: walking across a room
xt02od3	Difficulties doing activities: bathing or showering
xt02od4	Difficulties doing activities: eating, such as cutting up your food
xt02od5	Difficulties doing activities: getting in or out of bed
xt02od6	Difficulties doing activities: using the toilet, including getting up or down
xt02odno	Difficulties doing activities: none of these
xt62od1o	Difficulties doing activities II: continence over urination/defecation
xt62odno	Difficulties doing activities II: none of these
Wave 7 End of Life:	
xt02od1	Difficulties doing activities: dressing, including putting on shoes and socks
xt02od2	Difficulties doing activities: walking across a room
xt02od3	Difficulties doing activities: bathing or showering
xt02od4	Difficulties doing activities: eating, such as cutting up your food
xt02od5	Difficulties doing activities: getting in or out of bed
xt02od6	Difficulties doing activities: using the toilet, including getting up or down
xt02odno	Difficulties doing activities: none of these
xt62od1o	Difficulties doing activities II: continence over urination/defecation
xt62odno	Difficulties doing activities II: none of these
Wave 8 End of Life:	
xt02od1	Difficulties doing activities: dressing, including putting on shoes and socks
xt02od2	Difficulties doing activities: walking across a room
xt02od3	Difficulties doing activities: bathing or showering
xt02od4	Difficulties doing activities: eating, such as cutting up your food
xt02od5	Difficulties doing activities: getting in or out of bed
xt02od6	Difficulties doing activities: using the toilet, including getting up or down
xt02odno	Difficulties doing activities: none of these
xt62od1o	Difficulties doing activities II: continence over urination/defecation
xt62odno	Difficulties doing activities II: none of these
Wave 9 End of Life:	
xt02od1	Difficulties doing activities: dressing, including putting on shoes and socks
xt02od2	Difficulties doing activities: walking across a room
xt02od3	Difficulties doing activities: bathing or showering
xt02od4	Difficulties doing activities: eating, such as cutting up your food
xt02od5	Difficulties doing activities: getting in or out of bed
xt02od6	Difficulties doing activities: using the toilet, including getting up or down
xt02odno	Difficulties doing activities: none of these
xt62od1o	Difficulties doing activities II: continence over urination/defecation
xt62odno	Difficulties doing activities II: none of these
Wave 9 COVID End of Life:	
xt02od1	Difficulties doing activities: dressing, including putting on shoes and socks
xt02od2	Difficulties doing activities: walking across a room
xt02od3	Difficulties doing activities: bathing or showering
xt02od4	Difficulties doing activities: eating, such as cutting up your food
xt02od5	Difficulties doing activities: getting in or out of bed
xt02od6	Difficulties doing activities: using the toilet, including getting up or down
xt02odno	Difficulties doing activities: none of these
xt62od1o	Difficulties doing activities II: continence over urination/defecation
xt62odno	Difficulties doing activities II: none of these

Instrumental Activities of Daily Living: Whether Had Any Difficulty with IADLs in Final Year

Variable	Waves	Label	Type
raxmealsa1y	All	raxmealsa1y: r had difficulty with hot meals final 12 months	Categ
raxshopa1y	All	raxshopa1y: r had difficulty with grocery shopping final 12 months	Categ
raxphonea1y	All	raxphonea1y: r had difficulty with using the phone final 12 months	Categ
raxmedsa1y	All	raxmedsa1y: r had difficulty with medications final 12 months	Categ
raxmapa1y	All	raxmapa1y: r had difficulty using a map final 12 months	Categ
raxhousewka1y	All	raxhousewka1y: r had difficulty doing house/garden work final 12 months	Categ
raxmoneya1y	All	raxmoneya1y: r had difficulty managing money final 12 months	Categ
raxleavhsa1y	All	raxleavhsa1y: r had difficulty leaving house alone final 12 months	Categ
raxlaundrya1y	All	raxlaundrya1y: r had difficulty doing personal laundry final 12 months	Categ

How Constructed

raxmealsa1y, **raxshopa1y**, **raxphonea1y**, **raxmedsa1y**, **raxmapa1y**, **raxhousewka1y**, **raxmoneya1y**, **raxleavhsa1y**, and **raxlaundrya1y** indicate whether the deceased respondent had any difficulty with a specific instrumental activity of daily living in the last twelve months of life because of a physical, mental, emotional or memory problem, as reported by the proxy. The proxy was only asked to name difficulties that lasted at least three months. **raxmealsa1y** indicates whether the respondent had difficulty preparing hot meals in the last twelve months of life. **raxshopa1y** indicates whether the respondent had difficulty shopping for groceries in the last twelve months of life. **raxphonea1y** indicates whether the respondent had difficulty making phone calls in the last twelve months of life. **raxmedsa1y** indicates whether the respondent had difficulty taking medication in the last twelve months of life. **raxmapa1y** indicates whether the respondent had difficulty using a map to figure out how to get around in a strange place in the last twelve months of life. **raxhousewka1y** indicates whether the respondent had difficulty doing work around the house or garden in the last twelve months of life. **raxmoneya1y** indicates whether the respondent had difficulty managing money, such as paying bills and keeping track of expenses, in the last twelve months of life. **raxleavhsa1y** indicates whether the respondent had difficulty leaving the house independently and accessing transportation services in the last twelve months of life. **raxlaundrya1y** indicates whether the respondent had difficulty doing personal laundry in the last twelve months of life. **raxmealsa1y**, **raxshopa1y**, **raxphonea1y**, **raxmedsa1y**, **raxmapa1y**, **raxhousewka1y**, **raxmoneya1y**, **raxleavhsa1y**, and **raxlaundrya1y** are coded as 0 if the respondent had no difficulty with the specific activity in the last twelve months of life. **raxmealsa1y**, **raxshopa1y**, **raxphonea1y**, **raxmedsa1y**, **raxmapa1y**, **raxhousewka1y**, **raxmoneya1y**, **raxleavhsa1y**, and **raxlaundrya1y** are coded as 1 if the respondent had difficulty with the specific activity in the last twelve months of life. **raxmealsa1y**, **raxshopa1y**, **raxphonea1y**, and **raxmedsa1y** are asked in Waves 2 through 4, then again starting in Wave 6. As such in Wave 5, these variables are assigned special missing (.q). **raxmapa1y**, **raxhousewka1y**, **raxmoneya1y**, **raxleavhsa1y**, and **raxlaundrya1y** are asked starting in Wave 6. Therefore in Waves 2 through 5, these variables are assigned special missing (.q). Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Cross Wave Differences in SHARE EOL

Difficulty with preparing hot meals, grocery shopping, using the phone, and taking medications are asked in Waves 2 through 4, then again starting in Wave 6. In Waves 2 through 4, difficulty with these 4 activities is asked in the same question as difficulty with 6 activities of daily living (dressing, walking across a room, bathing, eating, getting in and out of bed, and using the toilet). In Wave 5, difficulty with these activities is not asked about at all. Starting in Wave 6, difficulty with these

activities is asked with 5 new instrumental activities of daily living and difficulty with continence over urination or defecation, as a separate question from difficulties with ADLs.

Difficulty with using a map, doing work around the house or garden, managing money, leaving the house independently, and doing personal laundry are asked starting in Wave 6.

Differences with the Harmonized HRS End of Life

The SHARE end of life interview asks about the respondent's difficulty with IADLs that lasted at least three months during the last twelve months of life. The HRS exit interview does not ask whether the respondent had difficulty with IADLs, but rather asks only if someone helped the respondent with the IADLs in the last three months of life. As such, the IADL difficulty variables in the Gateway Harmonized SHARE End of Life capture a different concept than the help received with IADL variables in the Harmonized HRS End of Life, and so are given different variable names.

The HRS exit interview asks only about help received with preparing hot meals, shopping for groceries, making telephone calls, and taking medication, the same activities in Waves 2 through 4 of the SHARE end of life interview. SHARE asks about difficulty with using a map, doing work around the house or garden, managing money, leaving the house independently, and doing personal laundry starting in Wave 6, which are not asked about in the HRS exit interview.

Categorical Variable Frequencies

	raxmealsa1y
0.no	8,719
1.yes	9,664
.d:dk	357
.m:missing	88
.q:not asked this wave	2,187
.r:refuse	83
Total	21,098
	raxshopa1y
0.no	7,694
1.yes	10,689
.d:dk	357
.m:missing	88
.q:not asked this wave	2,187
.r:refuse	83
Total	21,098
	raxphonea1y
0.no	12,440
1.yes	5,943
.d:dk	357
.m:missing	88
.q:not asked this wave	2,187
.r:refuse	83
Total	21,098
	raxmedsa1y
0.no	11,657
1.yes	6,726
.d:dk	357
.m:missing	88

Section B. Health

.q:not asked this wave	2,187
.r:refuse	83
Total	21,098
raxmapa1y	
o.no	7,946
1.yes	7,453
.d:dk	307
.m:missing	64
.q:not asked this wave	5,254
.r:refuse	74
Total	21,098
raxhousewka1y	
o.no	6,204
1.yes	9,195
.d:dk	307
.m:missing	64
.q:not asked this wave	5,254
.r:refuse	74
Total	21,098
raxmoneya1y	
o.no	8,140
1.yes	7,259
.d:dk	307
.m:missing	64
.q:not asked this wave	5,254
.r:refuse	74
Total	21,098
raxleavhsa1y	
o.no	6,502
1.yes	8,897
.d:dk	307
.m:missing	64
.q:not asked this wave	5,254
.r:refuse	74
Total	21,098
raxlaundrya1y	
o.no	7,205
1.yes	8,194
.d:dk	307
.m:missing	64
.q:not asked this wave	5,254
.r:refuse	74
Total	21,098

Original SHARE Variables Used

Wave 2 End of Life:

xt02od7	Difficulties doing activities: preparing a hot meal
xt02od8	Difficulties doing activities: shopping for groceries
xt02od9	Difficulties doing activities: making telephone calls
xt02od10	Difficulties doing activities: taking medication
xt02odno	Difficulties doing activities: none of these

Wave 3 End of Life:

sl_xt02od7	Difficulties: preparing a hot meal
sl_xt02od8	Difficulties: shopping for groceries
sl_xt02od9	Difficulties: making telephone calls
sl_xt02od10	Difficulties: taking medication
sl_xt02odno	Difficulties: none of these

Wave 4 End of Life:

xt02od7	Difficulties doing activities: preparing a hot meal
xt02od8	Difficulties doing activities: shopping for groceries
xt02od9	Difficulties doing activities: making telephone calls
xt02od10	Difficulties doing activities: taking medication
xt02odno	Difficulties doing activities: none of these

Wave 6 End of Life:

xt62od1	Difficulties doing activities II: preparing a hot meal
xt62od2	Difficulties doing activities II: shopping for groceries
xt62od3	Difficulties doing activities II: making telephone calls
xt62od4	Difficulties doing activities II: taking medication
xt62od5	Difficulties doing activities II: using a map
xt62od6	Difficulties doing activities II: doing work around the house/garden
xt62od7	Difficulties doing activities II: managing money
xt62od8	Difficulties doing activities II: leaving the house independently
xt62od9	Difficulties doing activities II: doing personal laundry
xt62odno	Difficulties doing activities II: none of these

Wave 7 End of Life:

xt62od1	Difficulties doing activities II: preparing a hot meal
xt62od2	Difficulties doing activities II: shopping for groceries
xt62od3	Difficulties doing activities II: making telephone calls
xt62od4	Difficulties doing activities II: taking medication
xt62od5	Difficulties doing activities II: using a map
xt62od6	Difficulties doing activities II: doing work around the house/garden
xt62od7	Difficulties doing activities II: managing money
xt62od8	Difficulties doing activities II: leaving the house independently
xt62od9	Difficulties doing activities II: doing personal laundry
xt62odno	Difficulties doing activities II: none of these

Wave 8 End of Life:

xt62od1	Difficulties doing activities II: preparing a hot meal
xt62od2	Difficulties doing activities II: shopping for groceries
xt62od3	Difficulties doing activities II: making telephone calls
xt62od4	Difficulties doing activities II: taking medication
xt62od5	Difficulties doing activities II: using a map
xt62od6	Difficulties doing activities II: doing work around the house/garden
xt62od7	Difficulties doing activities II: managing money
xt62od8	Difficulties doing activities II: leaving the house independently
xt62od9	Difficulties doing activities II: doing personal laundry
xt62odno	Difficulties doing activities II: none of these

Wave 9 End of Life:

Section B. Health

xt62od1	Difficulties doing activities II: preparing a hot meal
xt62od2	Difficulties doing activities II: shopping for groceries
xt62od3	Difficulties doing activities II: making telephone calls
xt62od4	Difficulties doing activities II: taking medication
xt62od5	Difficulties doing activities II: using a map
xt62od6	Difficulties doing activities II: doing work around the house/garden
xt62od7	Difficulties doing activities II: managing money
xt62od8	Difficulties doing activities II: leaving the house independently
xt62od9	Difficulties doing activities II: doing personal laundry
xt62odno	Difficulties doing activities II: none of these

Wave 9 COVID End of Life:

xt62od1	Difficulties doing activities II: preparing a hot meal
xt62od2	Difficulties doing activities II: shopping for groceries
xt62od3	Difficulties doing activities II: making telephone calls
xt62od4	Difficulties doing activities II: taking medication
xt62od5	Difficulties doing activities II: using a map
xt62od6	Difficulties doing activities II: doing work around the house/garden
xt62od7	Difficulties doing activities II: managing money
xt62od8	Difficulties doing activities II: leaving the house independently
xt62od9	Difficulties doing activities II: doing personal laundry
xt62odno	Difficulties doing activities II: none of these

Instrumental Activities of Daily Living: Summary Measures			
Variable	Waves	Label	Type
raxiadla	All	raxiadla: r some diff 3 item iadl summary	Cont
raxiadlam	All	raxiadlam: r missings in 3 item iadl summary	Cont
raxiadlaa	All	raxiadlaa: r any diff 3 item iadl summary	Categ
raxiadlfour	All	raxiadlfour: r some diff 4 item iadl summary	Cont
raxiadlfourm	All	raxiadlfourm: r missings in 4 item iadl summary	Cont
raxiadlfoura	All	raxiadlfoura: r any diff 4 item iadl summary	Categ
raxiadlza	All	raxiadlza: r some diff 5 item iadl summary	Cont
raxiadlzam	All	raxiadlzam: r missings in 5 item iadl summary	Cont
raxiadlzaa	All	raxiadlzaa: r any diff 5 item iadl summary	Categ
raxiadltot1_s	All	raxiadltot1_s: r some diff 4 item total iadl summary	Cont
raxiadltot1m_s	All	raxiadltot1m_s: r missings in 4 item total iadl summary	Cont
raxiadltot1a_s	All	raxiadltot1a_s: r any diff 4 item total iadl summary	Categ
raxiadltot2_s	All	raxiadltot2_s: r some diff 9 item total iadl summary	Cont
raxiadltot2m_s	All	raxiadltot2m_s: r missings in 9 item total iadl summary	Cont
raxiadltot2a_s	All	raxiadltot2a_s: r any diff 9 item total iadl summary	Categ

How Constructed

raxiadla is a 3-item summary of instrumental activities of daily living (IADLs). The IADLs include using the phone (**raxphonea1y**), managing money (**raxmoneya1y**), and taking medications (**raxmedsa1y**). Each limitation adds one to the summary measure, that is: **raxiadla** = **raxphonea1y** + **raxmoneya1y** + **raxmedsa1y**. **raxiadlam** is the number of IADL questions with missing values, ranging from 0 to 3. **raxiadla** is calculated for all respondents who answered at least one of the IADL component questions, that is respondents with **raxiadlam** values of less than three. **raxiadlaa** indicates whether the respondent had any difficulty with any of the component variables. **raxiadlaa** is assigned a 0 if the respondent had no difficulty with any of the component variables, and is assigned a 1 if the respondent had difficulty with at least one of the component variables. Difficulty with managing money is not asked until Wave 6. As such, in Waves 2 through 5, **raxiadla**, **raxiadlam**, and **raxiadlaa** are assigned special missing (.q). Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

raxiadlfour is a 4-item summary of instrumental activities of daily living. The IADLs include managing money (**raxmoneya1y**), taking medications (**raxmedsa1y**), grocery shopping (**raxshopa1y**), and preparing hot meals (**raxmealsa1y**). Each limitation adds one to the summary measure, that is: **raxiadlfour** = **raxmoneya1y** + **raxmedsa1y** + **raxshopa1y** + **raxmealsa1y**. **raxiadlfourm** is the number of IADL questions with missing values, ranging from 0 to 4. **raxiadlfour** is calculated for all respondents who answered at least one of the IADL component questions, that is respondents with **raxiadlfourm** values of less than four. **raxiadlfoura** indicates whether the respondent had any difficulty with any of the component variables. **raxiadlfoura** is assigned a 0 if the respondent had no difficulty with any of the component variables, and is assigned a 1 if the respondent had difficulty with at least one of the component variables. Difficulty with managing money is not asked until Wave 6. As such, in Waves 2 through 5, **raxiadlfour**, **raxiadlfourm**, and **raxiadlfoura** are assigned special missing (.q). Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

raxiadlza is a 5-item summary of instrumental activities of daily living. The IADLs include using the phone (**raxphonea1y**), managing money (**raxmoneya1y**), taking medications (**raxmedsa1y**), grocery shopping (**raxshopa1y**), and preparing hot meals (**raxmealsa1y**). Each limitation adds one to the summary measure, that is: **raxiadlza** = **raxphonea1y** + **raxmoneya1y** + **raxmedsa1y** + **raxshopa1y** + **raxmealsa1y**. **raxiadlzam** is the number of IADL questions with

missing values, ranging from 0 to 5. **raxiadlza** is calculated for all respondents who answered at least one of the IADL component questions, that is respondents with **raxiadlzam** values of less than five. **raxiadlzaa** indicates whether the respondent had any difficulty with any of the component variables. **raxiadlzaa** is assigned a 0 if the respondent had no difficulty with any of the component variables, and is assigned a 1 if the respondent had difficulty with at least one of the component variables. Difficulty with managing money is not asked until Wave 6. As such, in Waves 2 through 5, **raxiadlza**, **raxiadlzam**, and **raxiadlzaa** are assigned special missing (.q). Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

raxiadltot1_s is a 4-item summary of the most commonly provided instrumental activities of daily living. The IADLs include using the phone (**raxphonea1y**), taking medications (**raxmedsa1y**), grocery shopping (**raxshopa1y**), and preparing hot meals (**raxmealsa1y**). Each limitation adds one to the summary measure, that is: **raxiadltot1_s** = **raxphonea1y** + **raxmedsa1y** + **raxshopa1y** + **raxmealsa1y**. **raxiadltot1m_s** is the number of IADL questions with missing values, ranging from 0 to 4. **raxiadltot1_s** is calculated for all respondents who answered at least one of the IADL component questions, that is respondents with **raxiadltot1m_s** values of less than four. **raxiadltot1a_s** indicates whether the respondent had any difficulty with any of the component variables. **raxiadltot1a_s** is assigned a 0 if the respondent had no difficulty with any of the component variables, and is assigned a 1 if the respondent had difficulty with at least one of the component variables. Difficulty with IADL questions are not asked about in Wave 5, so in this wave **raxiadltot1_s**, **raxiadltot1m_s**, and **raxiadltot1a_s** are assigned special missing (.q). Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

raxiadltot2_s is a 9-item summary of all provided instrumental activities of daily living. The IADLs include using the phone (**raxphonea1y**), taking medications (**raxmedsa1y**), grocery shopping (**raxshopa1y**), preparing hot meals (**raxmealsa1y**), using a map (**raxmapa1y**), doing work around the house or garden (**raxhousewka1y**), managing money (**raxmoneya1y**), leaving the house independently (**raxleavhsa1y**), and doing personal laundry (**raxlaundrya1y**). Each limitation adds one to the summary measure, that is: **raxiadltot2_s** = **raxphonea1y** + **raxmedsa1y** + **raxshopa1y** + **raxmealsa1y** + **raxmapa1y** + **raxhousewka1y** + **raxmoneya1y** + **raxleavhsa1y** + **raxlaundrya1y**. **raxiadltot2m_s** is the number of IADL questions with missing values, ranging from 0 to 9. **raxiadltot2_s** is calculated for all respondents who answered at least one of the IADL component questions, that is respondents with **raxiadltot2m_s** values of less than nine. **raxiadltot2a_s** indicates whether the respondent had any difficulty with any of the component variables. **raxiadltot2a_s** is assigned a 0 if the respondent had no difficulty with any of the component variables, and is assigned a 1 if the respondent had difficulty with at least one of the component variables. Questions about difficulty using a map, doing work around the house or garden, managing money, leaving the house independently, and doing personal laundry are not asked until Wave 6. As such, in Waves 2 through 5, **raxiadltot2_s**, **raxiadltot2m_s**, and **raxiadltot2a_s** are assigned special missing (.q). Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Cross Wave Differences in SHARE EOL

Difficulty with preparing hot meals, grocery shopping, using the phone, and taking medications are asked in Waves 2 through 4, then again starting in Wave 6. In Waves 2 through 4, difficulty with these 4 activities is asked in the same question as difficulty with 6 activities of daily living (dressing, walking across a room, bathing, eating, getting in and out of bed, and using the toilet). In Wave 5, difficulty with these activities is not asked about at all. Starting in Wave 6, difficulty with these activities is asked with 5 new instrumental activities of daily living and difficulty with continence over urination or defecation, as a separate question from difficulties with ADLs.

Difficulty with using a map, doing work around the house or garden, managing money, leaving the house independently, and doing personal laundry are asked starting in Wave 6.

Differences with the Harmonized HRS End of Life

The SHARE end of life interview asks about the respondent's difficulty with IADLs that lasted at least three months during the last twelve months of life. The HRS exit interview does not ask whether the respondent had difficulty with IADLs, but rather asks only if someone helped the respondent with the IADLs in the last three months of life. As such, the IADL difficulty

variables in the Gateway Harmonized SHARE End of Life capture a different concept than the help received with IADL variables in the Harmonized HRS End of Life, and so are given different variable names.

The HRS exit interview asks only about help received with preparing hot meals, shopping for groceries, making telephone calls, and taking medication, the same activities in Waves 2 through 4 of the SHARE end of life interview. SHARE asks about difficulty with using a map, doing work around the house or garden, managing money, leaving the house independently, and doing personal laundry starting in Wave 6, which are not asked about in the HRS exit interview.

Continuous Variable Distributions

Variable	N	Mean	Std.Dev	Minimum	Maximum	Total missing
raxiadla	15,399	1.18	1.29	0.00	3.00	5,699
raxiadlam	15,844	0.08	0.50	0.00	3.00	5,254
raxiadlfour	15,399	1.96	1.72	0.00	4.00	5,699
raxiadlfourm	15,844	0.11	0.66	0.00	4.00	5,254
raxiadlza	15,399	2.29	2.08	0.00	5.00	5,699
raxiadlzam	15,844	0.14	0.83	0.00	5.00	5,254
raxiadltot1_s	18,383	1.80	1.66	0.00	4.00	2,715
raxiadltot1m_s	18,911	0.11	0.66	0.00	4.00	2,187
raxiadltot2_s	15,399	4.48	3.72	0.00	9.00	5,699
raxiadltot2m_s	15,844	0.25	1.49	0.00	9.00	5,254

Categorical Variable Frequencies

	raxiadlaa
0.no	7,411
1.yes	7,988
.d:dk	307
.m:missing	64
.q:not asked this wave	5,254
.r:refuse	74
Total	21,098
	raxiadlfoura
0.no	5,554
1.yes	9,845
.d:dk	307
.m:missing	64
.q:not asked this wave	5,254
.r:refuse	74
Total	21,098
	raxiadlzaa
0.no	5,529
1.yes	9,870
.d:dk	307

.m:missing	64
.q:not asked this wave	5,254
.r:refuse	74
Total	21,098
raxiadltot1a_s	
o.no	6,949
1.yes	11,434
.d:dk	357
.m:missing	88
.q:not asked this wave	2,187
.r:refuse	83
Total	21,098
raxiadltot2a_s	
o.no	4,709
1.yes	10,690
.d:dk	307
.m:missing	64
.q:not asked this wave	5,254
.r:refuse	74
Total	21,098

Original SHARE Variables Used

Wave 2 End of Life:

xt02od7	Difficulties doing activities: preparing a hot meal
xt02od8	Difficulties doing activities: shopping for groceries
xt02od9	Difficulties doing activities: making telephone calls
xt02od10	Difficulties doing activities: taking medication
xt02odno	Difficulties doing activities: none of these

Wave 3 End of Life:

sl_xt02od7	Difficulties: preparing a hot meal
sl_xt02od8	Difficulties: shopping for groceries
sl_xt02od9	Difficulties: making telephone calls
sl_xt02od10	Difficulties: taking medication
sl_xt02odno	Difficulties: none of these

Wave 4 End of Life:

xt02od7	Difficulties doing activities: preparing a hot meal
xt02od8	Difficulties doing activities: shopping for groceries
xt02od9	Difficulties doing activities: making telephone calls
xt02od10	Difficulties doing activities: taking medication
xt02odno	Difficulties doing activities: none of these

Wave 6 End of Life:

xt62od1	Difficulties doing activities II: preparing a hot meal
xt62od2	Difficulties doing activities II: shopping for groceries
xt62od3	Difficulties doing activities II: making telephone calls
xt62od4	Difficulties doing activities II: taking medication
xt62od5	Difficulties doing activities II: using a map
xt62od6	Difficulties doing activities II: doing work around the house/garden
xt62od7	Difficulties doing activities II: managing money
xt62od8	Difficulties doing activities II: leaving the house independently

xt62od9	Difficulties doing activities II: doing personal laundry
xt62odno	Difficulties doing activities II: none of these
Wave 7 End of Life:	
xt62od1	Difficulties doing activities II: preparing a hot meal
xt62od2	Difficulties doing activities II: shopping for groceries
xt62od3	Difficulties doing activities II: making telephone calls
xt62od4	Difficulties doing activities II: taking medication
xt62od5	Difficulties doing activities II: using a map
xt62od6	Difficulties doing activities II: doing work around the house/garden
xt62od7	Difficulties doing activities II: managing money
xt62od8	Difficulties doing activities II: leaving the house independently
xt62od9	Difficulties doing activities II: doing personal laundry
xt62odno	Difficulties doing activities II: none of these
Wave 8 End of Life:	
xt62od1	Difficulties doing activities II: preparing a hot meal
xt62od2	Difficulties doing activities II: shopping for groceries
xt62od3	Difficulties doing activities II: making telephone calls
xt62od4	Difficulties doing activities II: taking medication
xt62od5	Difficulties doing activities II: using a map
xt62od6	Difficulties doing activities II: doing work around the house/garden
xt62od7	Difficulties doing activities II: managing money
xt62od8	Difficulties doing activities II: leaving the house independently
xt62od9	Difficulties doing activities II: doing personal laundry
xt62odno	Difficulties doing activities II: none of these
Wave 9 End of Life:	
xt62od1	Difficulties doing activities II: preparing a hot meal
xt62od2	Difficulties doing activities II: shopping for groceries
xt62od3	Difficulties doing activities II: making telephone calls
xt62od4	Difficulties doing activities II: taking medication
xt62od5	Difficulties doing activities II: using a map
xt62od6	Difficulties doing activities II: doing work around the house/garden
xt62od7	Difficulties doing activities II: managing money
xt62od8	Difficulties doing activities II: leaving the house independently
xt62od9	Difficulties doing activities II: doing personal laundry
xt62odno	Difficulties doing activities II: none of these
Wave 9 COVID End of Life:	
xt62od1	Difficulties doing activities II: preparing a hot meal
xt62od2	Difficulties doing activities II: shopping for groceries
xt62od3	Difficulties doing activities II: making telephone calls
xt62od4	Difficulties doing activities II: taking medication
xt62od5	Difficulties doing activities II: using a map
xt62od6	Difficulties doing activities II: doing work around the house/garden
xt62od7	Difficulties doing activities II: managing money
xt62od8	Difficulties doing activities II: leaving the house independently
xt62od9	Difficulties doing activities II: doing personal laundry
xt62odno	Difficulties doing activities II: none of these

Doctor Diagnosed Conditions: Cancer

Variable	Waves	Label	Type
ralcancre	All	ralcancre: r ever had cancer as of last ivw	Categ

How Constructed

ralcancre indicates whether the deceased respondent had ever been told by a doctor that they had cancer or a malignant tumor, including leukemia or lymphoma, but excluding minor skin cancers, in their last completed core interview. **ralcancre** is based on **rWcancre** in the Gateway Harmonized SHARE and is taken from the last core wave completed. **ralcancre** is coded as 0 if the respondent reported never having been diagnosed with cancer, and is coded as 1 if the respondent reported being diagnosed with cancer. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Please note that in the SHARE core interview the respondent can dispute a previously positive report of cancer. Please see **rWcancre** in "Doctor Diagnosed Health Problems: Ever Have Condition" in the Gateway Harmonized SHARE codebook for more information on the flag variable or to determine whether the report of cancer has been disputed in a prior wave.

Cross Wave Differences in SHARE EOL

No differences known.

Differences with the Harmonized HRS End of Life

Unlike the HRS exit interview, the SHARE end of life interview does not ask proxies about the deceased respondent's cancer status by the time of their death. As such, we cannot provide variables indicating whether the respondent had been newly or ever diagnosed with cancer by the time of their death in the Gateway Harmonized SHARE End of Life as we do in the Harmonized HRS End of Life, we can only provide whether the respondent had ever been diagnosed with cancer by the time of their last core interview.

Please note that in the SHARE survey, the question wording specifically includes leukemia and lymphoma, whereas this distinction is not made in the HRS question wording.

For any differences in the construction of **rWcancre**, which is used to create **ralcancre**, please see "Doctor Diagnosed Health Problems: Ever Have Condition" in the Gateway Harmonized SHARE codebook.

Categorical Variable Frequencies

	ralcancre
0.no	17,306
1.yes	3,625
.d:dk	55
.m:missing	98
.r:refuse	14
Total	21,098

Original SHARE Variables Used

Harmonized SHARE:

r1cancre	r1cancre:w1 R ever had cancer
r2cancre	r2cancre:w2 R ever had cancer
r4cancre	r4cancre:w4 R ever had cancer
r5cancre	r5cancre:w5 R ever had cancer

Section B. Health

r6cancer	r6cancer:w6 R ever had cancer
r7cancer	r7cancer:w7 R ever had cancer
r8cancer	r8cancer:w8 R ever had cancer

Doctor Diagnosed Conditions: Lung Disease

Variable	Waves	Label	Type
rallunge	All	rallunge: r ever had lung disease as of last iwv	Categ

How Constructed

rallunge indicates whether the deceased respondent had ever been told by a doctor that they had chronic lung disease, such as chronic bronchitis or emphysema, in their last completed core interview. **rallunge** is based on **rWlunge** in the Gateway Harmonized SHARE and is taken from the last core wave completed. **rallunge** is coded as 0 if the respondent reported never having been diagnosed with chronic lung disease, and is coded as 1 if the respondent reported being diagnosed with chronic lung disease. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Cross Wave Differences in SHARE EOL

No differences known.

Differences with the Harmonized HRS End of Life

Unlike the HRS exit interview, the SHARE end of life interview does not ask proxies about the deceased respondent's chronic lung disease status by the time of their death. As such, we cannot provide variables indicating whether the respondent had been newly or ever diagnosed with chronic lung disease by the time of their death in the Gateway Harmonized SHARE End of Life as we do in the Harmonized HRS End of Life, we can only provide whether the respondent had ever been diagnosed with chronic lung disease by the time of their last core interview.

For any differences in the construction of **rWlunge**, which is used to create **rallunge**, please see "Doctor Diagnosed Health Problems: Ever Have Condition" in the Gateway Harmonized SHARE codebook.

Categorical Variable Frequencies

	rallunge
0.no	17,454
1.yes	3,477
.d:dk	54
.m:missing	97
.r:refuse	16
Total	21,098

Original SHARE Variables Used

Harmonized SHARE:

r1lunge	r1lunge:w1 R ever had lung disease
r2lunge	r2lunge:w2 R ever had lung disease
r4lunge	r4lunge:w4 R ever had lung disease
r5lunge	r5lunge:w5 R ever had lung disease
r6lunge	r6lunge:w6 R ever had lung disease
r7lunge	r7lunge:w7 R ever had lung disease
r8lunge	r8lunge:w8 R ever had lung disease

Doctor Diagnosed Conditions: Heart Conditions			
Variable	Waves	Label	Type
ralhearte	All	ralhearte: r ever had heart problems as of last iww	Categ
ralhrtatt	All	ralhrtatt: r had heart attack between first and last core iww	Categ

How Constructed

ralhearte indicates whether the deceased respondent had ever been told by a doctor that they had a heart attack, including myocardial infarction or coronary thrombosis, or any other heart problem, including congestive heart failure, in their last completed core interview. **ralhearte** is based on **rWhearte** in the Gateway Harmonized SHARE and is taken from the last core wave completed. **ralhearte** is coded as 0 if the respondent reported never having been diagnosed with heart problems, and is coded as 1 if the respondent reported being diagnosed with heart problems. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

ralhrtatt indicates whether the deceased respondent has been told by a doctor that they had a heart attack between their first and last completed core interviews. Starting in Wave 2 of the core interview, respondents who had been interviewed previously are asked if they had a heart attack since the last interview. **ralhrtatt** is based on **rWhrtatt** in the Gateway Harmonized SHARE and is taken from all core waves completed. **ralhrtatt** is coded as 0 if the respondent never reported having been diagnosed with a heart attack between their first and last completed core interviews, and is coded as 1 if the respondent reported being diagnosed with a heart attack between their first and last completed core interviews. If the respondent was new to the last completed core wave, meaning it was their baseline wave, then they are not asked whether they had a heart attack prior to their previous interview and **ralhrtatt** is assigned special missing (.b). The respondent is not asked whether they had a heart attack in the last two years in the Wave 1 core interview, and so **ralhrtatt** is assigned special missing (.q) if Wave 1 was the respondent's last core wave. If the respondent's last core interview was in Wave 7 and they were given the condensed questionnaire instead of the regular questionnaire because they participated in the Wave 7 Life History interview, then this question is not asked and **ralhrtatt** is assigned special missing (.c). Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Please note that in the SHARE core interview the respondent can dispute a previously positive report of heart problems. Please see **rWheartf** in "Doctor Diagnosed Health Problems: Ever Have Condition" in the Gateway Harmonized SHARE codebook for more information on the flag variable or to determine whether the report of heart problems has been disputed in a prior wave.

Cross Wave Differences in SHARE EOL

Respondents are not asked whether they have had a heart attack specifically in the Wave 1 core interview.

Respondents who completed the condensed interview in Wave 7 because they also completed the Wave 7 Life History interview are not asked whether they had a heart attack between their previous core interview and Wave 7.

Differences with the Harmonized HRS End of Life

Unlike the HRS exit interview, the SHARE end of life interview does not ask proxies about the deceased respondent's heart problem or heart attack status by the time of their death. As such, we cannot provide variables indicating whether the respondent had been newly or ever diagnosed with heart problems or heart attack by the time of their death in the Gateway Harmonized SHARE End of Life as we do in the Harmonized HRS End of Life, we can only provide whether the respondent had ever been diagnosed with a heart problem by the time of their last core interview and whether the respondent had ever been diagnosed with a heart attack between the respondent's first and last core interviews.

Please note that the SHARE question wording specifies "a heart attack including myocardial infarction or coronary thrombosis or any other heart problem including congestive heart failure", while the HRS question wording specifies "a heart attack, coronary heart disease, angina, congestive heart failure, or other heart problems".

For any differences in the construction of **rWhearte**, which is used to create **ralhearte**, please see "Doctor Diagnosed Health Problems: Ever Have Condition" in the Gateway Harmonized SHARE codebook. For any differences in the construction of **rWhrtatt**, which is used to create **ralhrtatt**, please see "Doctor Diagnosed Health Problems: Diagnosed Since Last Wave" in the Gateway Harmonized SHARE codebook.

Unlike in the Harmonized HRS End of Life, which has no variable indicating the respondent's heart attack status as of the last interview, **ralhrtatt** in the Gateway Harmonized SHARE End of Life indicates whether the respondent had a heart attack between their first and last core interviews.

Categorical Variable Frequencies

	ralhearte
0.no	13,389
1.yes	7,573
.d:dk	39
.m:missing	85
.r:refuse	12
Total	21,098

	ralhrtatt
0.no	9,590
1.yes	1,082
.b:baseline iww	5,136
.c:not asked condensed w7 i	2,558
.d:dk	17
.m:missing	1,121
.q:not asked this wave	1,591
.r:refuse	3
Total	21,098

Original SHARE Variables Used

Harmonized SHARE:

r1hearte	r1hearte:w1 R ever had heart problems
r2hearte	r2hearte:w2 R ever had heart problems
r4hearte	r4hearte:w4 R ever had heart problems
r5hearte	r5hearte:w5 R ever had heart problems
r6hearte	r6hearte:w6 R ever had heart problems
r7hearte	r7hearte:w7 R ever had heart problems
r8hearte	r8hearte:w8 R ever had heart problems
r2hrtatt	r2hrtatt:w2 R reports heart attack since last wave
r4hrtatt	r4hrtatt:w4 R reports heart attack since last wave
r5hrtatt	r5hrtatt:w5 R reports heart attack since last wave
r6hrtatt	r6hrtatt:w6 R reports heart attack since last wave
r7hrtatt	r7hrtatt:w7 R reports heart attack since last wave
r8hrtatt	r8hrtatt:w8 R reports heart attack since last wave

Doctor Diagnosed Conditions: Stroke

Variable	Waves	Label	Type
ralstroke	All	ralstroke: r ever had stroke as of last iwv	Categ

How Constructed

ralstroke indicates whether the deceased respondent had ever been told by a doctor that they had a stroke or cerebral vascular disease in their last completed core interview. **ralstroke** is based on **rWstroke** in the Gateway Harmonized SHARE and is taken from the last core wave completed. **ralstroke** is coded as 0 if the respondent reported never having been diagnosed with a stroke, and is coded as 1 if the respondent reported being diagnosed with a stroke. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Please note that in the SHARE core interview the respondent can dispute a previously positive report of a stroke. Please see **rWstrokef** in "Doctor Diagnosed Health Problems: Ever Have Condition" in the Gateway Harmonized SHARE codebook for more information on the flag variable or to determine whether the report of stroke has been disputed in a prior wave.

Cross Wave Differences in SHARE EOL

No differences known.

Differences with the Harmonized HRS End of Life

Unlike the HRS exit interview, the SHARE end of life interview does not ask proxies about the deceased respondent's stroke status by the time of their death. As such, we cannot provide variables indicating whether the respondent had been newly or ever diagnosed with a stroke by the time of their death in the Gateway Harmonized SHARE End of Life as we do in the Harmonized HRS End of Life, we can only provide whether the respondent had ever been diagnosed with a stroke by the time of their last core interview.

Please note that the SHARE question wording specifies "a stroke or cerebral vascular disease", while the HRS question wording specifies "a stroke".

For any differences in the construction of **rWstroke**, which is used to create **ralstroke**, please see "Doctor Diagnosed Health Problems: Ever Have Condition" in the Gateway Harmonized SHARE codebook.

Categorical Variable Frequencies

	ralstroke
0.no	17,470
1.yes	3,479
.d:dk	43
.m:missing	93
.r:refuse	13
Total	21,098

Original SHARE Variables Used

Harmonized SHARE:

r1stroke	r1stroke:w1 R ever had stroke
r2stroke	r2stroke:w2 R ever had stroke
r4stroke	r4stroke:w4 R ever had stroke
r5stroke	r5stroke:w5 R ever had stroke
r6stroke	r6stroke:w6 R ever had stroke

r7stroke
r8stroke

r7stroke:w7 R ever had stroke
r8stroke:w8 R ever had stroke

Doctor Diagnosed Conditions: Diabetes

Variable	Waves	Label	Type
raldiabe	All	raldiabe: r ever had diabetes as of last iwv	Categ

How Constructed

raldiabe indicates whether the deceased respondent had ever been told by a doctor that they had diabetes or high blood sugar, in their last completed core interview. **raldiabe** is based on **rWdiabe** in the Gateway Harmonized SHARE and is taken from the last core wave completed. **raldiabe** is coded as 0 if the respondent reported never having been diagnosed with diabetes, and is coded as 1 if the respondent reported being diagnosed with diabetes. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Cross Wave Differences in SHARE EOL

No differences known.

Differences with the Harmonized HRS End of Life

Unlike the HRS exit interview, the SHARE end of life interview does not ask proxies about the deceased respondent's diabetes status by the time of their death. As such, we cannot provide variables indicating whether the respondent had been newly or ever diagnosed with diabetes by the time of their death in the Gateway Harmonized SHARE End of Life as we do in the Harmonized HRS End of Life, we can only provide whether the respondent had ever been diagnosed with diabetes by the time of their last core interview.

For any differences in the construction of **rWdiabe**, which is used to create **raldiabe**, please see "Doctor Diagnosed Health Problems: Ever Have Condition" in the Gateway Harmonized SHARE codebook.

Categorical Variable Frequencies

	raldiabe
0.no	15,546
1.yes	5,401
.d:dk	49
.m:missing	91
.r:refuse	11
Total	21,098

Original SHARE Variables Used

Harmonized SHARE:

r1diabe	r1diabe:w1 R ever had diabetes
r2diabe	r2diabe:w2 R ever had diabetes
r4diabe	r4diabe:w4 R ever had diabetes
r5diabe	r5diabe:w5 R ever had diabetes
r6diabe	r6diabe:w6 R ever had diabetes
r7diabe	r7diabe:w7 R ever had diabetes
r8diabe	r8diabe:w8 R ever had diabetes

Doctor Diagnosed Conditions: High Blood Pressure

Variable	Waves	Label	Type
ralhibpe	All	ralhibpe: r ever had high bp as of last iww	Categ

How Constructed

ralhibpe indicates whether the deceased respondent had ever been told by a doctor that they had high blood pressure, in their last completed core interview. **ralhibpe** is based on **rWhibpe** in the Gateway Harmonized SHARE and is taken from the last core wave completed. **ralhibpe** is coded as 0 if the respondent reported never having been diagnosed with high blood pressure, and is coded as 1 if the respondent reported being diagnosed with high blood pressure. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Cross Wave Differences in SHARE EOL

No differences known.

Differences with the Harmonized HRS End of Life

Unlike the HRS exit interview, the SHARE end of life interview does not ask proxies about the deceased respondent's high blood pressure status by the time of their death. As such, we cannot provide variables indicating whether the respondent had been newly or ever diagnosed with high blood pressure by the time of their death in the Gateway Harmonized SHARE End of Life as we do in the Harmonized HRS End of Life, we can only provide whether the respondent had ever been diagnosed with high blood pressure by the time of their last core interview.

For any differences in the construction of **rWhibpe**, which is used to create **ralhibpe**, please see "Doctor Diagnosed Health Problems: Ever Have Condition" in the Gateway Harmonized SHARE codebook.

Categorical Variable Frequencies

	ralhibpe
0.no	8,350
1.yes	12,628
.d:dk	30
.m:missing	77
.r:refuse	13
Total	21,098

Original SHARE Variables Used

Harmonized SHARE:

r1hibpe	r1hibpe:w1 R ever had high blood pressure
r2hibpe	r2hibpe:w2 R ever had high blood pressure
r4hibpe	r4hibpe:w4 R ever had high blood pressure
r5hibpe	r5hibpe:w5 R ever had high blood pressure
r6hibpe	r6hibpe:w6 R ever had high blood pressure
r7hibpe	r7hibpe:w7 R ever had high blood pressure
r8hibpe	r8hibpe:w8 R ever had high blood pressure

Doctor Diagnosed Conditions: Memory Problems

Variable	Waves	Label	Type
ralmemrye	All	ralmemrye: r ever had memory problems as of last iww	Categ

How Constructed

ralmemrye indicates whether the deceased respondent had ever been told by a doctor that they had Alzheimer's disease, dementia, or senility, in their last completed core interview. **ralmemrye** is based on **rWalzdeme** in the Gateway Harmonized SHARE and is taken from the last core wave completed. **ralmemrye** is coded as 0 if the respondent reported never having been diagnosed with Alzheimer's disease, dementia, or senility, and is coded as 1 if the respondent reported being diagnosed with Alzheimer's disease, dementia, or senility. Questions on memory diseases were not asked in Wave 1, so if the respondent's last core wave was in Wave 1, then **ralmemrye** is assigned special missing (.q). Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Cross Wave Differences in SHARE EOL

No question on memory disease is asked in the SHARE core interview in Wave 1.

Differences with the Harmonized HRS End of Life

Unlike the HRS exit interview, the SHARE end of life interview does not ask proxies about the deceased respondent's memory disease status by the time of their death. As such, we cannot provide variables indicating whether the respondent had been newly or ever diagnosed with memory disease by the time of their death in the Gateway Harmonized SHARE End of Life as we do in the Harmonized HRS End of Life, we can only provide whether the respondent had ever been diagnosed with memory disease by the time of their last core interview.

Please note that the SHARE question wording specifies "Alzheimer's disease, dementia, organic brain syndrome, senility or any other serious memory impairment" in a single question, while the HRS question wording specifies a "memory-related disease" in Waves 4 through 9, and then specifies "Alzheimer's disease" and "dementia, senility or any other serious memory impairment" in two separate questions in Waves 10 and onward.

For any differences in the construction of **rWalzdeme**, which is used to create **ralmemrye**, please see "Doctor Diagnosed Health Problems: Alzheimer's Disease or Dementia" in the Gateway Harmonized SHARE codebook.

Categorical Variable Frequencies

	ralmemrye
0.no	16,838
1.yes	2,524
.d:dk	45
.m:missing	85
.q:not asked this wave	1,591
.r:refuse	15
Total	21,098

Original SHARE Variables Used

Harmonized SHARE:

r2alzdeme	r2alzdeme:w2 R ever had alzheimer's/dementia
r4alzdeme	r4alzdeme:w4 R ever had alzheimer's/dementia
r5alzdeme	r5alzdeme:w5 R ever had alzheimer's/dementia
r6alzdeme	r6alzdeme:w6 R ever had alzheimer's/dementia

Section B. Health

r7alzdeme
r8alzdeme

r7alzdeme:w7 R ever had alzheimer's/dementia
r8alzdeme:w8 R ever had alzheimer's/dementia

C. Health Care Utilization and Insurance

Hospital Stays

Variable	Waves	Label	Type
raxhosp1y	All	raxhosp1y: r any hospital stay prev yr	Categ

How Constructed

raxhosp1y indicates whether the deceased respondent received care at a hospital in the last 12 months before death. The proxy is first asked on how many different occasions the respondent stayed in a hospital, hospice, or nursing home in the last year before they died. In Waves 2 through 5 and starting in Wave 8, if the respondent stayed in one of these locations at least once, and if the respondent stayed in one of these locations at least twice in Waves 6 and 7, then the proxy is asked whether the respondent had any hospital stays. **raxhosp1y** is coded as 0 if the proxy reports that the respondent did not have any hospital stays in the last 12 months before death. **raxhosp1y** is coded as 1 if the proxy reports that the respondent did have a hospital stay in the last 12 months before death. In Waves 2 through 8, if the respondent's death was the result of an accident, then the proxy is not asked these questions and **raxhosp1y** is assigned special missing (.a). In Waves 6 and 7, **raxhosp1y** is assigned special missing (.s) if the proxy reports a single stay in a hospital, hospice, or nursing home in the first question, and the proxy is not asked further whether the respondent had any hospital stays. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Cross Wave Differences in SHARE EOL

In Waves 2 through 5, the proxy is asked to select the number of stays at a hospital, hospice, or nursing home in the last year before death from the following choices: 0 times, 1-2 times, 3-5 times, more than 5 times. Starting in Wave 6, the proxy is asked to directly report the number of stays at a hospital, hospice, or nursing home in the last year before death.

In Waves 2 through 8, if the respondent's death was the result of an accident, then the proxy is not asked about the number of hospital stays. Starting in Wave 9 the proxy is asked about the number of hospital stays regardless of the cause of death.

In Waves 6 and 7, the proxy is asked to directly report the number of stays at a hospital, hospice, or nursing home in the last year before death. In Waves 6 and 7, the proxy is not asked whether the respondent had any hospital stays if the proxy reported that the respondent stayed in a hospital, hospice, or nursing home only once or not at all. In this case, **raxhosp1y** is assigned a special missing (.s) if the proxy reported one stay.

Differences with the Harmonized HRS End of Life

The HRS exit interview inquires about hospital stays in the last 2 years before death or since the previous interview, which is coded in **raxhosp** in the Harmonized HRS End of life. The SHARE end of life interview inquires about hospital stays in the last 12 months before death, which is coded in **raxhosp1y** in the Gateway Harmonized SHARE End of Life. In the SHARE end of life interview, this question is not asked if the respondent's death was the result of an accident, whereas the HRS exit interview does not limit who is asked this question based on the cause of death. Additionally, the HRS exit interview asks this question slightly differently depending on whether the respondent died in the hospital or not, while it is asked the same whether or not the respondent died in a hospital in the SHARE end of life interview. Please see the Harmonized HRS End of Life codebook for more information about the differences in question wording in the HRS exit interview.

Categorical Variable Frequencies

	raxhosp1y
0.no	7,183
1.yes	10,798
.a:accident caused death	247
.d:dk	806
.m:missing	119
.r:refuse	101
.s:single stay	1,844

Total	21,098
-------	--------

Original SHARE Variables Used

Wave 2 End of Life:

xt011_	Main cause of death
xt015_	Times in hospital last year before dying
xt018_3	Type of medical care in last 12 months: hospital stays

Wave 3 End of Life:

sl_xt011_	Main cause of death
sl_xt015_	Times in hospital last year before dying
sl_xt018_3	Had medical care in last 12 months: hospital stays

Wave 4 End of Life:

xt011_	Main cause of death
xt015_	Times in hospital last year before dying
xt018_3	Type of medical care in last 12 months: hospital stays

Wave 5 End of Life:

xt011_	Main cause of death
xt015_	Times in hospital last year before dying
xt018_3	Type of medical care in last 12 months: hospital stays

Wave 6 End of Life:

xt011_	Main cause of death
xt615_	Times in hospital last year before dying
xt018_3	Type of medical care in last 12 months: hospital stays

Wave 7 End of Life:

xt011_	Main cause of death
xt615_	Times in hospital last year before dying
xt018_3	Type of medical care in last 12 months: hospital stays

Wave 8 End of Life:

xt011_	Main cause of death
xt615_	Times in hospital last year before dying
xt018_3	Type of medical care in last 12 months: hospital stays

Wave 9 End of Life:

xt011_	Main cause of death
xt615_	Times in hospital last year before dying
xt018_3	Type of medical care in last 12 months: hospital stays

Wave 9 COVID End of Life:

xt011_	Main cause of death
xt615_	Times in hospital last year before dying
xt018_3	Type of medical care in last 12 months: hospital stays

Nursing Home Stays: In the Last Year

Variable	Waves	Label	Type
raxnrshom1y	All	raxnrshom1y: r any nursing home stay prev yr	Categ

How Constructed

raxnrshom1y indicates whether the deceased respondent received care in a nursing home in the last 12 months before death. The proxy is first asked on how many different occasions the respondent stayed in a hospital, hospice, or nursing home in the last year before they died. In Waves 2 through 5 and starting in Wave 8, if the respondent stayed in one of these locations at least once, and if the respondent stayed in one of these locations at least twice in Waves 6 and 7, then the proxy is asked whether the respondent received care in a nursing home. **raxnrshom1y** is coded as 0 if the proxy reports that the respondent did not receive care in a nursing home in the last 12 months before death. **raxnrshom1y** is coded as 1 if the proxy reports that the respondent did receive care in a nursing home in the last 12 months before death. In Waves 2 through 8, if the respondent's death was the result of an accident, then the proxy is not asked these questions and **raxnrshom1y** is assigned special missing (.a). In Waves 6 and 7, **raxnrshom1y** is assigned special missing (.s) if the proxy reports a single stay in a hospital, hospice, or nursing home in the first question, and the proxy is not asked further whether the respondent received care in a nursing home. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Cross Wave Differences in SHARE EOL

In Waves 2 through 5, the proxy is asked to select the number of stays at a hospital, hospice, or nursing home in the last year before death from the following choices: 0 times, 1-2 times, 3-5 times, more than 5 times. Starting in Wave 6, the proxy is asked to directly report the number of stays at a hospital, hospice, or nursing home in the last year before death.

In Waves 2 through 8, if the respondent's death was the result of an accident, then the proxy is not asked about the number of hospital, hospice, or nursing home stays. Starting in Wave 9 the proxy is asked about the number of hospital, hospice, or nursing home stays regardless of the cause of death.

In Waves 6 and 7, the proxy is not asked whether the respondent had any nursing home stays if the proxy reported that the respondent stayed in a hospital, hospice, or nursing home only once or not at all. In this case, **raxnrshom1y** is assigned a special missing (.s) if the proxy reported one stay.

Differences with the Harmonized HRS End of Life

The HRS exit interview inquires about nursing home stays in the last 2 years before death or since the previous interview, which is coded in **raxnrshom** in the Harmonized HRS End of life. The SHARE end of life interview inquires about nursing home stays in the last 12 months before death, which is coded in **raxnrshom1y** in the Gateway Harmonized SHARE End of Life. In the SHARE end of life interview, this question is not asked if the respondent's death was the result of an accident, whereas the HRS exit interview does not limit who is asked this question based on the cause of death. The HRS exit interview inquires about nursing home, convalescent home, or other long-term health care facility stays as a single entity, while the SHARE end of life interview inquires only about a nursing home. Additionally, the HRS exit interview asks this question slightly differently depending on whether the respondent died in a nursing home or not, while it is asked the same whether or not the respondent died in a nursing home in the SHARE end of life interview. Please see the Harmonized HRS End of Life codebook for more information about the differences in question wording in the HRS exit interview.

Categorical Variable Frequencies

	raxnrshom1y
0.no	15,347
1.yes	2,648
.a:accident caused death	247
.d:dk	792

.m:missing	117
.r:refuse	103
.s:single stay	1,844
Total	21,098

Original SHARE Variables Used

Wave 2 End of Life:

xt011_	Main cause of death
xt015_	Times in hospital last year before dying
xt018_4	Type of medical care in last 12 months: care in a nursing home

Wave 3 End of Life:

sl_xt011_	Main cause of death
sl_xt015_	Times in hospital last year before dying
sl_xt018_4	Had medical care in last 12 months: care in a nursing home

Wave 4 End of Life:

xt011_	Main cause of death
xt015_	Times in hospital last year before dying
xt018_4	Type of medical care in last 12 months: care in a nursing home

Wave 5 End of Life:

xt011_	Main cause of death
xt015_	Times in hospital last year before dying
xt018_4	Type of medical care in last 12 months: care in a nursing home

Wave 6 End of Life:

xt011_	Main cause of death
xt615_	Times in hospital last year before dying
xt018_4	Type of medical care in last 12 months: care in a nursing home

Wave 7 End of Life:

xt011_	Main cause of death
xt615_	Times in hospital last year before dying
xt018_4	Type of medical care in last 12 months: care in a nursing home

Wave 8 End of Life:

xt011_	Main cause of death
xt615_	Times in hospital last year before dying
xt018_4	Type of medical care in last 12 months: care in a nursing home

Wave 9 End of Life:

xt011_	Main cause of death
xt615_	Times in hospital last year before dying
xt018_4	Type of medical care in last 12 months: care in a nursing home

Wave 9 COVID End of Life:

xt011_	Main cause of death
xt615_	Times in hospital last year before dying
xt018_4	Type of medical care in last 12 months: care in a nursing home

Hospice Stays: In the Last Year			
Variable	Waves	Label	Type
raxhospice1y	All	raxhospice1y: r any hospice stay prev yr	Categ

How Constructed

raxhospice1y indicates whether the deceased respondent had any hospice stays in the last 12 months before death. The proxy is first asked on how many different occasions the respondent stayed in a hospital, hospice, or nursing home in the last year before they died. In Waves 2 through 5 and starting in Wave 8, if the respondent stayed in one of these locations at least once, and if the respondent stayed in one of these locations at least twice in Waves 6 and 7, then the proxy is asked whether the respondent had any hospice stays. **raxhospice1y** is coded as 0 if the proxy reports that the respondent did not have any hospice stays in the last 12 months before death. **raxhospice1y** is coded as 1 if the proxy reports that the respondent did have a hospice stay in the last 12 months before death. In Waves 2 through 8, if the respondent's death was the result of an accident, then the proxy is not asked these questions and **raxhospice1y** is assigned special missing (.a). In Waves 6 and 7, **raxhospice1y** is assigned special missing (.s) if the proxy reports a single stay in a hospital, hospice, or nursing home in the first question, and the proxy is not asked further whether the respondent had any hospice stays. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Cross Wave Differences in SHARE EOL

In Waves 2 through 5, the proxy is asked to select the number of stays at a hospital, hospice, or nursing home in the last year before death from the following choices: 0 times, 1-2 times, 3-5 times, more than 5 times. Starting in Wave 6, the proxy is asked to directly report the number of stays at a hospital, hospice, or nursing home in the last year before death.

In Waves 2 through 8, if the respondent's death was the result of an accident, then the proxy is not asked about the number of hospital, hospice, or nursing home stays. Starting in Wave 9 the proxy is asked about the number of hospital, hospice, or nursing home stays regardless of the cause of death.

In Waves 6 and 7, the proxy is not asked whether the respondent had any hospice stays if the proxy reported that the respondent stayed in a hospital, hospice, or nursing home only once or not at all. In this case, **raxhospice1y** is assigned a special missing (.s) if the proxy reported one stay.

Differences with the Harmonized HRS End of Life

The HRS exit interview inquires about hospice stays in the last 2 years before death or since the previous interview, which is coded in **raxhospice** in the Harmonized HRS End of Life. The SHARE end of life interview inquires about hospice stays in the last 12 months before death, which is coded in **raxhospice1y** in the Gateway Harmonized SHARE End of Life. In the SHARE end of life interview, this question is not asked if the respondent's death was the result of an accident, whereas the HRS exit interview does not limit who is asked this question based on the cause of death. Additionally, the HRS exit interview asks this question slightly differently depending on whether the respondent died in a hospice or not, while it is asked the same whether or not the respondent died in a hospice in the SHARE end of life interview. In later waves, the HRS exit interview stops asking questions about hospice stays and instead asks about the use of hospice services in the last 2 years before death or since the previous interview, which is coded in **raxhpcserv** in the Harmonized HRS End of Life. Please see the Harmonized HRS End of Life codebook for more information about the differences in question wording in the HRS exit interview.

Categorical Variable Frequencies

	raxhospice1y
0.no	16,646
1.yes	1,292
.a:accident caused death	246
.d:dk	840
.m:missing	124

.r:refuse	106
.s:single stay	1,844
Total	21,098

Original SHARE Variables Used

Wave 2 End of Life:

xt011_	Main cause of death
xt015_	Times in hospital last year before dying
xt018_5	Type of medical care in last 12 months: hospice stays

Wave 3 End of Life:

sl_xt011_	Main cause of death
sl_xt015_	Times in hospital last year before dying
sl_xt018_5	Had medical care in last 12 months: hospice stays

Wave 4 End of Life:

xt011_	Main cause of death
xt015_	Times in hospital last year before dying
xt018_5	Type of medical care in last 12 months: hospice stays

Wave 5 End of Life:

xt011_	Main cause of death
xt015_	Times in hospital last year before dying
xt018_5	Type of medical care in last 12 months: hospice stays

Wave 6 End of Life:

xt011_	Main cause of death
xt615_	Times in hospital last year before dying
xt018_5	Type of medical care in last 12 months: hospice stays

Wave 7 End of Life:

xt011_	Main cause of death
xt615_	Times in hospital last year before dying
xt018_5	Type of medical care in last 12 months: hospice stays

Wave 8 End of Life:

xt011_	Main cause of death
xt615_	Times in hospital last year before dying
xt018_5	Type of medical care in last 12 months: hospice stays

Wave 9 End of Life:

xt011_	Main cause of death
xt615_	Times in hospital last year before dying
xt018_5	Type of medical care in last 12 months: hospice stays

Wave 9 COVID End of Life:

xt011_	Main cause of death
xt615_	Times in hospital last year before dying
xt018_5	Type of medical care in last 12 months: hospice stays

Hospice or Palliative Care: In the Last 4 Weeks

Variable	Waves	Label	Type
raxhsplc1m	All	raxhsplc1m: r recd hospice or palliative care last 4 wks	Cont

How Constructed

raxhsplc1m indicates whether the deceased respondent received any hospice or palliative care in the last 4 weeks before death. Starting in Wave 7, the proxy is asked whether the respondent received any hospice or palliative care in the last four weeks before death. In Waves 7 and 8, **raxhsplc1m** is coded as 1 if the proxy reports that the deceased respondent received any hospice or palliative care in the last 4 weeks before death; or if the proxy reports that the respondent died in a palliative care unit, inpatient hospital unit, or residential housing provided by hospice. In Wave 9, **raxhsplc1m** is coded as 1 if the proxy reports that the deceased respondent received any hospice or palliative care in the last 4 weeks before death; or if the proxy reports that the respondent died in a palliative care or inpatient hospital unit. **raxhsplc1m** is coded as 0 if the proxy reports that the respondent did not receive any hospice or palliative care in the last 4 weeks before death, or if the proxy reports that the respondent did not die in a hospice, palliative care unit, inpatient hospice unit, or housing provided by hospice. In Waves 2 through 6, **raxhsplc1m** is assigned special missing (.q) because the proxy is not asked whether the deceased respondent received any hospice or palliative care in the last 4 weeks before death. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Cross Wave Differences in SHARE EOL

Questions about hospice or palliative care in the last 4 weeks before death were not asked in SHARE until Wave 7. In Waves 7 and 8, the proxy is asked if the respondent died in a palliative care or inpatient hospital unit, or in an inpatient hospice unit, or in residential housing provided by hospice. Beginning in Wave 9, the proxy is asked only if the respondent died in a palliative care or inpatient hospital unit.

Differences with the Harmonized HRS End of Life

In the HRS exit interview, there is no specific question about hospice or palliative care received in the last 4 weeks of life. In HRS Waves 2-11, the proxy is asked about the use of a hospice facility. Starting in HRS Wave 12, the proxy is asked if the deceased respondent received hospice services. Beginning in Wave 14, the proxy is asked if the deceased respondent received palliative care in the months or years prior to death. Please see the Harmonized HRS End of Life codebook for more information about the differences in question wording in the HRS exit interview.

Continuous Variable Distributions

Variable	N	Mean	Std.Dev	Minimum	Maximum	Total missing
raxhsplc1m	12,208	0.33	0.47	0.00	1.00	8,890

Original SHARE Variables Used

Wave 7 End of Life:

xt014_	Place of dying
xt757_	Hospice or palliative care
xt751_	Palliative care or inpatient hospice unit
xt752_	Inpatient hospice unit
xt753_	Residential provided by hospice

Wave 8 End of Life:

xt014_	Place of dying
xt757_	Hospice or palliative care
xt751_	Palliative care or inpatient hospice unit
xt752_	Inpatient hospice unit

xt753_	Residential provided by hospice
Wave 9 End of Life:	
xt014_	Place of dying
xt757_	Hospice or palliative care
xt751_	Palliative care or inpatient hospice unit
Wave 9 COVID End of Life:	
xt014_	Place of dying
xt757_	Hospice or palliative care
xt751_	Palliative care or inpatient hospice unit

Hospital, Nursing Home, or Hospice Stays			
Variable	Waves	Label	Type
raxhnh1y	All	raxhnh1y: r any hospital, hospice, nursing home stay prev yr	Categ
raxhntim1y	All	raxhntim1y: r # hospital, hospice, nursing home stays grouped prev yr	Categ
raxhntim1y	All	raxhntim1y: r # hospital, hospice, nursing home stays prev yr	Cont
raxhnnit1y	All	raxhnnit1y: r total duration hospital, hospice, nursing home stays prev yr	Categ

How Constructed

raxhnh1y indicates whether the deceased respondent stayed at a hospital, hospice, or nursing home in the last year before death. The proxy is asked on how many different occasions the respondent stayed in a hospital, hospice, or nursing home. **raxhnh1y** is coded as 0 if the proxy reports that the respondent never stayed in a hospital, hospice, or nursing home in the last year before death. **raxhnh1y** is coded as 1 if the proxy reports that the respondent stayed at least once in a hospital, hospice, or nursing home in the last year before death. In Waves 2 through 8, if the respondent's death was the result of an accident, then the proxy is not asked these questions and **raxhnh1y** is assigned special missing (.a). Starting in Wave 6, if the number of occasions provided was greater than 365, then **raxhnh1y** is assigned special missing (.i) in this wave. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

raxhntim1y and **raxhntim1y** indicate the number of times the deceased respondent stayed at a hospital, hospice, or nursing home in the last year before death. **raxhntim1y** is coded as follows: 0.0 times, 1.1-2 times, 2.3-5 times, 3.more than 5 times. In Waves 2 through 5, the proxy is asked to select the number of stays at a hospital, hospice, or nursing home in the last year before death from these choices. Starting in Wave 6, the proxy is asked to directly report the number of stays at a hospital, hospice, or nursing home in the last year before death, which is recorded in **raxhntim1y**. Starting in Wave 6, the responses provided in **raxhntim1y** are grouped to assign values to **raxhntim1y**. In Waves 2 through 8, if the respondent's death was the result of an accident, then the proxy is not asked these questions and **raxhntim1y** and **raxhntim1y** are assigned special missing (.a). In Waves 2 through 5, **raxhntim1y** is assigned special missing (.q) because the proxy cannot respond with continuous values in these waves. Starting in Wave 6, if the number of occasions provided was greater than 365, then **raxhntim1y** and **raxhntim1y** are assigned special missing (.i). Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

raxhnnit1y indicates how long the deceased respondent stayed in a hospital, hospice, or nursing home in total in the last year before death. In Waves 2 through 5, the proxy is asked how long altogether the respondent stayed at hospitals, hospices, or nursing homes during the last year of their life as long as the proxy reported at least one stay. Starting in Wave 6, if the proxy reported that the respondent had at least one stay in a hospital, hospice, or nursing home in the last year before death, then the proxy is asked how long altogether the respondent stayed at hospitals, hospices, or nursing homes during the last year of their life. The proxy **raxhnnit1y** is coded as follows: 0.0 nights, 1.less than 1 week, 2.1 week or more but less than 1 month, 3.1 month or more but less than 3 months, 4.3 months or more but less than 6 months, 5.6 months or more but less than a year, 6.a full year. If the proxy reported that the respondent had not stayed at all in a hospital, hospice, or nursing home, then **raxhnnit1y** is set to 0. In Waves 2 through 8, if the respondent's death was the result of an accident, then the proxy is not asked these questions and **raxhnnit1y** is assigned special missing (.a). In Wave 6, if the respondent had a single stay at a hospital, hospice, or nursing home and this question is not asked, then **raxhnnit1y** is assigned special missing (.s). Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Cross Wave Differences in SHARE EOL

In Waves 2 through 5, the proxy is asked to select the number of stays at a hospital, hospice, or nursing home in the last year before death from the following choices: 0 times, 1-2 times, 3-5 times, more than 5 times, which is recorded in **raxhntim1y**. Starting in Wave 6, the proxy is asked to directly report the number of stays at a hospital, hospice, or nursing home in the last year before death, which is recorded in **raxhntim1y**.

ing home in the last year before death, which is recorded in **raxhhntim1y**. Starting in Wave 6, the responses provided in **raxhhntim1y** are grouped to assign values to **raxhhntimd1y**.

In Waves 2 through 5, the proxy is asked how long altogether the respondent stayed at hospitals, hospices, or nursing homes during the last year of their life as long as the proxy reported at least one stay. In Wave 6, if the proxy reported that the respondent had more than one stay in a hospital, hospice, or nursing home in the last year before death, then the proxy is asked to report how long altogether the respondent stayed at hospitals, hospices, or nursing homes during the last year of their life, which is recorded in **raxhhnntd1y**. As a result, a special missing (.s) is assigned to **raxhhnntd1y** in Wave 6 for those who had a single stay. Starting in Wave 7, if the proxy reported that the respondent had at least one stay in a hospital, hospice, or nursing home in the last year before death, then the proxy is asked to report how long altogether the respondent stayed at hospitals, hospices, or nursing homes during the last year of their life.

In Waves 2 through 8, if the respondent's death was the result of an accident, then the proxy is not asked any questions about hospital, hospice, or nursing home stays. Starting in Wave 9 the proxy is asked about hospital, hospice, and nursing home stays regardless of the cause of death.

Differences with the Harmonized HRS End of Life

The HRS exit interview inquires about hospital, hospice, and nursing home stays separately for the last 2 years before death or since the previous interview, while the SHARE end of life interview inquires about hospital, hospice, and nursing home stays together for the last year before death. In the SHARE end of life interview, this question is not asked if the respondent's death was the result of an accident, whereas the HRS exit interview does not limit who is asked this question based on the cause of death. Additionally, the HRS exit interview asks the question for each location slightly differently depending on whether the respondent died in that location, while it is asked the same whether or not the respondent died in a hospital, hospice, or nursing home in the SHARE end of life interview. Please see the Harmonized HRS End of Life codebook for more information about the differences in question wording in the HRS exit interview.

In the HRS exit interview, the proxy is first asked whether the respondent had any stays at that location, followed by the number of stays, and then the total length of stay. The number of stays at each location takes continuous values in all waves of the Harmonized HRS End of Life, whereas it takes categorical and continuous values in the Gateway Harmonized SHARE End of Life. Also, the total length of stay is asked in various ways but with generally more detail in the HRS exit interview than the categories available in the SHARE end of life interview.

Continuous Variable Distributions

Variable	N	Mean	Std.Dev	Minimum	Maximum	Total missing
raxhhntim1y	14,550	4.54	28.73	0.00	365.00	6,548

Categorical Variable Frequencies

	raxhhnh1y
0.no	6,041
1.yes	13,493
.a:accident caused death	340
.d:dk	947
.i:invalid	27
.m:missing	139
.r:refuse	111
Total	21,098
	raxhhntimd1y
0.0 times	6,041
1.1-2 times	8,567
2.3-5 times	3,157

3.more than 5 times	1,769
.a:accident caused death	340
.d:dk	947
.i:records inaccurate	27
.m:missing	139
.r:refuse	111
Total	21,098
raxhhnnitd1y	
0.0 nights	6,059
1.less than 1 week	2,182
2.1 week to lt 1 month	4,563
3.1 month to lt 3 months	3,238
4.3 months to lt 6 months	1,166
5.6 months to lt 1 year	553
6.1 year	827
.a:accident caused death	340
.d:dk	1,016
.m:missing	153
.r:refuse	123
.s:single stay	878
Total	21,098

Original SHARE Variables Used

Wave 2 End of Life:

xt011_	Main cause of death
xt015_	Times in hospital last year before dying
xt016_	Total time in hospital last year before dying

Wave 3 End of Life:

sl_xt011_	Main cause of death
sl_xt015_	Times in hospital last year before dying
sl_xt016_	Total time in hospital last year before dying

Wave 4 End of Life:

xt011_	Main cause of death
xt015_	Times in hospital last year before dying
xt016_	Total time in hospital last year before dying

Wave 5 End of Life:

xt011_	Main cause of death
xt015_	Times in hospital last year before dying
xt016_	Total time in hospital last year before dying

Wave 6 End of Life:

xt011_	Main cause of death
xt615_	Times in hospital last year before dying
xt016_	Total time in hospital last year before dying

Wave 7 End of Life:

xt011_	Main cause of death
xt615_	Times in hospital last year before dying
xt016_	Total time in hospital last year before dying

Wave 8 End of Life:

xt011_	Main cause of death
--------	---------------------

xt615_	Times in hospital last year before dying
xt016_	Total time in hospital last year before dying
Wave 9 End of Life:	
xt011_	Main cause of death
xt615_	Times in hospital last year before dying
xt016_	Total time in hospital last year before dying
Wave 9 COVID End of Life:	
xt011_	Main cause of death
xt615_	Times in hospital last year before dying
xt016_	Total time in hospital last year before dying

Doctor Visits: In the Last Year			
Variable	Waves	Label	Type
raxdoctor1y	All	raxdoctor1y: r any doctor visits prev yr	Categ

How Constructed

raxdoctor1y indicates whether the deceased respondent had any doctor visits in the last 12 months before death. **raxdoctor1y** is coded as 0 if the proxy reports that the respondent did not have any visits with a general practitioner or specialist physician in the last 12 months before death. **raxdoctor1y** is coded as 1 if the proxy reports that the respondent did have a visit with a general practitioner or specialist physician in the last 12 months before death. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Cross Wave Differences in SHARE EOL

No differences known.

Differences with the Harmonized HRS End of Life

The HRS asks how many times the respondent saw or talked to a medical doctor since the previous wave or in the last two years, whereas SHARE asks whether the respondent had any care from a general practitioner or from a specialist physician separately in the last 12 months of life. As such, the Harmonized HRS End of Life includes **raxdoctor** and **raxdoctim**, while the Gateway Harmonized SHARE End of Life includes **raxdoctor1y**.

Categorical Variable Frequencies

	raxdoctor1y
0.no	2,077
1.yes	18,566
.d:dk	298
.m:missing	124
.r:refuse	33
Total	21,098

Original SHARE Variables Used

Wave 2 End of Life:

xto18_1	Type of medical care in last 12 months: care from a general practitioner
xto18_2	Type of medical care in last 12 months: care from specialist physicians

Wave 3 End of Life:

sl_xto18_1	Had medical care in last 12 months: care from a general practitioner
sl_xto18_2	Had medical care in last 12 months: care from specialist physicians

Wave 4 End of Life:

xto18_1	Type of medical care in last 12 months: care from a general practitioner
xto18_2	Type of medical care in last 12 months: care from specialist physicians

Wave 5 End of Life:

xto18_1	Type of medical care in last 12 months: care from a general practitioner
xto18_2	Type of medical care in last 12 months: care from specialist physicians

Wave 6 End of Life:

xto18_1	Type of medical care in last 12 months: care from a general practitioner
xto18_2	Type of medical care in last 12 months: care from specialist physicians

Wave 7 End of Life:

xt018_1

Type of medical care in last 12 months: care from a general practitioner

xt018_2

Type of medical care in last 12 months: care from specialist physicians

Wave 8 End of Life:

xt018_1

Type of medical care in last 12 months: care from a general practitioner

xt018_2

Type of medical care in last 12 months: care from specialist physicians

Wave 9 End of Life:

xt018_1

Type of medical care in last 12 months: care from a general practitioner

xt018_2

Type of medical care in last 12 months: care from specialist physicians

Wave 9 COVID End of Life:

xt018_1

Type of medical care in last 12 months: care from a general practitioner

xt018_2

Type of medical care in last 12 months: care from specialist physicians

Medication Use: In the Last Year			
Variable	Waves	Label	Type
raxdrugs1y	All	raxdrugs1y: r any medication use prev yr	Categ

How Constructed

raxdrugs1y indicates whether the deceased respondent used medication in the last 12 months before death. **raxdrugs1y** is coded as 0 if the proxy reports that the respondent did not use medication in the last 12 months before death. **raxdrugs1y** is coded as 1 if the proxy reports that the respondent did use medication in the last 12 months before death. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Cross Wave Differences in SHARE EOL

No differences known.

Differences with the Harmonized HRS End of Life

The HRS asks whether the respondent regularly took medication before their death, whereas SHARE asks whether the respondent used medication in the last 12 months of life. As such, the Harmonized HRS End of Life includes **raxdrugs**, while the Gateway Harmonized SHARE End of Life includes **raxdrugs1y**.

Categorical Variable Frequencies

	raxdrugs1y
0.no	1,607
1.yes	18,955
.d.dk	365
.m:missing	127
.r:refuse	44
Total	21,098

Original SHARE Variables Used

Wave 2 End of Life:	
xto18_6	Type of medical care in last 12 months: medication
Wave 3 End of Life:	
sl_xto18_6	Had medical care in last 12 months: medication
Wave 4 End of Life:	
xto18_6	Type of medical care in last 12 months: medication
Wave 5 End of Life:	
xto18_6	Type of medical care in last 12 months: medication
Wave 6 End of Life:	
xto18_6	Type of medical care in last 12 months: medication
Wave 7 End of Life:	
xto18_6	Type of medical care in last 12 months: medication
Wave 8 End of Life:	
xto18_6	Type of medical care in last 12 months: medication
Wave 9 End of Life:	
xto18_6	Type of medical care in last 12 months: medication

Wave 9 COVID End of Life:

xt018_6

Type of medical care in last 12 months: medication

Home Care: In the Last Year			
Variable	Waves	Label	Type
raxhomcar1y	All	raxhomcar1y: r any home care prev yr	Categ

How Constructed

raxhomcar1y indicates whether the deceased respondent received any home care in the last 12 months before death. In Waves 2 through 5, the proxy is asked whether the respondent received home care or home help due to disability. Starting in Wave 6, the proxy is asked separately whether the respondent received help with personal care due to disability and whether they received help with domestic tasks due to disability. **raxhomcar1y** is coded as 0 if the proxy reports that the respondent did not receive home care in the last 12 months before death. **raxhomcar1y** is coded as 1 if the proxy reports that the respondent received home care in the last 12 months before death. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Cross Wave Differences in SHARE EOL

No differences known.

Differences with the Harmonized HRS End of Life

The HRS asks whether the respondent had any medically-trained person come to their home to help them since the previous wave or in the last two years, whereas SHARE asks whether the respondent received home care or home help due to disability in Waves 2 through 5 and whether the respondent received help with personal care due to disability or help with domestic tasks due to disability separately starting in Wave 6 in the last 12 months of life. As such, the Harmonized HRS End of Life includes **raxhomcar**, while the Gateway Harmonized SHARE End of Life includes **raxhomcar1y**.

Categorical Variable Frequencies

	raxhomcar1y
0.no	11,592
1.yes	9,099
.d:dk	240
.m:missing	125
.r:refuse	42
Total	21,098

Original SHARE Variables Used

Wave 2 End of Life:	
xto18_8	Type of medical care in last 12 months: home care or help due to disability
Wave 3 End of Life:	
sl_xto18_8	Had medical care in last 12 months: home care or home help due to disability
Wave 4 End of Life:	
xto18_8	Type of medical care in last 12 months: home care or help due to disability
Wave 5 End of Life:	
xto18_8	Type of medical care in last 12 months: home care or help due to disability
Wave 6 End of Life:	
xto18_8	Type of medical care last 12 months: help with personal care due to disability
xto18_9	Type of medical care last 12 months: help with domestic tasks due to disability
Wave 7 End of Life:	

xto18_8	Type of medical care last 12 months: help with personal care due to disability
xto18_9	Type of medical care last 12 months: help with domestic tasks due to disability
Wave 8 End of Life:	
xto18_8	Type of medical care last 12 months: help with personal care due to disability
xto18_9	Type of medical care last 12 months: help with domestic tasks due to disability
Wave 9 End of Life:	
xto18_8	Type of medical care last 12 months: help with personal care due to disability
xto18_9	Type of medical care last 12 months: help with domestic tasks due to disability
Wave 9 COVID End of Life:	
xto18_8	Type of medical care last 12 months: help with personal care due to disability
xto18_9	Type of medical care last 12 months: help with domestic tasks due to disability

Out-Of-Pocket Medical Costs: Total			
Variable	Waves	Label	Type
raxoophos1y	All	raxoophos1y: r OOP cost: hospital	Cont
raxoophosf1y	All	raxoophosf1y: r OOP cost flag: hospital	Categ
raxoophnh1y	All	raxoophnh1y: r OOP cost: nursing home	Cont
raxoophnhf1y	All	raxoophnhf1y: r OOP cost flag: nursing home	Categ
raxoophpc1y	All	raxoophpc1y: r OOP cost: hospice	Cont
raxoophpcf1y	All	raxoophpcf1y: r OOP cost flag: hospice	Categ
raxoopdoc1y	All	raxoopdoc1y: r OOP cost: doctor visits	Cont
raxoopdocf1y	All	raxoopdocf1y: r OOP cost flag: doctor visits	Categ
raxoopdrug1y	All	raxoopdrug1y: r OOP cost: drugs	Cont
raxoopdrugf1y	All	raxoopdrugf1y: r OOP cost flag: drugs	Categ
raxoophmcr1y	All	raxoophmcr1y: r OOP cost: home care	Cont
raxoophmcrf1y	All	raxoophmcrf1y: r OOP cost flag: home care	Categ
raxoopome1y	All	raxoopome1y: r OOP cost: other med exp	Cont
raxoopomef1y	All	raxoopomef1y: r OOP cost flag: other med exp	Categ
raxoopmd1y	All	raxoopmd1y: r OOP cost: total major medical expenses	Cont
raxoopmdf1y	All	raxoopmdf1y: r OOP cost flag: total major medical expenses	Categ

How Constructed

raxoophos1y, **raxoophnh1y**, **raxoophpc1y**, **raxoopdoc1y**, **raxoopdrug1y**, **raxoopome1y**, **raxoophmcr1y**, and **raxoopmd1y** have been adjusted to 2010 Euros based on the consumer price index for the year and country of death.

raxoophos1y, **raxoophnh1y**, **raxoophpc1y**, **raxoopdoc1y**, **raxoopdrug1y**, **raxoopome1y**, and **raxoophmcr1y** are the out-of-pocket costs for specific major medical expenses and include imputed values. The proxy is asked about how much the respondent paid out-of-pocket for each major medical expenditure in the last year of life. In Wave 3 and onward, if the proxy answers don't know or refuses to answer, a series of unfolding bracket questions is asked to obtain a minimum and maximum bracket value of the out-of-pocket expenses. If the respondent did not use one of the specified services, then the out-of-pocket cost for that item is assumed to be 0. For **raxoophos1y**, **raxoophnh1y**, and **raxoophpc1y**, if the proxy reports that the respondent did not have any stays at a hospital, hospice, or nursing home, then these questions are not asked and are assumed to be 0. All out-of-pocket expenses are elicited in Euros in all SHARE countries except in Denmark, Sweden, Switzerland, Poland, Czech Republic, and Israel where local currencies are used and which have been converted to Euros by the SHARE team. However, all reported or imputed values in these variables have been converted and adjusted to 2010 Euros based on the consumer price index for the year and country of death. In Waves 6 and 7, if the respondent had 1 stay at a hospital, hospice, or nursing home and the cost for receiving care at these facilities is not asked, then **raxoophos1y**, **raxoophnh1y**, and **raxoophpc1y** are assigned special missing (.s). **raxoophos1y**, **raxoophnh1y**, **raxoophpc1y**, **raxoopdoc1y**, **raxoopdrug1y**, **raxoopome1y**, and **raxoophmcr1y** are assigned special missing (.m) if the values are not able to be imputed due to a lack of reference observations in the donor pool or missing covariates necessary for the imputation procedure.

raxoophos1y is the out-of-pocket costs for hospital stays and includes imputed values. **raxoophnh1y** is the out-of-pocket costs for nursing home stays and includes imputed values. **raxoophpc1y** is the out-of-pocket costs for hospice and includes imputed values. **raxoopdoc1y** is the out-of-pocket costs for general practitioners and specialist physicians and includes imputed values. **raxoopdrug1y** is the out-of-pocket costs for medication and includes imputed values. **raxoopome1y** is the out-of-pocket costs for aids and appliances and includes imputed values. **raxoophmcr1y** is the out-of-pocket costs

for home care or help due to disability in Waves 2 through 5, and for help with personal care due to disability and help with domestic tasks due to disability starting in Wave 6 and includes imputed values.

In Wave 2, the proxy is not asked any additional questions if the proxy does not give an exact amount for each out-of-pocket expenditure. For this wave, don't know and refused answers are imputed with no bracket information. In Wave 3 and onward, if the proxy does not give an exact amount for each out-of-pocket expenditure, the proxy is asked a set of bracket questions and answers if the true value was less than, about equal to, or more than the threshold value. In Wave 3, the threshold values vary by country, even within Euro-using countries, and vary by expense type. In Wave 4, the threshold values vary by country, but all Euro-using countries use the same threshold values, and the values vary by expense type. In Waves 5-7, the threshold values vary by country, even within Euro-using countries, but the values are constant across all expense types. Starting in Wave 8, the threshold values vary by country and expense type. This information obtained from the unfolding bracket sequence is used in the imputation of **raxoophos1y**, **raxoophnh1y**, **raxoophpc1y**, **raxoopdoc1y**, **raxoopdrug1y**, **raxoopome1y**, and **raxoophmcr1y** to address item-level missingness in these variables.

raxoophosf1y, **raxoophnhf1y**, **raxoophpcf1y**, **raxoopdocf1y**, **raxoopdrugf1y**, **raxoopomef1y**, and **raxoophmcrf1y** are flag variables indicating the level of imputation for the specified out-of-pocket expense. A code of 1 indicates the proxy reported a continuous value and no imputation was necessary. A code of 2 indicates that the component was imputed based on a closed bracket. A code of 3 indicates that the component was imputed based on an open bracket. A code of 5 indicates that the component was imputed without any bracket information. A code of 6 indicates the proxy reported the respondent not having the expense and the value is 0. A code of 7 indicates that the proxy was not sure whether the respondent had this expense. A code of -1 indicates that the expense could not be imputed due to a lack of reference observations in the donor pool. A code of -2 indicates that the expense could not be imputed due to one or more missing values in the covariates used in the imputation procedure. In Waves 6 and 7, a special missing code of (.s) for **raxoophosf1y**, **raxoophnhf1y**, or **raxoophpcf1y** indicates that the proxy reported a single hospital, hospice, or nursing home stay for the respondent, and these questions were not asked nor were values imputed.

raxoopmd1y is the total out-of-pocket costs for major medical expenses in the last year of life and includes imputed values. The proxy is first asked whether the respondent received care from a list of sources, and for each one the proxy reports the respondent used, the proxy is asked about how much the respondent paid out-of-pocket for each major medical expenditure in the last year before death. In Wave 3 and onward, if the proxy answers don't know or refuses to answer, a series of unfolding bracket questions is asked to obtain a minimum and maximum bracket value of the out-of-pocket expenses. **raxoopmd1y** is the sum of the reported or imputed out-of-pocket expenses for hospital stays, care in a nursing home, hospice stays, care from a general practitioner, care from specialist physicians, medication, aids and appliances, and help with personal care and domestic tasks due to disability (**raxoopmd1y** = **raxoophos1y** + **raxoophnh1y** + **raxoophpc1y** + **raxoopdoc1y** + **raxoopdrug1y** + **raxoopome1y** + **raxoophmcr1y**). Please note that in Waves 2 through 5, the proxy is asked about the cost of home care or help due to disability in a single question, and starting in Wave 6, the proxy is asked about the cost of help with personal care due to disability and help with domestic tasks due to disability in separate questions. In Waves 6 and 7, if the respondent had 1 stay at a hospital, hospice, or nursing home and the cost of receiving care at these facilities is not asked, then **raxoopmd1y** is assigned special missing (.s). **raxoopmd1y** is assigned special missing (.m) if at least one of the comprising values could be imputed due to a lack of reference observations in the donor pool or missing covariates necessary for the imputation procedure.

raxoopmdf1y is a flag variable indicating the highest level of imputation of the components of **raxoopmd1y**. A code of 1 indicates the proxy reported continuous values for all components and no imputation was necessary. A code of 2 indicates that at least one component was imputed based on a closed bracket. A code of 3 indicates that at least one component was imputed based on an open bracket. A code of 5 indicates that at least one component was imputed without any bracket information. A code of 6 indicates the proxy reported the respondent not having any of the expenses and the value is 0. A code of 7 indicates that the proxy was unsure whether the respondent had at least one of the expenses. A code of -1 indicates that none of the expenses could be imputed due to a lack of reference observations in the donor pool. A code of -2 indicates that none of the expenses could be imputed due to one or more missing values in the covariates used in the imputation procedure. In Waves 6 and 7, a special missing code of (.s) indicates that the proxy reported a single hospital, hospice, or nursing home stay for the respondent, and questions about hospital, hospice, and nursing home stays were not asked nor were values imputed.

Cross Wave Differences in SHARE EOL

In Waves 2 through 5 and starting in Wave 8, if the proxy reports that the respondent did not have any stays at a hospital, hospice, or nursing home, then the proxy is not asked whether the respondent stayed in a hospital, hospice, or nursing home or how much was spent out-of-pocket for these services. In Waves 6 and 7, if the proxy reports that the respondent had 0 or 1 stays at a hospital, hospice, or nursing home then the proxy is not asked whether the respondent stayed in a hospital, hospice, or nursing home or how much was spent out-of-pocket for these services. In Waves 6 and 7, if the respondent had 1 stay at a hospital, hospice, or nursing home, then the out-of-pocket costs for these expenses are not imputed and are assigned special missing (.s) in the cost and in the flag variables.

In Waves 2 through 5, the proxy is asked about the cost of home care or help due to disability in a single question, and starting in Wave 6, the proxy is asked about the cost of help with personal care due to disability and help with domestic tasks due to disability in separate questions.

In Wave 2, the proxy is not asked any additional questions if the proxy does not give an exact amount for each out-of-pocket expenditure. For this wave, don't know and refused answers are imputed with no bracket information. In Wave 3, the threshold values vary by country, even within Euro-using countries, and vary by expense type. In Wave 4, the threshold values vary by country, but all Euro-using countries use the same threshold values, and the values vary by expense type. In Waves 5 through 7, the threshold values vary by country, even within Euro-using countries, and the values are constant across all expense types. Starting in Wave 8, the threshold values vary by country, even within Euro-using countries, and vary by expense type.

Please note that the threshold values in Wave 4 are significantly higher for each country than the values used in any other wave.

Differences with the Harmonized HRS End of Life

All out-of-pocket costs in the Harmonized HRS End of Life are denominated in 2010 U.S. dollars and are for the last 2 years of life or since the last interview, whereas all out-of-pocket costs in the Gateway Harmonized SHARE End of Life are denominated in 2010 Euros and are for the last year of life. Therefore, conversion into a common currency and an adjustment for the period of time is necessary before comparison of these data. Additionally, if there were insufficient observations for the donor pool by country in the SHARE, if there were missing variables required for the imputation procedure, or if the respondent had a single hospital, hospice or nursing home stay in Waves 6 and 7, then expenses could not be imputed and are assigned special missing (.m) and (.s), respectively, in the Gateway Harmonized SHARE End of Life, whereas all out-of-pocket costs were either reported or imputed in the Harmonized HRS End of Life.

raxoopdoc in the Harmonized HRS End of Life includes the costs of doctor visits from a single question, whereas **raxoopdoc1y** in the Gateway Harmonized SHARE End of Life includes the cost of general practitioners and the cost of specialist physicians, asked in 2 separate questions.

raxoopmcr in the Harmonized HRS End of Life includes the costs of in-home medical care from a single question, whereas **raxoopmcr1y** in the Gateway Harmonized SHARE End of Life includes the cost of home care or help due to disability from a single question in Wave 2 through 5, or it includes the cost of help with personal care due to disability and help with domestic tasks due to disability from 2 separate questions starting in Wave 6.

raxoopome in the Harmonized HRS End of Life includes the costs of other medical expenses not covered by insurance, including medications, special food, equipment, visits by health professionals, or other costs. In the Gateway Harmonized SHARE End of Life, **raxoopome1y** includes the costs of aids and appliances.

raxoopmd in the Harmonized HRS End of Life incorporates the reported and imputed costs of special facilities and services (**raxoopspec**), which has no corollary in the Gateway Harmonized SHARE End of Life.

Continuous Variable Distributions

Variable	N	Mean	Std.Dev	Minimum	Maximum	Total missing
raxoopphos1y	19,176	166.99	3,430.37	0.00	244,379.27	1,922

Section C. Health Care Utilization and Insurance

raxoopnh1y	19,182	572.76	3,486.27	0.00	140,432.77	1,916
raxoophpc1y	19,181	33.82	458.31	0.00	19,550.34	1,917
raxoopdoc1y	21,010	405.89	8,651.42	0.00	1,011,235.94	88
raxoopdrug1y	21,020	315.07	2,561.02	0.00	234,126.23	78
raxoophmcr1y	21,005	526.02	3,225.23	0.00	230,202.58	93
raxoopome1y	21,024	85.13	871.25	0.00	103,528.32	74
raxoopmd1y	19,102	2,099.25	11,987.56	0.00	1,011,349.31	1,996

Categorical Variable Frequencies

	raxoopphosf1y
-2.not imputed, missing cov	70
-1.not imputed, missing nei	8
1.continuous value	9,144
2.closed bracket	451
3.open bracket	271
5.no bracket info	919
6.no expense	7,183
7.dk had expense	1,208
.s:single stay	1,844
Total	21,098

	raxoopnhf1y
-2.not imputed, missing cov	66
-1.not imputed, missing nei	6
1.continuous value	2,000
2.closed bracket	180
3.open bracket	132
5.no bracket info	329
6.no expense	15,347
7.dk had expense	1,194
.s:single stay	1,844
Total	21,098

	raxoophpcf1y
-2.not imputed, missing cov	65
-1.not imputed, missing nei	8
1.continuous value	983
2.closed bracket	67
3.open bracket	49
5.no bracket info	185
6.no expense	16,646
7.dk had expense	1,251
.s:single stay	1,844
Total	21,098

	raxoopdocf1y
-2.not imputed, missing cov	68
-1.not imputed, missing nei	20

1.continuous value	14,736
2.closed bracket	1,037
3.open bracket	540
5.no bracket info	1,774
6.no expense	2,026
7.dk had expense	897
Total	21,098
raxoopdrugf1y	
-2.not imputed, missing cov	71
-1.not imputed, missing nei	7
1.continuous value	13,651
2.closed bracket	1,830
3.open bracket	853
5.no bracket info	2,601
6.no expense	1,607
7.dk had expense	478
Total	21,098
raxoophmcrf1y	
-2.not imputed, missing cov	64
-1.not imputed, missing nei	29
1.continuous value	7,141
2.closed bracket	355
3.open bracket	317
5.no bracket info	1,159
6.no expense	11,492
7.dk had expense	541
Total	21,098
raxoopomef1y	
-2.not imputed, missing cov	65
-1.not imputed, missing nei	9
1.continuous value	6,615
2.closed bracket	479
3.open bracket	229
5.no bracket info	969
6.no expense	12,224
7.dk had expense	508
Total	21,098
raxoopmdf1y	
-2.not imputed, missing cov	75
-1.not imputed, missing nei	77
1.continuous value	11,221
2.closed bracket	1,807
3.open bracket	964
5.no bracket info	2,672
6.no expense	573
7.dk had expense	1,865
.s:single stay	1,844
Total	21,098

Original SHARE Variables Used

Wave 2 End of Life:

xt018_1	Type of medical care in last 12 months: care from a general practitioner
xt018_2	Type of medical care in last 12 months: care from specialist physicians
xt018_3	Type of medical care in last 12 months: hospital stays
xt018_4	Type of medical care in last 12 months: care in a nursing home
xt018_5	Type of medical care in last 12 months: hospice stays
xt018_6	Type of medical care in last 12 months: medication
xt018_7	Type of medical care in last 12 months: aids and appliances
xt018_8	Type of medical care in last 12 months: home care or help due to disability
xt019e_1	Costs: care from a general practitioner
xt019e_2	Costs: care from specialist physicians
xt019e_3	Costs: hospital stays
xt019e_4	Costs: care in a nursing home
xt019e_5	Costs: hospice stays
xt019e_6	Costs: medication
xt019e_7	Costs: aids and appliances
xt019e_8	Costs: home care or help due to disability

Wave 3 End of Life:

sl_xt018_1	Had medical care in last 12 months: care from a general practitioner
sl_xt018_2	Had medical care in last 12 months: care from specialist physicians
sl_xt018_3	Had medical care in last 12 months: hospital stays
sl_xt018_4	Had medical care in last 12 months: care in a nursing home
sl_xt018_5	Had medical care in last 12 months: hospice stays
sl_xt018_6	Had medical care in last 12 months: medication
sl_xt018_7	Had medical care in last 12 months: aids and appliances
sl_xt018_8	Had medical care in last 12 months: home care or home help due to disability
sl_xt019e_1	Costs medical care in last 12 months: care from a general practitioner
sl_xt019e_2	Costs medical care in last 12 months: care from specialist physicians
sl_xt019e_3	Costs medical care in last 12 months: hospital stays
sl_xt019e_4	Costs medical care in last 12 months: care in a nursing home
sl_xt019e_5	Costs medical care in last 12 months: hospice stays
sl_xt019e_6	Costs medical care in last 12 months: medication
sl_xt019e_7	Costs medical care in last 12 months: aids and appliances
sl_xt019e_8	Costs medical care in last 12 months: home care or home help due to disability
sl_xt019ub_1	Costs medical care in last 12 months: care from a general practitioner ub
sl_xt019ub_2	Costs medical care in last 12 months: care from specialist physicians ub
sl_xt019ub_3	Costs medical care in last 12 months: hospital stays ub
sl_xt019ub_4	Costs medical care in last 12 months: care in a nursing home ub
sl_xt019ub_5	Costs medical care in last 12 months: hospice stays ub
sl_xt019ub_6	Costs medical care in last 12 months: medication ub
sl_xt019ub_7	Costs medical care in last 12 months: aids and appliances ub
sl_xt019ub_8	Costs medical care in last 12 months: home care or home help due to disability u
sl_xt019v1_1	Bracket value 1: care from a general practitioner
sl_xt019v1_2	Bracket value 1: care from specialist physicians
sl_xt019v1_3	Bracket value 1: hospital stays
sl_xt019v1_4	Bracket value 1: care in a nursing home
sl_xt019v1_5	Bracket value 1: hospice stays
sl_xt019v1_6	Bracket value 1: medication
sl_xt019v1_7	Bracket value 1: aids and appliances
sl_xt019v1_8	Bracket value 1: home care or home help due to disability
sl_xt019v2_1	Bracket value 2: care from a general practitioner

sl_xto19v2_2	Bracket value 2: care from specialist physicians
sl_xto19v2_3	Bracket value 2: hospital stays
sl_xto19v2_4	Bracket value 2: care in a nursing home
sl_xto19v2_5	Bracket value 2: hospice stays
sl_xto19v2_6	Bracket value 2: medication
sl_xto19v2_7	Bracket value 2: aids and appliances
sl_xto19v2_8	Bracket value 2: home care or home help due to disability
sl_xto19v3_1	Bracket value 3: care from a general practitioner
sl_xto19v3_2	Bracket value 3: care from specialist physicians
sl_xto19v3_3	Bracket value 3: hospital stays
sl_xto19v3_4	Bracket value 3: care in a nursing home
sl_xto19v3_5	Bracket value 3: hospice stays
sl_xto19v3_6	Bracket value 3: medication
sl_xto19v3_7	Bracket value 3: aids and appliances
sl_xto19v3_8	Bracket value 3: home care or home help due to disability
country	Country identifier
Wave 4 End of Life:	
xt018_1	Type of medical care in last 12 months: care from a general practitioner
xt018_2	Type of medical care in last 12 months: care from specialist physicians
xt018_3	Type of medical care in last 12 months: hospital stays
xt018_4	Type of medical care in last 12 months: care in a nursing home
xt018_5	Type of medical care in last 12 months: hospice stays
xt018_6	Type of medical care in last 12 months: medication
xt018_7	Type of medical care in last 12 months: aids and appliances
xt018_8	Type of medical care in last 12 months: home care or help due to disability
xt019e_1	Costs: care from a general practitioner
xt019e_2	Costs: care from specialist physicians
xt019e_3	Costs: hospital stays
xt019e_4	Costs: care in a nursing home
xt019e_5	Costs: hospice stays
xt019e_6	Costs: medication
xt019e_7	Costs: aids and appliances
xt019e_8	Costs: home care or help due to disability
xt019ub_1	Costs: care from a general practitioner ub
xt019ub_2	Costs: care from specialist physicians ub
xt019ub_3	Costs: hospital stays ub
xt019ub_4	Costs: care in a nursing home ub
xt019ub_5	Costs: hospice stays ub
xt019ub_6	Costs: medication ub
xt019ub_7	Costs: aids and appliances ub
xt019ub_8	Costs: home care or help due to disability ub
xt019v1_1	Bracket value 1: care from a general practitioner
xt019v1_2	Bracket value 1: care from specialist physicians
xt019v1_3	Bracket value 1: hospital stays
xt019v1_4	Bracket value 1: care in a nursing home
xt019v1_5	Bracket value 1: hospice stays
xt019v1_6	Bracket value 1: medication
xt019v1_7	Bracket value 1: aids and appliances
xt019v1_8	Bracket value 1: home care or help due to disability
xt019v2_1	Bracket value 2: care from a general practitioner
xt019v2_2	Bracket value 2: care from specialist physicians
xt019v2_3	Bracket value 2: hospital stays
xt019v2_4	Bracket value 2: care in a nursing home
xt019v2_5	Bracket value 2: hospice stays

xt019v2_6	Bracket value 2: medication
xt019v2_7	Bracket value 2: aids and appliances
xt019v2_8	Bracket value 2: home care or help due to disability
xt019v3_1	Bracket value 3: care from a general practitioner
xt019v3_2	Bracket value 3: care from specialist physicians
xt019v3_3	Bracket value 3: hospital stays
xt019v3_4	Bracket value 3: care in a nursing home
xt019v3_5	Bracket value 3: hospice stays
xt019v3_6	Bracket value 3: medication
xt019v3_7	Bracket value 3: aids and appliances
xt019v3_8	Bracket value 3: home care or help due to disability
country	Country identifier
Wave 5 End of Life:	
xt018_1	Type of medical care in last 12 months: care from a general practitioner
xt018_2	Type of medical care in last 12 months: care from specialist physicians
xt018_3	Type of medical care in last 12 months: hospital stays
xt018_4	Type of medical care in last 12 months: care in a nursing home
xt018_5	Type of medical care in last 12 months: hospice stays
xt018_6	Type of medical care in last 12 months: medication
xt018_7	Type of medical care in last 12 months: aids and appliances
xt018_8	Type of medical care in last 12 months: home care or help due to disability
xt019e_1	Costs: care from a general practitioner
xt019e_2	Costs: care from specialist physicians
xt019e_3	Costs: hospital stays
xt019e_4	Costs: care in a nursing home
xt019e_5	Costs: hospice stays
xt019e_6	Costs: medication
xt019e_7	Costs: aids and appliances
xt019e_8	Costs: home care or help due to disability
xt019ub_1	Costs: care from a general practitioner ub
xt019ub_2	Costs: care from specialist physicians ub
xt019ub_3	Costs: hospital stays ub
xt019ub_4	Costs: care in a nursing home ub
xt019ub_5	Costs: hospice stays ub
xt019ub_6	Costs: medication ub
xt019ub_7	Costs: aids and appliances ub
xt019ub_8	Costs: home care or help due to disability ub
xt019v1	Bracket value 1
xt019v2	Bracket value 2
xt019v3	Bracket value 3
country	Country identifier
Wave 6 End of Life:	
xt018_1	Type of medical care in last 12 months: care from a general practitioner
xt018_2	Type of medical care in last 12 months: care from specialist physicians
xt018_3	Type of medical care in last 12 months: hospital stays
xt018_4	Type of medical care in last 12 months: care in a nursing home
xt018_5	Type of medical care in last 12 months: hospice stays
xt018_6	Type of medical care in last 12 months: medication
xt018_7	Type of medical care in last 12 months: aids and appliances
xt018_8	Type of medical care last 12 months: help with personal care due to disability
xt018_9	Type of medical care last 12 months: help with domestic tasks due to disability
xt119e_1	Costs: care from a general practitioner
xt119e_2	Costs: care from specialist physicians

xt119e_3	Costs: hospital stays
xt119e_4	Costs: care in a nursing home
xt119e_5	Costs: hospice stays
xt119e_6	Costs: medication
xt119e_7	Costs: aids and appliances
xt119e_8	Costs: help with personal care due to disability
xt119e_9	Costs: help with domestic tasks due to disability
xt119ub_1	Costs: care from a general practitioner ub
xt119ub_2	Costs: care from specialist physicians ub
xt119ub_3	Costs: hospital stays ub
xt119ub_4	Costs: care in a nursing home ub
xt119ub_5	Costs: hospice stays ub
xt119ub_6	Costs: medication ub
xt119ub_7	Costs: aids and appliances ub
xt119ub_8	Costs: help with personal care due to disability ub
xt119ub_9	Costs: help with domestic tasks due to disability ub
xt119v1_1	Bracket value 1
xt119v1_2	Bracket value 1
xt119v1_3	Bracket value 1
xt119v1_4	Bracket value 1
xt119v1_5	Bracket value 1
xt119v1_6	Bracket value 1
xt119v1_7	Bracket value 1
xt119v1_8	Bracket value 1
xt119v1_9	Bracket value 1
xt119v2_1	Bracket value 2
xt119v2_2	Bracket value 2
xt119v2_3	Bracket value 2
xt119v2_4	Bracket value 2
xt119v2_5	Bracket value 2
xt119v2_6	Bracket value 2
xt119v2_7	Bracket value 2
xt119v2_8	Bracket value 2
xt119v2_9	Bracket value 2
xt119v3_1	Bracket value 3
xt119v3_2	Bracket value 3
xt119v3_3	Bracket value 3
xt119v3_4	Bracket value 3
xt119v3_5	Bracket value 3
xt119v3_6	Bracket value 3
xt119v3_7	Bracket value 3
xt119v3_8	Bracket value 3
xt119v3_9	Bracket value 3
country	Country identifier

Wave 7 End of Life:

xt018_1	Type of medical care in last 12 months: care from a general practitioner
xt018_2	Type of medical care in last 12 months: care from specialist physicians
xt018_3	Type of medical care in last 12 months: hospital stays
xt018_4	Type of medical care in last 12 months: care in a nursing home
xt018_5	Type of medical care in last 12 months: hospice stays
xt018_6	Type of medical care in last 12 months: medication
xt018_7	Type of medical care in last 12 months: aids and appliances
xt018_8	Type of medical care last 12 months: help with personal care due to disability
xt018_9	Type of medical care last 12 months: help with domestic tasks due to disability

xt119e_1	Costs: care from a general practitioner
xt119e_2	Costs: care from specialist physicians
xt119e_3	Costs: hospital stays
xt119e_4	Costs: care in a nursing home
xt119e_5	Costs: hospice stays
xt119e_6	Costs: medication
xt119e_7	Costs: aids and appliances
xt119e_8	Costs: help with personal care due to disability
xt119e_9	Costs: help with domestic tasks due to disability
xt119ub_1	Costs: care from a general practitioner ub
xt119ub_2	Costs: care from specialist physicians ub
xt119ub_3	Costs: hospital stays ub
xt119ub_4	Costs: care in a nursing home ub
xt119ub_5	Costs: hospice stays ub
xt119ub_6	Costs: medication ub
xt119ub_7	Costs: aids and appliances ub
xt119ub_8	Costs: help with personal care due to disability ub
xt119ub_9	Costs: help with domestic tasks due to disability ub
xt119v1_1	Bracket value 1
xt119v1_2	Bracket value 1
xt119v1_3	Bracket value 1
xt119v1_4	Bracket value 1
xt119v1_5	Bracket value 1
xt119v1_6	Bracket value 1
xt119v1_7	Bracket value 1
xt119v1_8	Bracket value 1
xt119v1_9	Bracket value 1
xt119v2_1	Bracket value 2
xt119v2_2	Bracket value 2
xt119v2_3	Bracket value 2
xt119v2_4	Bracket value 2
xt119v2_5	Bracket value 2
xt119v2_6	Bracket value 2
xt119v2_7	Bracket value 2
xt119v2_8	Bracket value 2
xt119v2_9	Bracket value 2
xt119v3_1	Bracket value 3
xt119v3_2	Bracket value 3
xt119v3_3	Bracket value 3
xt119v3_4	Bracket value 3
xt119v3_5	Bracket value 3
xt119v3_6	Bracket value 3
xt119v3_7	Bracket value 3
xt119v3_8	Bracket value 3
xt119v3_9	Bracket value 3
country	Country identifier

Wave 8 End of Life:

xt018_1	Type of medical care in last 12 months: care from a general practitioner
xt018_2	Type of medical care in last 12 months: care from specialist physicians
xt018_3	Type of medical care in last 12 months: hospital stays
xt018_4	Type of medical care in last 12 months: care in a nursing home
xt018_5	Type of medical care in last 12 months: hospice stays
xt018_6	Type of medical care in last 12 months: medication
xt018_7	Type of medical care in last 12 months: aids and appliances

xt018_8	Type of medical care last 12 months: help with personal care due to disability
xt018_9	Type of medical care last 12 months: help with domestic tasks due to disability
xt119e_1	Costs: care from a general practitioner
xt119e_2	Costs: care from specialist physicians
xt119e_3	Costs: hospital stays
xt119e_4	Costs: care in a nursing home
xt119e_5	Costs: hospice stays
xt119e_6	Costs: medication
xt119e_7	Costs: aids and appliances
xt119e_8	Costs: help with personal care due to disability
xt119e_9	Costs: help with domestic tasks due to disability
xt119ub_1	Costs: care from a general practitioner ub
xt119ub_2	Costs: care from specialist physicians ub
xt119ub_3	Costs: hospital stays ub
xt119ub_4	Costs: care in a nursing home ub
xt119ub_5	Costs: hospice stays ub
xt119ub_6	Costs: medication ub
xt119ub_7	Costs: aids and appliances ub
xt119ub_8	Costs: help with personal care due to disability ub
xt119ub_9	Costs: help with domestic tasks due to disability ub
xt119v1_1	Bracket value 1
xt119v1_2	Bracket value 1
xt119v1_3	Bracket value 1
xt119v1_4	Bracket value 1
xt119v1_5	Bracket value 1
xt119v1_6	Bracket value 1
xt119v1_7	Bracket value 1
xt119v1_8	Bracket value 1
xt119v1_9	Bracket value 1
xt119v2_1	Bracket value 2
xt119v2_2	Bracket value 2
xt119v2_3	Bracket value 2
xt119v2_4	Bracket value 2
xt119v2_5	Bracket value 2
xt119v2_6	Bracket value 2
xt119v2_7	Bracket value 2
xt119v2_8	Bracket value 2
xt119v2_9	Bracket value 2
xt119v3_1	Bracket value 3
xt119v3_2	Bracket value 3
xt119v3_3	Bracket value 3
xt119v3_4	Bracket value 3
xt119v3_5	Bracket value 3
xt119v3_6	Bracket value 3
xt119v3_7	Bracket value 3
xt119v3_8	Bracket value 3
xt119v3_9	Bracket value 3
country	Country identifier
Wave 9 End of Life:	
xt018_1	Type of medical care in last 12 months: care from a general practitioner
xt018_2	Type of medical care in last 12 months: care from specialist physicians
xt018_3	Type of medical care in last 12 months: hospital stays
xt018_4	Type of medical care in last 12 months: care in a nursing home
xt018_5	Type of medical care in last 12 months: hospice stays

xt018_6	Type of medical care in last 12 months: medication
xt018_7	Type of medical care in last 12 months: aids and appliances
xt018_8	Type of medical care last 12 months: help with personal care due to disability
xt018_9	Type of medical care last 12 months: help with domestic tasks due to disability
xt119e_1	Costs: care from a general practitioner
xt119e_2	Costs: care from specialist physicians
xt119e_3	Costs: hospital stays
xt119e_4	Costs: care in a nursing home
xt119e_5	Costs: hospice stays
xt119e_6	Costs: medication
xt119e_7	Costs: aids and appliances
xt119e_8	Costs: help with personal care due to disability
xt119e_9	Costs: help with domestic tasks due to disability
xt119ub_1	Costs: care from a general practitioner ub
xt119ub_2	Costs: care from specialist physicians ub
xt119ub_3	Costs: hospital stays ub
xt119ub_4	Costs: care in a nursing home ub
xt119ub_5	Costs: hospice stays ub
xt119ub_6	Costs: medication ub
xt119ub_7	Costs: aids and appliances ub
xt119ub_8	Costs: help with personal care due to disability ub
xt119ub_9	Costs: help with domestic tasks due to disability ub
xt119v1_1	Bracket value 1
xt119v1_2	Bracket value 1
xt119v1_3	Bracket value 1
xt119v1_4	Bracket value 1
xt119v1_5	Bracket value 1
xt119v1_6	Bracket value 1
xt119v1_7	Bracket value 1
xt119v1_8	Bracket value 1
xt119v1_9	Bracket value 1
xt119v2_1	Bracket value 2
xt119v2_2	Bracket value 2
xt119v2_3	Bracket value 2
xt119v2_4	Bracket value 2
xt119v2_5	Bracket value 2
xt119v2_6	Bracket value 2
xt119v2_7	Bracket value 2
xt119v2_8	Bracket value 2
xt119v2_9	Bracket value 2
xt119v3_1	Bracket value 3
xt119v3_2	Bracket value 3
xt119v3_3	Bracket value 3
xt119v3_4	Bracket value 3
xt119v3_5	Bracket value 3
xt119v3_6	Bracket value 3
xt119v3_7	Bracket value 3
xt119v3_8	Bracket value 3
xt119v3_9	Bracket value 3
country	Country identifier

Wave 9 COVID End of Life:

xt018_1	Type of medical care in last 12 months: care from a general practitioner
xt018_2	Type of medical care in last 12 months: care from specialist physicians
xt018_3	Type of medical care in last 12 months: hospital stays

xt018_4	Type of medical care in last 12 months: care in a nursing home
xt018_5	Type of medical care in last 12 months: hospice stays
xt018_6	Type of medical care in last 12 months: medication
xt018_7	Type of medical care in last 12 months: aids and appliances
xt018_8	Type of medical care last 12 months: help with personal care due to disability
xt018_9	Type of medical care last 12 months: help with domestic tasks due to disability
xt119e_1	Costs: care from a general practitioner
xt119e_2	Costs: care from specialist physicians
xt119e_3	Costs: hospital stays
xt119e_4	Costs: care in a nursing home
xt119e_5	Costs: hospice stays
xt119e_6	Costs: medication
xt119e_7	Costs: aids and appliances
xt119e_8	Costs: help with personal care due to disability
xt119e_9	Costs: help with domestic tasks due to disability
xt119ub_1	Costs: care from a general practitioner ub
xt119ub_2	Costs: care from specialist physicians ub
xt119ub_3	Costs: hospital stays ub
xt119ub_4	Costs: care in a nursing home ub
xt119ub_5	Costs: hospice stays ub
xt119ub_6	Costs: medication ub
xt119ub_7	Costs: aids and appliances ub
xt119ub_8	Costs: help with personal care due to disability ub
xt119ub_9	Costs: help with domestic tasks due to disability ub
xt119v1_1	Bracket value 1
xt119v1_2	Bracket value 1
xt119v1_3	Bracket value 1
xt119v1_4	Bracket value 1
xt119v1_5	Bracket value 1
xt119v1_6	Bracket value 1
xt119v1_7	Bracket value 1
xt119v1_8	Bracket value 1
xt119v1_9	Bracket value 1
xt119v2_1	Bracket value 2
xt119v2_2	Bracket value 2
xt119v2_3	Bracket value 2
xt119v2_4	Bracket value 2
xt119v2_5	Bracket value 2
xt119v2_6	Bracket value 2
xt119v2_7	Bracket value 2
xt119v2_8	Bracket value 2
xt119v2_9	Bracket value 2
xt119v3_1	Bracket value 3
xt119v3_2	Bracket value 3
xt119v3_3	Bracket value 3
xt119v3_4	Bracket value 3
xt119v3_5	Bracket value 3
xt119v3_6	Bracket value 3
xt119v3_7	Bracket value 3
xt119v3_8	Bracket value 3
xt119v3_9	Bracket value 3
country	Country identifier

D. Cognition

Whether R Had Difficulty Remembering in Last Year of Life			
Variable	Waves	Label	Type
raxdfrloc	All	raxdfrloc: r difficulty remembering location in last year	Categ
raxdfryr	All	raxdfryr: r difficulty remembering year in last year	Categ
raxdfrfam	All	raxdfrfam: r difficulty recognizing family/friends in last year	Categ

How Constructed

raxdfrloc, **raxdfryr**, and **raxdfrfam** indicate whether the respondent had difficulty remembering certain things in their last year of life that lasted at least three months because of physical, mental, emotional, or memory problems. **raxdfrloc** indicates whether the respondent had difficulty remembering where they were. **raxdfryr** indicates whether the respondent had difficulty remembering what year it was. **raxdfrfam** indicates whether the respondent had difficulty recognizing family members or good friends. **raxdfrloc**, **raxdfryr**, and **raxdfrfam** are coded as 0 if the respondent had no difficulty with the specific activity, and are coded as 1 if the respondent had difficulty with the specific activity. **raxdfrloc**, **raxdfryr**, and **raxdfrfam** are assigned special missing (.q) in Wave 2 when these questions were not asked. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Cross Wave Differences in SHARE EOL

Questions about having difficulty remembering where they were, remembering what year it was, and recognizing family members or good friends are asked starting in Wave 3.

Differences with the Harmonized HRS End of Life

The HRS exit interview asks different questions to ascertain the respondent's cognitive status, so there are no comparable variables in the Harmonized HRS End of Life to those found in the Gateway Harmonized SHARE End of Life.

Categorical Variable Frequencies

raxdfrloc	
0.no	14,359
1.yes	5,572
.d:dk	271
.m:missing	118
.q:not asked this wave	726
.r:refuse	52
Total	21,098
raxdfryr	
0.no	14,660
1.yes	5,126
.d:dk	403
.m:missing	132
.q:not asked this wave	726
.r:refuse	51
Total	21,098
raxdfrfam	
0.no	16,292
1.yes	3,661
.d:dk	252

.m:missing	118
.q:not asked this wave	726
.r:refuse	49
Total	21,098

Original SHARE Variables Used

Wave 3 End of Life:

sl_xt105_	Difficulties remembering where
sl_xt106_	Difficulties remembering the year
sl_xt107_	Difficulties recognizing

Wave 4 End of Life:

xt105_	Difficulties remembering where he/she was
xt106_	Difficulties remembering the year
xt107_	Difficulties recognizing

Wave 5 End of Life:

xt105_	Difficulties remembering where he/she was
xt106_	Difficulties remembering the year
xt107_	Difficulties recognizing

Wave 6 End of Life:

xt105_	Difficulties remembering where he/she was
xt106_	Difficulties remembering the year
xt107_	Difficulties recognizing

Wave 7 End of Life:

xt105_	Difficulties remembering where he/she was
xt106_	Difficulties remembering the year
xt107_	Difficulties recognizing

Wave 8 End of Life:

xt105_	Difficulties remembering where he/she was
xt106_	Difficulties remembering the year
xt107_	Difficulties recognizing

Wave 9 End of Life:

xt105_	Difficulties remembering where he/she was
xt106_	Difficulties remembering the year
xt107_	Difficulties recognizing

Wave 9 COVID End of Life:

xt105_	Difficulties remembering where he/she was
xt106_	Difficulties remembering the year
xt107_	Difficulties recognizing

E. Financial and Housing Wealth

Inflation Multiplier			
Variable	Waves	Label	Type
c2004cpindex	All	2004 consumer price index, 2010=100	Cont
c2005cpindex	All	2005 consumer price index, 2010=100	Cont
c2006cpindex	All	2006 consumer price index, 2010=100	Cont
c2007cpindex	All	2007 consumer price index, 2010=100	Cont
c2008cpindex	All	2008 consumer price index, 2010=100	Cont
c2009cpindex	All	2009 consumer price index, 2010=100	Cont
c2010cpindex	All	2010 consumer price index, 2010=100	Cont
c2011cpindex	All	2011 consumer price index, 2010=100	Cont
c2012cpindex	All	2012 consumer price index, 2010=100	Cont
c2013cpindex	All	2013 consumer price index, 2010=100	Cont
c2014cpindex	All	2014 consumer price index, 2010=100	Cont
c2015cpindex	All	2015 consumer price index, 2010=100	Cont
c2016cpindex	All	2016 consumer price index, 2010=100	Cont
c2017cpindex	All	2017 consumer price index, 2010=100	Cont
c2018cpindex	All	2018 consumer price index, 2010=100	Cont
c2019cpindex	All	2019 consumer price index, 2010=100	Cont
c2020cpindex	All	2020 consumer price index, 2010=100	Cont
c2021cpindex	All	2021 consumer price index, 2010=100	Cont
c2022cpindex	All	2022 consumer price index, 2010=100	Cont

How Constructed

cYYYYcpindex is the annual consumer price index for Euros for the year of the respondent's death. cYYYYcpindex uses 2010 as its base year so the consumer price index for a respondent who died in 2010 would be 100. This consumer price index is used as an inflation multiplier for the comparison of financial values between different years for all financial variables in the Gateway Harmonized SHARE End of Life dataset. The consumer price index of the year of death has already been used to adjust the monetary values provided in the SHARE end of life survey.

cYYYYcpindex values were provided by the World Bank as part of the World Development Indicators (WDI) dataset. The index measures monthly changes in the general level of prices of goods and services that households acquire for consumption. For more information on the calculation of the consumer price index see <http://databank.worldbank.org>.

Cross Wave Differences in SHARE EOL

Consumer price index values are not based on any SHARE end of life survey question.

Differences with the Harmonized HRS End of Life

No differences known.

Continuous Variable Distributions

Variable	N	Mean	Std.Dev	Minimum	Maximum	Total missing
c2004cpindex	19,512	86.15	5.06	74.30	94.60	0
c2005cpindex	19,512	88.18	4.55	76.90	95.70	0
c2006cpindex	19,512	90.28	4.01	79.90	96.80	0
c2007cpindex	19,512	92.80	2.90	86.30	101.50	0
c2008cpindex	19,512	97.09	1.96	91.50	105.70	0

Section E. Financial and Housing Wealth

c2009cpindex	19,512	97.87	0.99	95.40	100.90	0
c2010cpindex	19,512	100.03	0.43	100.00	105.90	0
c2011cpindex	19,512	103.10	1.00	100.20	105.00	0
c2012cpindex	19,512	105.80	1.93	99.50	109.80	0
c2013cpindex	19,512	107.10	2.62	99.30	112.10	0
c2014cpindex	19,969	107.28	2.74	99.30	112.00	0
c2015cpindex	21,098	107.20	3.08	98.20	114.20	0
c2016cpindex	21,098	107.37	3.35	97.70	112.40	0
c2017cpindex	21,098	109.32	3.98	98.30	115.50	0
c2018cpindex	21,098	111.36	4.77	99.20	119.40	0
c2019cpindex	21,098	113.08	5.60	99.50	123.80	0
c2020cpindex	21,098	113.86	6.51	98.80	127.00	0
c2021cpindex	21,098	117.16	7.95	99.40	133.50	0
c2022cpindex	21,098	129.66	13.12	102.20	151.90	0

Main House: Ownership

Variable	Waves	Label	Type
raxahown	All	raxahown: r owned home at death	Categ

How Constructed

raxahown indicates whether the deceased respondent owned their home or apartment, either in total or a share of it, according to the proxy. **raxahown** is coded as 0 if the proxy reports that the respondent did not own their home or apartment, either in total or a share of it. **raxahown** is coded as 1 if the proxy reports that the respondent did own their home or apartment, either in total or a share of it. Don't know, refused, and other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Cross Wave Differences in SHARE EOL

No differences known.

Differences with the Harmonized HRS End of Life

In the SHARE end of life interview, the proxy is asked a single direct question to ascertain the deceased respondent's home ownership, the answers to which are used to assign values in **raxahown** in the Gateway Harmonized SHARE End of Life. In the HRS exit interview, the proxy is asked whether the respondent still owned their main residence when they died, if a preload indicates that the respondent owned their main residence at the previous interview. This question is used to assign values in **raxahown** in the Harmonized HRS End of Life. As such, **raxahown** captures a slightly different concept in the Harmonized HRS End of Life because it specified whether the residence was still owned at the time of death specifically and does not allow for the possibility of becoming a homeowner in the time between the previous interview and death, as compared to a more general question of homeownership without limiting the time period in the Gateway Harmonized SHARE End of Life. Additionally, **raxahown** in the Gateway Harmonized SHARE End of Life definitely captures partial ownership, whereas this is not specified for the variable in the Harmonized HRS End of Life.

Categorical Variable Frequencies

	raxahown
0.no	7,623
1.yes	12,709
.d:dk	397
.m:missing	153
.r:refuse	216
Total	21,098

Original SHARE Variables Used

Wave 2 End of Life:	
xt03o_	The deceased owned home
Wave 3 End of Life:	
sl_xt03o_	The deceased owned home
Wave 4 End of Life:	
xt03o_	The deceased owned home
Wave 5 End of Life:	
xt03o_	The deceased owned home
Wave 6 End of Life:	
xt03o_	The deceased owned home

Section E. Financial and Housing Wealth

Wave 7 End of Life:	
xt03O_	The deceased owned home
Wave 8 End of Life:	
xt03O_	The deceased owned home
Wave 9 End of Life:	
xt03O_	The deceased owned home
Wave 9 COVID End of Life:	
xt03O_	The deceased owned home

Main House: Who Received or Inherited

Variable	Waves	Label	Type
raxhomesp	All	raxhomesp: spouse received or inherited r's home	Categ
raxhomech	All	raxhomech: child/grandchild received or inherited r's home	Categ
raxhomerl	All	raxhomerl: relative received or inherited r's home	Categ
raxhomeot	All	raxhomeot: other person received or inherited r's home	Categ

How Constructed

raxhomesp, **raxhomech**, **raxhomerl**, and **raxhomeot** indicate whether a person of the specified relationship inherited the deceased respondent's home or apartment, as reported by the proxy. The proxy is asked to select from a list all those who inherited the respondent's home or apartment. **raxhomesp** indicates whether the respondent's spouse or partner inherited the home or apartment. **raxhomech** indicates whether the respondent's sons, daughters, son- or daughter-in-law, son or daughter of husband/wife/partner, or grandchildren inherited the home or apartment. **raxhomerl** indicates if siblings or other relatives inherited the home or apartment. **raxhomeot** indicates if other non-relatives inherited the home or apartment. **raxhomesp**, **raxhomech**, **raxhomerl**, and **raxhomeot** are coded as 0 if a person of the specified relationship did not inherit the home or apartment, and are coded as 1 if a person of the specified relationship did inherit the home or apartment. **raxhomesp**, **raxhomech**, **raxhomerl**, and **raxhomeot** are assigned special missing (.x) if the respondent did not own all or part of their home or apartment. Don't know, refused, and other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Cross Wave Differences in SHARE EOL

No differences known.

Differences with the Harmonized HRS End of Life

In the SHARE end of life interview, the proxy is first asked whether the respondent owned all or part of their house or apartment and then asked to select the relationship of all those who inherited the house or apartment from a list. In the HRS exit interview, the proxy is asked whether the respondent still owned their home at the time of death based on a preload indicating home ownership in the previous interview, then asked what happened to the house separately if the respondent still or no longer owned the home at the time of death. The proxy is asked the relationship of who was given or inherited the house only if the respondent gave the house to someone prior to death or if someone inherited the house after death, and not if the house was sold, foreclosed, etc. Therefore, **raxhomesp**, **raxhomech**, **raxhomerl**, and **raxhomeot** capture slightly different concepts in the Gateway Harmonized SHARE End of Life and in the Harmonized HRS End of Life because different subsets of proxies are asked the questions.

Categorical Variable Frequencies

	raxhomesp
0.no	5,784
1.yes	6,167
.d:dk	902
.m:missing	226
.r:refuse	397
.x:didn't own home	7,622
Total	21,098
	raxhomech
0.no	5,265
1.yes	6,686

.d:dk	919
.m:missing	197
.r:refuse	409
.x:didn't own home	7,622
Total	21,098

raxhomer1

o.no	11,039
1.yes	912
.d:dk	919
.m:missing	197
.r:refuse	409
.x:didn't own home	7,622
Total	21,098

raxhomeot

o.no	11,785
1.yes	166
.d:dk	902
.m:missing	226
.r:refuse	397
.x:didn't own home	7,622
Total	21,098

Original SHARE Variables Used

Wave 2 End of Life:

xto3o_	The deceased owned home
xto02_	Relationship to the deceased
xto32d1	Who inherited the home of the deceased: yourself (proxy)
xto32d2	Who inherited the home of the deceased: spouse/partner of the deceased
xto32d3	Who inherited the home of the deceased: sons/daughters
xto32d4	Who inherited the home of the deceased: grandchildren of the deceased
xto32d5	Who inherited the home of the deceased: siblings of the deceased
xto32d6	Who inherited the home of the deceased: other relatives
xto32d7	Who inherited the home of the deceased: other non-relatives

Wave 3 End of Life:

sl_xto3o_	The deceased owned home
sl_xto02_	Relationship to the deceased
sl_xto32d1	Who inherited home of the deceased: proxy respondent
sl_xto32d2	Who inherited home of the deceased: husband/wife/partner of the deceased
sl_xto32d3	Who inherited home of the deceased: sons or daughters
sl_xto32d4	Who inherited home of the deceased: grandchildren
sl_xto32d5	Who inherited home of the deceased: siblings
sl_xto32d6	Who inherited home of the deceased: other relatives
sl_xto32d7	Who inherited home of the deceased: other non-relatives

Wave 4 End of Life:

xto3o_	The deceased owned home
xto02_	Relationship to the deceased
xto32d1	Who inherited the home of the deceased: yourself (proxy)
xto32d2	Who inherited the home of the deceased: spouse/partner of the deceased
xto32d3	Who inherited the home of the deceased: sons/daughters

xto32d4	Who inherited the home of the deceased: grandchildren of the deceased
xto32d5	Who inherited the home of the deceased: siblings of the deceased
xto32d6	Who inherited the home of the deceased: other relatives
xto32d7	Who inherited the home of the deceased: other non-relatives
Wave 5 End of Life:	
xto3o_	The deceased owned home
xto02_	Relationship to the deceased
xto32d1	Who inherited the home of the deceased: yourself (proxy)
xto32d2	Who inherited the home of the deceased: spouse/partner of the deceased
xto32d3	Who inherited the home of the deceased: sons/daughters
xto32d4	Who inherited the home of the deceased: grandchildren of the deceased
xto32d5	Who inherited the home of the deceased: siblings of the deceased
xto32d6	Who inherited the home of the deceased: other relatives
xto32d7	Who inherited the home of the deceased: other non-relatives
Wave 6 End of Life:	
xto3o_	The deceased owned home
xto02_	Relationship to the deceased
xto32d1	Who inherited the home of the deceased: yourself (proxy)
xto32d2	Who inherited the home of the deceased: spouse/partner of the deceased
xto32d3	Who inherited the home of the deceased: sons/daughters
xto32d4	Who inherited the home of the deceased: grandchildren of the deceased
xto32d5	Who inherited the home of the deceased: siblings of the deceased
xto32d6	Who inherited the home of the deceased: other relatives
xto32d7	Who inherited the home of the deceased: other non-relatives
Wave 7 End of Life:	
xto3o_	The deceased owned home
xto02_	Relationship to the deceased
xto32d1	Who inherited the home of the deceased: yourself (proxy)
xto32d2	Who inherited the home of the deceased: spouse/partner of the deceased
xto32d3	Who inherited the home of the deceased: sons/daughters
xto32d4	Who inherited the home of the deceased: grandchildren of the deceased
xto32d5	Who inherited the home of the deceased: siblings of the deceased
xto32d6	Who inherited the home of the deceased: other relatives
xto32d7	Who inherited the home of the deceased: other non-relatives
Wave 8 End of Life:	
xto3o_	The deceased owned home
xto02_	Relationship to the deceased
xto32d1	Who inherited the home of the deceased: yourself (proxy)
xto32d2	Who inherited the home of the deceased: spouse/partner of the deceased
xto32d3	Who inherited the home of the deceased: sons/daughters
xto32d4	Who inherited the home of the deceased: grandchildren of the deceased
xto32d5	Who inherited the home of the deceased: siblings of the deceased
xto32d6	Who inherited the home of the deceased: other relatives
xto32d7	Who inherited the home of the deceased: other non-relatives
Wave 9 End of Life:	
xto3o_	The deceased owned home
xto02_	Relationship to the deceased
xto32d1	Who inherited the home of the deceased: yourself (proxy)
xto32d2	Who inherited the home of the deceased: spouse/partner of the deceased
xto32d3	Who inherited the home of the deceased: sons/daughters
xto32d4	Who inherited the home of the deceased: grandchildren of the deceased
xto32d5	Who inherited the home of the deceased: siblings of the deceased
xto32d6	Who inherited the home of the deceased: other relatives

xt032d7	Who inherited the home of the deceased: other non-relatives
Wave 9 COVID End of Life:	
xt030_	The deceased owned home
xt002_	Relationship to the deceased
xt032d1	Who inherited the home of the deceased: yourself (proxy)
xt032d2	Who inherited the home of the deceased: spouse/partner of the deceased
xt032d3	Who inherited the home of the deceased: sons/daughters
xt032d4	Who inherited the home of the deceased: grandchildren of the deceased
xt032d5	Who inherited the home of the deceased: siblings of the deceased
xt032d6	Who inherited the home of the deceased: other relatives
xt032d7	Who inherited the home of the deceased: other non-relatives

Main House: Net Value

Variable	Waves	Label	Type
raxatoth	All	raxatoth: asset: net value of r's home	Cont
raxaftoth	All	raxaftoth: asset flag: net value of r's home	Categ

How Constructed

raxatoth has been adjusted to 2010 Euros based on the consumer price index for the year and country of death.

raxatoth is the net value of the respondent's home or apartment or the share of it owned by the deceased, that is the value after any outstanding mortgages, and includes imputed values to address item-level missingness. Starting in Wave 3, if the proxy answers don't know or refuses to answer, a series of unfolding bracket questions is asked to obtain a minimum and maximum bracket value for the net value of the main residence. **raxatoth** reported home net values are elicited in Euros in all SHARE countries except in Denmark, Sweden, Switzerland, Poland, Czech Republic, and Israel where local currencies are used and have been converted to Euros by the SHARE team; however all reported and imputed values in **raxatoth** have been adjusted to 2010 Euros based on the consumer price index for the year and country of death. **raxatoth** is assigned a value of 0 if the deceased respondent did not own their home or apartment, either in total or a share of it. **raxatoth** is assigned special missing (.m) if the values are not able to be imputed due to a lack of reference observations in the donor pool or due to one or more missing values in the covariates used in the imputation procedure.

In Wave 2, the proxy is not asked any additional questions if the proxy does not give an exact amount for net home value. For this wave, don't know and refused answers are imputed with no bracket information. In Waves 3 and onward, if the proxy does not give an exact amount for the net value of the respondent's residence, the proxy is asked a set of bracket questions and answers if the true value was less than, about equal to, or more than the threshold value. In Wave 3, the threshold values vary by country, even within Euro-using countries. In Wave 4, the threshold values vary by country, but all Euro-using countries use the same threshold values. Starting in Wave 5, the threshold values vary by country, even within Euro-using countries, and the values used across waves are not the same. This bracket information is used in the imputation of **raxatoth**.

raxaftoth is a flag variable indicating the level of imputation used for **raxatoth**. A code of 1 indicates the proxy reported a continuous value and no imputation was necessary. A code of 2 indicates that the value was imputed based on a closed bracket. A code of 3 indicates that the value was imputed based on an open bracket. A code of 5 indicates that the value was imputed without any bracket information. A code of 6 indicates the proxy reported the respondent not owning a residence and the value is 0. A code of 7 indicates that the ownership of the residence is unknown. A code of -1 indicates that the value could not be imputed due to a lack of reference observations in the donor pool. A code of -2 indicates that the value could not be imputed due to one or more missing values in the covariates used in the imputation procedure.

Please note that there are several extremely high values for **raxatoth** especially in Wave 4, which impact the mean value. We have included these values as reported, adjusted by the appropriate consumer price index, and leave their inclusion in any analysis to the discretion of the user.

Cross Wave Differences in SHARE EOL

In Wave 2, the proxy is not asked any additional questions if the proxy does not give an exact amount for net home value. For this wave, don't know and refused answers are imputed with no bracket information. In Wave 3, the threshold values vary by country, even within Euro-using countries. In Wave 4, the threshold values vary by country, but all Euro-using countries use the same threshold values. Starting in Wave 5, the threshold values vary by country, even within Euro-using countries, and the values used change between waves.

Differences with the Harmonized HRS End of Life

The Gateway Harmonized SHARE End of Life has the net value of the respondent's home or apartment, meaning the value after any outstanding mortgages, in **raxatoth**. There is no comparable value in the Harmonized HRS End of Life, rather that dataset contains the value of the respondent's home, not taking into account any outstanding mortgages, in **raxahous**.

The value of the respondent's house in the Harmonized HRS End of Life is denominated in 2010 U.S. dollars. The net value of the respondent's house in the Gateway Harmonized SHARE End of Life is denominated in 2010 Euros.

Additionally, if there were insufficient observations for the donor pool by country in the SHARE or if there were one or more missing values in the covariates used in the imputation procedure, then the net value of the home could not be imputed and is assigned special missing (.m) in the Gateway Harmonized SHARE End of Life, whereas the home value is either reported or imputed in the Harmonized HRS End of Life.

Continuous Variable Distributions

Variable	N	Mean	Std.Dev	Minimum	Maximum	Total missing
raxatoth	21,029	77,551.80	394,127.91	-44,802,868.00	9,503,696.00	69

Categorical Variable Frequencies

	raxaftoth
-2.not imputed, missing cov	66
-1.not imputed, missing nei	3
1.continuous value	7,252
2.closed bracket	2,219
3.open bracket	622
5.no bracket info	2,604
6.no asset	7,623
7.ownership unknown	709
Total	21,098

Original SHARE Variables Used

Wave 2 End of Life:

xto3o_	The deceased owned home
xto31e	Value of home after mortgages

Wave 3 End of Life:

sl_xto3o_	The deceased owned home
sl_xto31e	Value home after mortgages
sl_xto31ub	Value home after mortgages ub
sl_xto31v1	Bracket value 1
sl_xto31v2	Bracket value 2
sl_xto31v3	Bracket value 3

Wave 4 End of Life:

xto3o_	The deceased owned home
xto31e	Value of home after mortgages
xto31ub	Values home after mortgages ub
xto31v1	Bracket value 1
xto31v2	Bracket value 2
xto31v3	Bracket value 3

Wave 5 End of Life:

xto3o_	The deceased owned home
xto31e	Value of home after mortgages
xto31ub	Values home after mortgages ub
xto31v1	Bracket value 1
xto31v2	Bracket value 2
xto31v3	Bracket value 3

Wave 6 End of Life:

xto3o_	The deceased owned home
xto31e	Value of home after mortgages
xto31ub	Values home after mortgages ub
xto31v1	Bracket value 1
xto31v2	Bracket value 2
xto31v3	Bracket value 3

Wave 7 End of Life:

xto3o_	The deceased owned home
xto31e	Value of home after mortgages
xto31ub	Values home after mortgages ub
xto31v1	Bracket value 1
xto31v2	Bracket value 2
xto31v3	Bracket value 3

Wave 8 End of Life:

xto3o_	The deceased owned home
xto31e	Value of home after mortgages
xto31ub	Values home after mortgages ub
xto31v1	Bracket value 1
xto31v2	Bracket value 2
xto31v3	Bracket value 3

Wave 9 End of Life:

xto3o_	The deceased owned home
xto31e	Value of home after mortgages
xto31ub	Values home after mortgages ub
xto31v1	Bracket value 1
xto31v2	Bracket value 2
xto31v3	Bracket value 3

Wave 9 COVID End of Life:

xto3o_	The deceased owned home
xto31e	Value of home after mortgages
xto31ub	Values home after mortgages ub
xto31v1	Bracket value 1
xto31v2	Bracket value 2
xto31v3	Bracket value 3

Value of Assets: Individual Assets			
Variable	Waves	Label	Type
raxabsns	All	raxabsns: asset: business	Cont
raxafbsns	All	raxafbsns: asset flag: business	Categ
raxarles	All	raxarles: asset: other real estate	Cont
raxafirles	All	raxafirles: asset flag: other real estate	Categ
raxatran	All	raxatran: asset: cars	Cont
raxaftran	All	raxaftran: asset flag: cars	Categ
raxafnas	All	raxafnas: asset: financial assets	Cont
raxaffnas	All	raxaffnas: asset flag: financial assets	Categ
raxaphas	All	raxaphas: asset: physical assets	Cont
raxafphas	All	raxafphas: asset flag: physical assets	Categ

How Constructed

raxabsns, **raxarles**, **raxatran**, **raxafnas**, and **raxaphas** have been adjusted to 2010 Euros based on the consumer price index for the year and country of death.

raxabsns, **raxarles**, **raxatran**, **raxafnas**, and **raxaphas** are the values of the respondent's specific assets and include imputed values to address item-missingness. The proxy is first asked whether the respondent owned any of the following types of assets: businesses including land or premises, other real estate, cars except leased cars, financial assets (e.g. cash, bonds or stocks), and jewelry or antiquities. If the proxy reported that the respondent owned the asset, the proxy is then asked for the value of the asset at the time of their death. Starting in Wave 3, if the proxy answers don't know or refuses to answer for the value of the specific asset, a series of unfolding bracket questions is asked to obtain a minimum and maximum bracket value for the value of each asset. If the respondent did not own one of the assets, then the value for that asset is assumed to be 0. These variables are assigned special missing (.m) if the values are not able to be imputed due to a lack of reference observations in the donor pool or due to one or more missing values in the covariates used in the imputation procedure.

raxabsns is the value of the respondent's business, including land or premises, and includes imputed values. **raxarles** is the value of the respondent's other real estate, and includes imputed values. **raxatran** is the value of the respondent's cars, and includes imputed values. **raxafnas** is the value of the respondent's financial assets, e.g. cash, money, or stocks, and includes imputed values. **raxaphas** is the value of the respondent's physical assets, such as jewelry or antiquities, and includes imputed values.

In Wave 2, the proxy is not asked any additional questions if the proxy does not give an exact amount for each asset value. For this wave, don't know and refused answers are imputed with no bracket information. In Waves 3 and onward, if the proxy does not give an exact amount for the value of each asset, the proxy is asked a set of bracket questions and answers if the true value was less than, about equal to, or more than the threshold value. In Wave 3 and starting in Wave 5, the threshold values vary by country, even within Euro-using countries, and vary by asset type. In Wave 4, the threshold values vary by country, but all Euro-using countries use the same threshold values, and vary by asset type. This bracket information is used in the imputation of **raxabsns**, **raxarles**, **raxatran**, **raxafnas**, and **raxaphas** to address item-level missingness in these variables.

raxafbsns, **raxafirles**, **raxaftran**, **raxaffnas**, and **raxafphas** are flag variables indicating the level of imputation used for the specified asset. A code of 1 indicates the proxy reported only continuous values and no imputation was necessary. A code of 2 indicates that the value was imputed based on a closed bracket. A code of 3 indicates that the value was imputed based on an open bracket. A code of 5 indicates that the value was imputed without any bracket information. A code of 6 indicates the proxy reported the respondent not having the asset and the value is 0. A code of 7 indicates that the ownership of the asset is unknown. A code of -1 indicates that the value could not be imputed due to a lack of reference observations

in the donor pool. A code of -2 indicates that the value could not be imputed due to one or more missing values in the covariates used in the imputation procedure.

Please note that there are several extremely high values for some of these variables. We have included these values as reported, adjusted by the appropriate consumer price index, and leave their inclusion in any analysis to the discretion of the user.

Cross Wave Differences in SHARE EOL

In Waves 2 through 5, the proxy is asked if the respondent owned "cars". Starting in Wave 6, the proxy is asked if the respondent owned "cars, except leased cars".

In Wave 2, the proxy is not asked any additional questions if the proxy does not give an exact amount for each asset value. For this wave, don't know and refused answers are imputed with no bracket information. In Wave 3 and starting in Wave 5, the threshold values vary by country, even within Euro-using countries, and vary by asset type. In Wave 4, the threshold values vary by country, but all Euro-using countries use the same threshold values, and vary by asset type.

Differences with the Harmonized HRS End of Life

The HRS exit interview asks for the value of the respondent's total estate, excluding life insurance, in a single question, which is imputed in **raxestatev** for the Harmonized HRS End of Life. Please note that some waves exclude the value of the respondent's home, while other waves do not exclude the value of the respondent's home in the question text in the HRS exit interview. The SHARE end of life interview asks for the value of 5 specific assets after any outstanding debts, which are imputed for the Gateway Harmonized SHARE End of Life. As such, there are no comparable variables in the Harmonized HRS End of Life to these variables in the Gateway Harmonized SHARE End of Life.

Continuous Variable Distributions

Variable	N	Mean	Std.Dev	Minimum	Maximum	Total missing
raxabsns	21,016	6,599.65	80,584.54	-62,141.45	3,479,188.75	82
raxarles	21,023	9,815.77	86,268.07	-62,826.44	4,664,179.00	75
raxatran	21,039	1,090.45	6,177.76	-17,376.20	442,477.88	59
raxafnas	21,023	14,210.08	100,991.00	-3,286.39	4,651,163.00	75
raxaphas	21,012	1,012.83	43,474.88	0.00	6,109,980.00	86

Categorical Variable Frequencies

	raxafbsns
-2.not imputed, missing cov	58
-1.not imputed, missing nei	24
1.continuous value	582
2.closed bracket	185
3.open bracket	61
5.no bracket info	304
6.no asset	19,093
7.ownership unknown	791
Total	21,098
	raxafrles
-2.not imputed, missing cov	56

-1.not imputed, missing nei	19
1.continuous value	946
2.closed bracket	201
3.open bracket	100
5.no bracket info	341
6.no asset	18,475
7.ownership unknown	960
Total	21,098

raxaftran

-2.not imputed, missing cov	55
-1.not imputed, missing nei	4
1.continuous value	3,294
2.closed bracket	402
3.open bracket	89
5.no bracket info	512
6.no asset	16,179
7.ownership unknown	563
Total	21,098

raxaffnas

-2.not imputed, missing cov	63
-1.not imputed, missing nei	12
1.continuous value	2,910
2.closed bracket	509
3.open bracket	280
5.no bracket info	967
6.no asset	14,629
7.ownership unknown	1,728
Total	21,098

raxafphas

-2.not imputed, missing cov	64
-1.not imputed, missing nei	22
1.continuous value	1,015
2.closed bracket	401
3.open bracket	119
5.no bracket info	591
6.no asset	17,277
7.ownership unknown	1,609
Total	21,098

Original SHARE Variables Used

Wave 2 End of Life:

xto37_1	Deceased owned assets: business, including land or permises
xto37_2	Deceased owned assets: other real estate
xto37_3	Deceased owned assets: cars
xto37_4	Deceased owned assets: financial assets, e.g. cash/bonds/stocks
xto37_5	Deceased owned assets: jewelry or antiques
xto38e_1	Value of assets: business, including land or permises
xto38e_2	Value of assets: other real estate

xto38e_3	Value of assets: cars
xto38e_4	Value of assets: financial assets, e.g. cash, bonds or stocks
xto38e_5	Value of assets: jewelry or antiquities
Wave 3 End of Life:	
sl_xto37_1	Deceased owned assets: businesses (incl. land or premises)
sl_xto37_2	Deceased owned assets: other real estate
sl_xto37_3	Deceased owned assets: cars
sl_xto37_4	Deceased owned assets: financial assets (e.g. cash/money/stocks)
sl_xto37_5	Deceased owned assets: jewelry or antiquities
sl_xto38e_1	Value: businesses (incl. land or premises)
sl_xto38e_2	Value: other real estate
sl_xto38e_3	Value: cars
sl_xto38e_4	Value: financial assets (e.g. cash/money/stocks)
sl_xto38e_5	Value: jewelry or antiquities
sl_xto38ub_1	Value: businesses (incl. land or premises) ub
sl_xto38ub_2	Value: other real estate ub
sl_xto38ub_3	Value: cars ub
sl_xto38ub_4	Value: financial assets (e.g. cash/money/stocks) ub
sl_xto38ub_5	Value: jewelry or antiquities ub
sl_xto38v1_1	Bracket value 1: businesses (incl. land or premises)
sl_xto38v1_2	Bracket value 1: other real estate
sl_xto38v1_3	Bracket value 1: cars
sl_xto38v1_4	Bracket value 1: financial assets (e.g. cash/money/stocks)
sl_xto38v1_5	Bracket value 1: jewelry or antiquities
sl_xto38v2_1	Bracket value 2: businesses (incl. land or premises)
sl_xto38v2_2	Bracket value 2: other real estate
sl_xto38v2_3	Bracket value 2: cars
sl_xto38v2_4	Bracket value 2: financial assets (e.g. cash/money/stocks)
sl_xto38v2_5	Bracket value 2: jewelry or antiquities
sl_xto38v3_1	Bracket value 3: businesses (incl. land or premises)
sl_xto38v3_2	Bracket value 3: other real estate
sl_xto38v3_3	Bracket value 3: cars
sl_xto38v3_4	Bracket value 3: financial assets (e.g. cash/money/stocks)
sl_xto38v3_5	Bracket value 3: jewelry or antiquities
Wave 4 End of Life:	
xto37_1	Deceased owned assets: business, including land or premises
xto37_2	Deceased owned assets: other real estate
xto37_3	Deceased owned assets: cars
xto37_4	Deceased owned assets: financial assets, e.g. cash/bonds/stocks
xto37_5	Deceased owned assets: jewelry or antiquities
xto38e_1	Value of assets: business, including land or premises
xto38e_2	Value of assets: other real estate
xto38e_3	Value of assets: cars
xto38e_4	Value of assets: financial assets, e.g. cash, bonds or stocks
xto38e_5	Value of assets: jewelry or antiquities
xto38ub_1	Value of assets: business, including land or premises ub
xto38ub_2	Value of assets: other real estate ub
xto38ub_3	Value of assets: cars ub
xto38ub_4	Value of assets: financial assets, e.g. cash, bonds or stocks ub
xto38ub_5	Value of assets: jewelry or antiquities ub
xto38v1_1	Bracket value 1: business, including land or premises
xto38v1_2	Bracket value 1: other real estate
xto38v1_3	Bracket value 1: cars

xt038v1_4	Bracket value 1: financial assets, e.g. cash/bonds/stocks
xt038v1_5	Bracket value 1: jewelry or antiques
xt038v2_1	Bracket value 2: business, including land or premises
xt038v2_2	Bracket value 2: other real estate
xt038v2_3	Bracket value 2: cars
xt038v2_4	Bracket value 2: financial assets, e.g. cash/bonds/stocks
xt038v2_5	Bracket value 2: jewelry or antiques
xt038v3_1	Bracket value 3: business, including land or premises
xt038v3_2	Bracket value 3: other real estate
xt038v3_3	Bracket value 3: cars
xt038v3_4	Bracket value 3: financial assets, e.g. cash/bonds/stocks
xt038v3_5	Bracket value 3: jewelry or antiques
Wave 5 End of Life:	
xt037_1	Deceased owned assets: business, including land or premises
xt037_2	Deceased owned assets: other real estate
xt037_3	Deceased owned assets: cars
xt037_4	Deceased owned assets: financial assets, e.g. cash/bonds/stocks
xt037_5	Deceased owned assets: jewelry or antiques
xt038e_1	Value of assets: business, including land or premises
xt038e_2	Value of assets: other real estate
xt038e_3	Value of assets: cars
xt038e_4	Value of assets: financial assets, e.g. cash, bonds or stocks
xt038e_5	Value of assets: jewelry or antiques
xt038ub_1	Value of assets: business, including land or premises ub
xt038ub_2	Value of assets: other real estate ub
xt038ub_3	Value of assets: cars ub
xt038ub_4	Value of assets: financial assets, e.g. cash, bonds or stocks ub
xt038ub_5	Value of assets: jewelry or antiques ub
xt038v1_1	Bracket value 1: business, including land or premises
xt038v1_2	Bracket value 1: other real estate
xt038v1_3	Bracket value 1: cars
xt038v1_4	Bracket value 1: financial assets, e.g. cash/bonds/stocks
xt038v1_5	Bracket value 1: jewelry or antiques
xt038v2_1	Bracket value 2: business, including land or premises
xt038v2_2	Bracket value 2: other real estate
xt038v2_3	Bracket value 2: cars
xt038v2_4	Bracket value 2: financial assets, e.g. cash/bonds/stocks
xt038v2_5	Bracket value 2: jewelry or antiques
xt038v3_1	Bracket value 3: business, including land or premises
xt038v3_2	Bracket value 3: other real estate
xt038v3_3	Bracket value 3: cars
xt038v3_4	Bracket value 3: financial assets, e.g. cash/bonds/stocks
xt038v3_5	Bracket value 3: jewelry or antiques
Wave 6 End of Life:	
xt637_1	Deceased owned assets: business, including land or premises
xt637_2	Deceased owned assets: other real estate
xt637_3	Deceased owned assets: cars
xt637_4	Deceased owned assets: financial assets, e.g. cash/bonds/stocks
xt637_5	Deceased owned assets: jewelry or antiques
xt638e_1	Value of assets: business, including land or premises
xt638e_2	Value of assets: other real estate
xt638e_3	Value of assets: cars
xt638e_4	Value of assets: financial assets, e.g. cash, bonds or stocks

xt638e_5	Value of assets: jewelry or antiques
xt638ub_1	Value of assets: business, including land or premises ub
xt638ub_2	Value of assets: other real estate ub
xt638ub_3	Value of assets: cars ub
xt638ub_4	Value of assets: financial assets, e.g. cash, bonds or stocks ub
xt638ub_5	Value of assets: jewelry or antiques ub
xt638v1_1	Bracket value 1
xt638v1_2	Bracket value 1
xt638v1_3	Bracket value 1
xt638v1_4	Bracket value 1
xt638v1_5	Bracket value 1
xt638v2_1	Bracket value 2
xt638v2_2	Bracket value 2
xt638v2_3	Bracket value 2
xt638v2_4	Bracket value 2
xt638v2_5	Bracket value 2
xt638v3_1	Bracket value 3
xt638v3_2	Bracket value 3
xt638v3_3	Bracket value 3
xt638v3_4	Bracket value 3
xt638v3_5	Bracket value 3
Wave 7 End of Life:	
xt637_1	Deceased owned assets: business, including land or premises
xt637_2	Deceased owned assets: other real estate
xt637_3	Deceased owned assets: cars
xt637_4	Deceased owned assets: financial assets, e.g. cash/bonds/stocks
xt637_5	Deceased owned assets: jewelry or antiques
xt638e_1	Value of assets: business, including land or premises
xt638e_2	Value of assets: other real estate
xt638e_3	Value of assets: cars
xt638e_4	Value of assets: financial assets, e.g. cash, bonds or stocks
xt638e_5	Value of assets: jewelry or antiques
xt638ub_1	Value of assets: business, including land or premises ub
xt638ub_2	Value of assets: other real estate ub
xt638ub_3	Value of assets: cars ub
xt638ub_4	Value of assets: financial assets, e.g. cash, bonds or stocks ub
xt638ub_5	Value of assets: jewelry or antiques ub
xt638v1_1	Bracket value 1
xt638v1_2	Bracket value 1
xt638v1_3	Bracket value 1
xt638v1_4	Bracket value 1
xt638v1_5	Bracket value 1
xt638v2_1	Bracket value 2
xt638v2_2	Bracket value 2
xt638v2_3	Bracket value 2
xt638v2_4	Bracket value 2
xt638v2_5	Bracket value 2
xt638v3_1	Bracket value 3
xt638v3_2	Bracket value 3
xt638v3_3	Bracket value 3
xt638v3_4	Bracket value 3
xt638v3_5	Bracket value 3
Wave 8 End of Life:	

xt637_1	Deceased owned assets: business, including land or premises
xt637_2	Deceased owned assets: other real estate
xt637_3	Deceased owned assets: cars
xt637_4	Deceased owned assets: financial assets, e.g. cash/bonds/stocks
xt637_5	Deceased owned assets: jewelry or antiques
xt638e_1	Value of assets: business, including land or premises
xt638e_2	Value of assets: other real estate
xt638e_3	Value of assets: cars
xt638e_4	Value of assets: financial assets, e.g. cash, bonds or stocks
xt638e_5	Value of assets: jewelry or antiques
xt638ub_1	Value of assets: business, including land or premises ub
xt638ub_2	Value of assets: other real estate ub
xt638ub_3	Value of assets: cars ub
xt638ub_4	Value of assets: financial assets, e.g. cash, bonds or stocks ub
xt638ub_5	Value of assets: jewelry or antiques ub
xt638v1_1	Bracket value 1
xt638v1_2	Bracket value 1
xt638v1_3	Bracket value 1
xt638v1_4	Bracket value 1
xt638v1_5	Bracket value 1
xt638v2_1	Bracket value 2
xt638v2_2	Bracket value 2
xt638v2_3	Bracket value 2
xt638v2_4	Bracket value 2
xt638v2_5	Bracket value 2
xt638v3_1	Bracket value 3
xt638v3_2	Bracket value 3
xt638v3_3	Bracket value 3
xt638v3_4	Bracket value 3
xt638v3_5	Bracket value 3
Wave 9 End of Life:	
xt637_1	Deceased owned assets: business, including land or premises
xt637_2	Deceased owned assets: other real estate
xt637_3	Deceased owned assets: cars
xt637_4	Deceased owned assets: financial assets, e.g. cash/bonds/stocks
xt637_5	Deceased owned assets: jewelry or antiques
xt638e_1	Value of assets: business, including land or premises
xt638e_2	Value of assets: other real estate
xt638e_3	Value of assets: cars
xt638e_4	Value of assets: financial assets, e.g. cash, bonds or stocks
xt638e_5	Value of assets: jewelry or antiques
xt638ub_1	Value of assets: business, including land or premises ub
xt638ub_2	Value of assets: other real estate ub
xt638ub_3	Value of assets: cars ub
xt638ub_4	Value of assets: financial assets, e.g. cash, bonds or stocks ub
xt638ub_5	Value of assets: jewelry or antiques ub
xt638v1_1	Bracket value 1
xt638v1_2	Bracket value 1
xt638v1_3	Bracket value 1
xt638v1_4	Bracket value 1
xt638v1_5	Bracket value 1
xt638v2_1	Bracket value 2
xt638v2_2	Bracket value 2
xt638v2_3	Bracket value 2

xt638v2_4	Bracket value 2
xt638v2_5	Bracket value 2
xt638v3_1	Bracket value 3
xt638v3_2	Bracket value 3
xt638v3_3	Bracket value 3
xt638v3_4	Bracket value 3
xt638v3_5	Bracket value 3

Wave 9 COVID End of Life:

xt637_1	Deceased owned assets: business, including land or permises
xt637_2	Deceased owned assets: other real estate
xt637_3	Deceased owned assets: cars
xt637_4	Deceased owned assets: financial assets, e.g. cash/bonds/stocks
xt637_5	Deceased owned assets: jewelry or antiquities
xt638e_1	Value of assets: business, including land or permises
xt638e_2	Value of assets: other real estate
xt638e_3	Value of assets: cars
xt638e_4	Value of assets: financial assets, e.g. cash, bonds or stocks
xt638e_5	Value of assets: jewelry or antiquities
xt638ub_1	Value of assets: business, including land or permises ub
xt638ub_2	Value of assets: other real estate ub
xt638ub_3	Value of assets: cars ub
xt638ub_4	Value of assets: financial assets, e.g. cash, bonds or stocks ub
xt638ub_5	Value of assets: jewelry or antiquities ub
xt638v1_1	Bracket value 1
xt638v1_2	Bracket value 1
xt638v1_3	Bracket value 1
xt638v1_4	Bracket value 1
xt638v1_5	Bracket value 1
xt638v2_1	Bracket value 2
xt638v2_2	Bracket value 2
xt638v2_3	Bracket value 2
xt638v2_4	Bracket value 2
xt638v2_5	Bracket value 2
xt638v3_1	Bracket value 3
xt638v3_2	Bracket value 3
xt638v3_3	Bracket value 3
xt638v3_4	Bracket value 3
xt638v3_5	Bracket value 3

Value of Assets: Total Assets			
Variable	Waves	Label	Type
raxatotb	All	raxatotb: asset: r's total assets	Cont
raxaftotb	All	raxaftotb: asset flag: r's total assets	Categ
raxatotn	All	raxatotn: asset: r's total non-housing assets	Cont
raxaftotn	All	raxaftotn: asset flag: r's total non-housing assets	Categ

How Constructed

raxatotb and **raxatotn** have been adjusted to 2010 Euros based on the consumer price index for the year and country of death.

raxatotb is the value of all the respondent's assets, as reported by the proxy, and includes imputed values to address item-missingness. The proxy is first asked whether the respondent owned any of the following types of assets: house, businesses including land or premises, other real estate, cars except leased cars, financial assets (e.g. cash, bonds or stocks), and jewelry or antiquities. If the proxy reported that the respondent owned the asset, the proxy is then asked for the value of the asset at the time of their death. Starting in Wave 3, if the proxy answers don't know or refuses to answer for the value of the specific asset, a series of unfolding bracket questions is asked to obtain a minimum and maximum bracket value for the value of each asset. If the respondent did not own one of the assets, then the value for that asset is assumed to be 0. The value of each asset is then summed to obtain a value of the respondent's total assets (**raxatotb** = **raxatoth** + **raxabsns** + **raxarles** + **raxatran** + **raxafnas** + **raxaphas**). Each asset value is elicited in Euros in all SHARE countries except in Denmark, Sweden, Switzerland, Poland, Czech Republic, and Israel where local currencies are used and values have been converted to Euros by the SHARE team, however all reported and imputed values in **raxatotb** have been adjusted to 2010 Euros based on the consumer price index for the year and country of death. **raxatotb** is assigned special missing (.m) if at least one of the comprising values could not be imputed due to a lack of reference observations in the donor pool or due to one or more missing values in the covariates used in the imputation procedure.

raxatotn is the value of the respondent's non-housing assets, as reported by the proxy, and includes imputed values to address item-missingness. The proxy is first asked whether the respondent owned any of the following types of assets: businesses including land or premises, cars except leased cars, financial assets (e.g. cash, bonds or stocks), and jewelry or antiquities. If the proxy reported that the respondent owned the asset, the proxy is then asked for the value of the asset at the time of their death. Starting in Wave 3, if the proxy answers don't know or refuses to answer for the value of the specific asset, a series of unfolding bracket questions is asked to obtain a minimum and maximum bracket value for the value of each asset. If the respondent did not own one of the assets, then the value for that asset is assumed to be 0. The value of each asset is then summed to obtain a value of the respondent's non-housing assets (**raxatotn** = **raxabsns** + **raxatran** + **raxafnas** + **raxaphas**). Each asset value is elicited in Euros in all SHARE countries except in Denmark, Sweden, Switzerland, Poland, Czech Republic, and Israel where local currencies are used and values have been converted to Euros by the SHARE team, however all reported and imputed values in **raxatotn** have been adjusted to 2010 Euros based on the consumer price index for the year and country of death. **raxatotn** is assigned special missing (.m) if at least one of the comprising values could not be imputed due to a lack of reference observations in the donor pool or due to one or more missing values in the covariates used in the imputation procedure.

raxaftotb and **raxaftotn** are flag variables indicating the level of imputation used for **raxatotb** and **raxatotn**, respectively. A code of 1 indicates the proxy reported only continuous values and no imputation was necessary. A code of 2 indicates that the value was imputed based on at least one closed bracket. A code of 3 indicates that the value was imputed based on at least one open bracket. A code of 5 indicates that the value was imputed without any bracket information for at least one component. A code of 6 indicates the proxy reported the respondent not having any assets and the value is 0. A code of 7 indicates that the ownership of at least one of the assets is unknown. A code of -1 indicates that at least one of the values could not be imputed due to a lack of reference observations in the donor pool. A code of -2 indicates that at least one of the values could not be imputed due to one or more missing values in the covariates used in the imputation procedure.

Please note that there are several extremely high values for **raxatotb** and **raxatotn** and its components. We have included

these values as reported, adjusted by the appropriate consumer price index, and leave their inclusion in any analysis to the discretion of the user.

Cross Wave Differences in SHARE EOL

In Waves 2 through 5, the proxy is asked if the respondent owned "cars". Starting in Wave 6, the proxy is asked if the respondent owned "cars, except leased cars".

In Wave 2, the proxy is not asked any additional questions if the proxy does not give an exact amount for each asset value. For this wave, don't know and refused answers are imputed with no bracket information. In Wave 3 and starting in Wave 5, the threshold values vary by country, even within Euro-using countries, and vary by asset type. In Wave 4, the threshold values vary by country, but all Euro-using countries use the same threshold values, and vary by asset type.

Differences with the Harmonized HRS End of Life

The HRS exit interview asks for the value of the respondent's total estate, excluding life insurance, in a single question, which is imputed in **raxestatev** for the Harmonized HRS End of Life. Please note that some waves exclude the value of the respondent's home, while other waves do not exclude the value of the respondent's home in the question text in the HRS exit interview. The SHARE end of life interview asks for the value of 6 specific assets after any outstanding debts, which are imputed and summed in different combinations in **raxatotb** and **raxatotn** for the Gateway Harmonized SHARE End of Life.

The value of the respondent's estate in the Harmonized HRS End of Life is denominated in 2010 U.S. dollars, whereas the value of the respondent's total assets and non-housing assets in the Gateway Harmonized SHARE End of Life are denominated in 2010 Euros.

Additionally, if there were insufficient observations for the donor pool by country in the SHARE or if there were one or more missing values in the covariates used in the imputation procedure, then asset values could not be imputed and are assigned special missing (.m) in the Gateway Harmonized SHARE End of Life, whereas all estate values were either reported or imputed in the Harmonized HRS End of Life.

Continuous Variable Distributions

Variable	N	Mean	Std.Dev	Minimum	Maximum	Total missing
raxatotb	20,949	109,517.12	445,489.16	-44,802,868.00	9,610,863.00	149
raxatotn	20,970	22,866.94	142,861.37	-61,958.68	6,120,531.50	128

Categorical Variable Frequencies

	raxaftotb
-2.not imputed, missing cov	69
-1.not imputed, missing nei	80
1.continuous value	7,955
2.closed bracket	2,541
3.open bracket	820
5.no bracket info	3,098
6.no asset	4,359
7.ownership unknown	2,176
Total	21,098
	raxaftotn
-2.not imputed, missing cov	66
-1.not imputed, missing nei	62

1.continuous value	5,007
2.closed bracket	1,118
3.open bracket	422
5.no bracket info	1,580
6.no asset	10,810
7.ownership unknown	2,033
Total	21,098

Original SHARE Variables Used

Wave 2 End of Life:

xto37_1	Deceased owned assets: business, including land or premises
xto37_2	Deceased owned assets: other real estate
xto37_3	Deceased owned assets: cars
xto37_4	Deceased owned assets: financial assets, e.g. cash/bonds/stocks
xto37_5	Deceased owned assets: jewelry or antiques
xto38e_1	Value of assets: business, including land or premises
xto38e_2	Value of assets: other real estate
xto38e_3	Value of assets: cars
xto38e_4	Value of assets: financial assets, e.g. cash, bonds or stocks
xto38e_5	Value of assets: jewelry or antiques
xto30_	The deceased owned home
xto31e	Value of home after mortgages

Wave 3 End of Life:

sl_xto37_1	Deceased owned assets: businesses (incl. land or premises)
sl_xto37_2	Deceased owned assets: other real estate
sl_xto37_3	Deceased owned assets: cars
sl_xto37_4	Deceased owned assets: financial assets (e.g. cash/money/stocks)
sl_xto37_5	Deceased owned assets: jewelry or antiques
sl_xto38e_1	Value: businesses (incl. land or premises)
sl_xto38e_2	Value: other real estate
sl_xto38e_3	Value: cars
sl_xto38e_4	Value: financial assets (e.g. cash/money/stocks)
sl_xto38e_5	Value: jewelry or antiques
sl_xto38ub_1	Value: businesses (incl. land or premises) ub
sl_xto38ub_2	Value: other real estate ub
sl_xto38ub_3	Value: cars ub
sl_xto38ub_4	Value: financial assets (e.g. cash/money/stocks) ub
sl_xto38ub_5	Value: jewelry or antiques ub
sl_xto38v1_1	Bracket value 1: businesses (incl. land or premises)
sl_xto38v1_2	Bracket value 1: other real estate
sl_xto38v1_3	Bracket value 1: cars
sl_xto38v1_4	Bracket value 1: financial assets (e.g. cash/money/stocks)
sl_xto38v1_5	Bracket value 1: jewelry or antiques
sl_xto38v2_1	Bracket value 2: businesses (incl. land or premises)
sl_xto38v2_2	Bracket value 2: other real estate
sl_xto38v2_3	Bracket value 2: cars
sl_xto38v2_4	Bracket value 2: financial assets (e.g. cash/money/stocks)
sl_xto38v2_5	Bracket value 2: jewelry or antiques
sl_xto38v3_1	Bracket value 3: businesses (incl. land or premises)
sl_xto38v3_2	Bracket value 3: other real estate
sl_xto38v3_3	Bracket value 3: cars

sl_xto38v3_4	Bracket value 3: financial assets (e.g. cash/money/stocks)
sl_xto38v3_5	Bracket value 3: jewelry or antiques
sl_xto3o_	The deceased owned home
sl_xto31e	Value home after mortgages
sl_xto31ub	Value home after mortgages ub
sl_xto31v1	Bracket value 1
sl_xto31v2	Bracket value 2
sl_xto31v3	Bracket value 3
Wave 4 End of Life:	
xt037_1	Deceased owned assets: business, including land or premises
xt037_2	Deceased owned assets: other real estate
xt037_3	Deceased owned assets: cars
xt037_4	Deceased owned assets: financial assets, e.g. cash/bonds/stocks
xt037_5	Deceased owned assets: jewelry or antiques
xt038e_1	Value of assets: business, including land or premises
xt038e_2	Value of assets: other real estate
xt038e_3	Value of assets: cars
xt038e_4	Value of assets: financial assets, e.g. cash, bonds or stocks
xt038e_5	Value of assets: jewelry or antiques
xt038ub_1	Value of assets: business, including land or premises ub
xt038ub_2	Value of assets: other real estate ub
xt038ub_3	Value of assets: cars ub
xt038ub_4	Value of assets: financial assets, e.g. cash, bonds or stocks ub
xt038ub_5	Value of assets: jewelry or antiques ub
xt038v1_1	Bracket value 1: business, including land or premises
xt038v1_2	Bracket value 1: other real estate
xt038v1_3	Bracket value 1: cars
xt038v1_4	Bracket value 1: financial assets, e.g. cash/bonds/stocks
xt038v1_5	Bracket value 1: jewelry or antiques
xt038v2_1	Bracket value 2: business, including land or premises
xt038v2_2	Bracket value 2: other real estate
xt038v2_3	Bracket value 2: cars
xt038v2_4	Bracket value 2: financial assets, e.g. cash/bonds/stocks
xt038v2_5	Bracket value 2: jewelry or antiques
xt038v3_1	Bracket value 3: business, including land or premises
xt038v3_2	Bracket value 3: other real estate
xt038v3_3	Bracket value 3: cars
xt038v3_4	Bracket value 3: financial assets, e.g. cash/bonds/stocks
xt038v3_5	Bracket value 3: jewelry or antiques
xt03o_	The deceased owned home
xt031e	Value of home after mortgages
xt031ub	Values home after mortgages ub
xt031v1	Bracket value 1
xt031v2	Bracket value 2
xt031v3	Bracket value 3
Wave 5 End of Life:	
xt037_1	Deceased owned assets: business, including land or premises
xt037_2	Deceased owned assets: other real estate
xt037_3	Deceased owned assets: cars
xt037_4	Deceased owned assets: financial assets, e.g. cash/bonds/stocks
xt037_5	Deceased owned assets: jewelry or antiques
xt038e_1	Value of assets: business, including land or premises
xt038e_2	Value of assets: other real estate

xto38e_3	Value of assets: cars
xto38e_4	Value of assets: financial assets, e.g. cash, bonds or stocks
xto38e_5	Value of assets: jewelry or antiques
xto38ub_1	Value of assets: business, including land or premises ub
xto38ub_2	Value of assets: other real estate ub
xto38ub_3	Value of assets: cars ub
xto38ub_4	Value of assets: financial assets, e.g. cash, bonds or stocks ub
xto38ub_5	Value of assets: jewelry or antiques ub
xto38v1_1	Bracket value 1: business, including land or premises
xto38v1_2	Bracket value 1: other real estate
xto38v1_3	Bracket value 1: cars
xto38v1_4	Bracket value 1: financial assets, e.g. cash/bonds/stocks
xto38v1_5	Bracket value 1: jewelry or antiques
xto38v2_1	Bracket value 2: business, including land or premises
xto38v2_2	Bracket value 2: other real estate
xto38v2_3	Bracket value 2: cars
xto38v2_4	Bracket value 2: financial assets, e.g. cash/bonds/stocks
xto38v2_5	Bracket value 2: jewelry or antiques
xto38v3_1	Bracket value 3: business, including land or premises
xto38v3_2	Bracket value 3: other real estate
xto38v3_3	Bracket value 3: cars
xto38v3_4	Bracket value 3: financial assets, e.g. cash/bonds/stocks
xto38v3_5	Bracket value 3: jewelry or antiques
xto3o_	The deceased owned home
xto31e	Value of home after mortgages
xto31ub	Values home after mortgages ub
xto31v1	Bracket value 1
xto31v2	Bracket value 2
xto31v3	Bracket value 3

Wave 6 End of Life:

xt637_1	Deceased owned assets: business, including land or premises
xt637_2	Deceased owned assets: other real estate
xt637_3	Deceased owned assets: cars
xt637_4	Deceased owned assets: financial assets, e.g. cash/bonds/stocks
xt637_5	Deceased owned assets: jewelry or antiques
xt638e_1	Value of assets: business, including land or premises
xt638e_2	Value of assets: other real estate
xt638e_3	Value of assets: cars
xt638e_4	Value of assets: financial assets, e.g. cash, bonds or stocks
xt638e_5	Value of assets: jewelry or antiques
xt638ub_1	Value of assets: business, including land or premises ub
xt638ub_2	Value of assets: other real estate ub
xt638ub_3	Value of assets: cars ub
xt638ub_4	Value of assets: financial assets, e.g. cash, bonds or stocks ub
xt638ub_5	Value of assets: jewelry or antiques ub
xt638v1_1	Bracket value 1
xt638v1_2	Bracket value 1
xt638v1_3	Bracket value 1
xt638v1_4	Bracket value 1
xt638v1_5	Bracket value 1
xt638v2_1	Bracket value 2
xt638v2_2	Bracket value 2
xt638v2_3	Bracket value 2
xt638v2_4	Bracket value 2

xt638v2_5	Bracket value 2
xt638v3_1	Bracket value 3
xt638v3_2	Bracket value 3
xt638v3_3	Bracket value 3
xt638v3_4	Bracket value 3
xt638v3_5	Bracket value 3
xto3o_	The deceased owned home
xto31e	Value of home after mortgages
xto31ub	Values home after mortgages ub
xto31v1	Bracket value 1
xto31v2	Bracket value 2
xto31v3	Bracket value 3

Wave 7 End of Life:

xt637_1	Deceased owned assets: business, including land or premises
xt637_2	Deceased owned assets: other real estate
xt637_3	Deceased owned assets: cars
xt637_4	Deceased owned assets: financial assets, e.g. cash/bonds/stocks
xt637_5	Deceased owned assets: jewelry or antiques
xt638e_1	Value of assets: business, including land or premises
xt638e_2	Value of assets: other real estate
xt638e_3	Value of assets: cars
xt638e_4	Value of assets: financial assets, e.g. cash, bonds or stocks
xt638e_5	Value of assets: jewelry or antiques
xt638ub_1	Value of assets: business, including land or premises ub
xt638ub_2	Value of assets: other real estate ub
xt638ub_3	Value of assets: cars ub
xt638ub_4	Value of assets: financial assets, e.g. cash, bonds or stocks ub
xt638ub_5	Value of assets: jewelry or antiques ub
xt638v1_1	Bracket value 1
xt638v1_2	Bracket value 1
xt638v1_3	Bracket value 1
xt638v1_4	Bracket value 1
xt638v1_5	Bracket value 1
xt638v2_1	Bracket value 2
xt638v2_2	Bracket value 2
xt638v2_3	Bracket value 2
xt638v2_4	Bracket value 2
xt638v2_5	Bracket value 2
xt638v3_1	Bracket value 3
xt638v3_2	Bracket value 3
xt638v3_3	Bracket value 3
xt638v3_4	Bracket value 3
xt638v3_5	Bracket value 3
xto3o_	The deceased owned home
xto31e	Value of home after mortgages
xto31ub	Values home after mortgages ub
xto31v1	Bracket value 1
xto31v2	Bracket value 2
xto31v3	Bracket value 3

Wave 8 End of Life:

xt637_1	Deceased owned assets: business, including land or premises
xt637_2	Deceased owned assets: other real estate
xt637_3	Deceased owned assets: cars

xt637_4	Deceased owned assets: financial assets, e.g. cash/bonds/stocks
xt637_5	Deceased owned assets: jewelry or antiques
xt638e_1	Value of assets: business, including land or premises
xt638e_2	Value of assets: other real estate
xt638e_3	Value of assets: cars
xt638e_4	Value of assets: financial assets, e.g. cash, bonds or stocks
xt638e_5	Value of assets: jewelry or antiques
xt638ub_1	Value of assets: business, including land or premises ub
xt638ub_2	Value of assets: other real estate ub
xt638ub_3	Value of assets: cars ub
xt638ub_4	Value of assets: financial assets, e.g. cash, bonds or stocks ub
xt638ub_5	Value of assets: jewelry or antiques ub
xt638v1_1	Bracket value 1
xt638v1_2	Bracket value 1
xt638v1_3	Bracket value 1
xt638v1_4	Bracket value 1
xt638v1_5	Bracket value 1
xt638v2_1	Bracket value 2
xt638v2_2	Bracket value 2
xt638v2_3	Bracket value 2
xt638v2_4	Bracket value 2
xt638v2_5	Bracket value 2
xt638v3_1	Bracket value 3
xt638v3_2	Bracket value 3
xt638v3_3	Bracket value 3
xt638v3_4	Bracket value 3
xt638v3_5	Bracket value 3
xt03o_	The deceased owned home
xt031e	Value of home after mortgages
xt031ub	Values home after mortgages ub
xt031v1	Bracket value 1
xt031v2	Bracket value 2
xt031v3	Bracket value 3

Wave 9 End of Life:

xt637_1	Deceased owned assets: business, including land or premises
xt637_2	Deceased owned assets: other real estate
xt637_3	Deceased owned assets: cars
xt637_4	Deceased owned assets: financial assets, e.g. cash/bonds/stocks
xt637_5	Deceased owned assets: jewelry or antiques
xt638e_1	Value of assets: business, including land or premises
xt638e_2	Value of assets: other real estate
xt638e_3	Value of assets: cars
xt638e_4	Value of assets: financial assets, e.g. cash, bonds or stocks
xt638e_5	Value of assets: jewelry or antiques
xt638ub_1	Value of assets: business, including land or premises ub
xt638ub_2	Value of assets: other real estate ub
xt638ub_3	Value of assets: cars ub
xt638ub_4	Value of assets: financial assets, e.g. cash, bonds or stocks ub
xt638ub_5	Value of assets: jewelry or antiques ub
xt638v1_1	Bracket value 1
xt638v1_2	Bracket value 1
xt638v1_3	Bracket value 1
xt638v1_4	Bracket value 1
xt638v1_5	Bracket value 1

xt638v2_1	Bracket value 2
xt638v2_2	Bracket value 2
xt638v2_3	Bracket value 2
xt638v2_4	Bracket value 2
xt638v2_5	Bracket value 2
xt638v3_1	Bracket value 3
xt638v3_2	Bracket value 3
xt638v3_3	Bracket value 3
xt638v3_4	Bracket value 3
xt638v3_5	Bracket value 3
xto3o_	The deceased owned home
xto31e	Value of home after mortgages
xto31ub	Values home after mortgages ub
xto31v1	Bracket value 1
xto31v2	Bracket value 2
xto31v3	Bracket value 3

Wave 9 COVID End of Life:

xt637_1	Deceased owned assets: business, including land or permises
xt637_2	Deceased owned assets: other real estate
xt637_3	Deceased owned assets: cars
xt637_4	Deceased owned assets: financial assets, e.g. cash/bonds/stocks
xt637_5	Deceased owned assets: jewelry or antiques
xt638e_1	Value of assets: business, including land or permises
xt638e_2	Value of assets: other real estate
xt638e_3	Value of assets: cars
xt638e_4	Value of assets: financial assets, e.g. cash, bonds or stocks
xt638e_5	Value of assets: jewelry or antiques
xt638ub_1	Value of assets: business, including land or permises ub
xt638ub_2	Value of assets: other real estate ub
xt638ub_3	Value of assets: cars ub
xt638ub_4	Value of assets: financial assets, e.g. cash, bonds or stocks ub
xt638ub_5	Value of assets: jewelry or antiques ub
xt638v1_1	Bracket value 1
xt638v1_2	Bracket value 1
xt638v1_3	Bracket value 1
xt638v1_4	Bracket value 1
xt638v1_5	Bracket value 1
xt638v2_1	Bracket value 2
xt638v2_2	Bracket value 2
xt638v2_3	Bracket value 2
xt638v2_4	Bracket value 2
xt638v2_5	Bracket value 2
xt638v3_1	Bracket value 3
xt638v3_2	Bracket value 3
xt638v3_3	Bracket value 3
xt638v3_4	Bracket value 3
xt638v3_5	Bracket value 3
xto3o_	The deceased owned home
xto31e	Value of home after mortgages
xto31ub	Values home after mortgages ub
xto31v1	Bracket value 1
xto31v2	Bracket value 2
xto31v3	Bracket value 3

G. Family History

Number of Children

Variable	Waves	Label	Type
raxchild	All	raxchild: r number of living children when r died	Cont

How Constructed

raxchild provides the number of living children the respondent had at the time of their death, including natural children, fostered children, adopted children, and stepchildren. **raxchild** is assigned special missing (.i) if the number reported is a series of 9's and thought to be an incorrectly entered don't know or refused code. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Cross Wave Differences in SHARE EOL

No differences known.

Differences with the Harmonized HRS End of Life

The HRS exit interview does not directly ask the proxy for the number of living children the respondent had at the time of their death; rather, the count of children is based on a count of resident children and non-resident children.

Continuous Variable Distributions

Variable	N	Mean	Std.Dev	Minimum	Maximum	Total missing
raxchild	20,387	2.15	2.11	0.00	99.00	711

Original SHARE Variables Used

Wave 2 End of Life:	
xt039_	Number of children (still alive) deceased had at the end
Wave 3 End of Life:	
sl_xt039_	Number of children the deceased had at the end
Wave 4 End of Life:	
xt039_	Number of children (still alive) deceased had at the end
Wave 5 End of Life:	
xt039_	Number of children (still alive) deceased had at the end
Wave 6 End of Life:	
xt039_	Number of children (still alive) deceased had at the end
Wave 7 End of Life:	
xt039_	Number of children (still alive) deceased had at the end
Wave 8 End of Life:	
xt039_	Number of children (still alive) deceased had at the end
Wave 9 End of Life:	
xt039_	Number of children (still alive) deceased had at the end
Wave 9 COVID End of Life:	
xt039_	Number of children (still alive) deceased had at the end

L. Assistance and Caregiving

Activities of Daily Living: Whether Received Care for ADLs

Variable	Waves	Label	Type
raxracany1y	All	raxracany1y: r recvd any care for 6 adls final 12 months	Categ

How Constructed

raxracany1y indicates whether anyone regularly provided care to the deceased respondent for difficulty with any activity of daily living (ADL) in the last twelve months of life because of a physical, mental, emotional or memory problem, as reported by the proxy. In Wave 5 through 7, the proxy is first asked whether the deceased respondent had any difficulty with activities of daily living, including dressing, walking across a room, bathing, eating, getting in and out of bed, and using the toilet. If the respondent had difficulty with at least one ADL, then the proxy is asked whether anyone regularly helped with these activities. **raxracany1y** is coded as 0 if nobody regularly provided care to the respondent for difficulty with an activity of daily living, or if the respondent had no difficulty with ADLs and the proxy was not asked if anyone helped. **raxracany1y** is coded as 1 if somebody regularly provided care to the respondent for difficulty with ADLs. This question was asked together with help received for IADLs in Waves 2 through 4 and beginning in Wave 8, so in these waves **raxracany1y** is assigned special missing (.q). Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Cross Wave Differences in SHARE EOL

Whether anyone helped the respondent with ADLs alone was not asked in Waves 2 through 4 or beginning in Wave 8, instead the proxy is asked whether anyone helped the respondent with a combination of ADLs and IADLs in these waves.

Differences with the Harmonized HRS End of Life

In the Harmonized HRS End of Life, this variable indicates whether anyone helped the respondent with at least one ADL in the last three months of life. In the HRS exit interview, the proxy is asked whether anyone helped the respondent with each ADL and this variable summarizes these responses. In the Gateway Harmonized SHARE End of Life, **raxracany1y** is taken from a single question asking whether anyone helped the respondent with ADLs in the last 12 months of life, which is asked after questions ascertaining whether the respondent had any difficulty with activities of daily living. Please note that the "1Y" is added to the SHARE variable to denote the 12 month time period.

In the HRS exit interview, the proxy is only asked whether the respondent received help with the ADLs but not whether they had difficulty with the ADLs. In the SHARE end of life interview, the proxy is asked whether the respondent received help with the ADLs and whether they had difficulty with the ADLs.

Categorical Variable Frequencies

	raxracany1y
0.no	3,592
1.yes	5,192
.d:dk	235
.m:missing	112
.q:not asked this wave	11,908
.r:refuse	59
Total	21,098

Original SHARE Variables Used

Wave 5 End of Life:

xt022_	Anyone helped with ADL
xt020dno	Difficulties doing activities: none of these

Wave 6 End of Life:

xto22_	Anyone helped with ADL
xto2odno	Difficulties doing activities: none of these

Wave 7 End of Life:

xto22_	Anyone helped with ADL
xto2odno	Difficulties doing activities: none of these

Activities of Daily Living: Informal Caregiving

Variable	Waves	Label	Type
raxracaany1y	All	raxracaany1y: r recvd informal care for 6 adls final 12 months	Categ
raxrascare1y	All	raxrascare1y: r recvd informal care from spouse for 6 adls final 12 months	Categ
raxraccare1y	All	raxraccare1y: r recvd informal care from children/grandchildren for 6 adls final	Categ
raxrarcare1y	All	raxrarcare1y: r recvd informal care from relatives for 6 adls final 12 months	Categ
raxrafcare1y	All	raxrafcare1y: r recvd informal care from non-relatives for 6 adls final 12 month	Categ

How Constructed

raxracaany1y indicates whether the respondent received any informal care for difficulty with an activity of daily living (ADL) in the last twelve months of life, as reported by the proxy. In Waves 5 through 7, if the proxy reports that the respondent had difficulty with at least one ADL and that someone helped the respondent with an activity of daily living, then the proxy is asked for the relationship of the top three people who helped the respondent. **raxracaany1y** is coded as 1 if the respondent's spouse, children, grandchildren, relatives, or non-relatives provided any informal care to the respondent. **raxracaany1y** is coded as 0 if the respondent's spouse, children, grandchildren, relatives, or non-relatives did not provide any informal care to the respondent; if nobody helped the respondent with any ADL in the last twelve months of life; or if the respondent had no difficulty with any ADL in the last twelve months of life. This question is asked together with help received for IADLs in Waves 2 through 4 and in Wave 8 onwards, and so **raxracaany1y** is assigned special missing (.q) in these waves. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

raxrascare1y, **raxraccare1y**, **raxrarcare1y**, and **raxrafcare1y** indicate whether a person with the specified relationship provided informal care to the deceased respondent for difficulty with an activity of daily living (ADL) in the last twelve months of life, as reported by the proxy. In Waves 5 through 7, if the proxy reports that the respondent had difficulty with at least one ADL and that someone helped the respondent with an activity of daily living, then the proxy is asked for the relationship of the top three people who helped the respondent. **raxrascare1y** indicates whether the respondent's spouse or partner, or if the proxy, who is the respondent's husband/wife/partner, provided informal care to the respondent for difficulty with ADLs. **raxraccare1y** indicates whether the respondent's son, son-in-law, daughter, daughter-in-law, grandson, granddaughter, or if the proxy, who is the respondent's son or daughter, son- or daughter-in-law, son or daughter of husband/wife/partner, or grandchild, provided informal care to the respondent for difficulty with ADLs. **raxrarcare1y** indicates whether the respondent's mother or father, sister, brother, other relative, or if the proxy, who is the respondent's sibling or other relative, provided care to the respondent for difficulty with ADLs. **raxrafcare1y** indicates whether a friend or neighbor of the deceased, other person, or if the proxy, who is the respondent's other non-relative, provided care to the respondent for difficulty with ADLs. **raxrascare1y**, **raxraccare1y**, **raxrarcare1y**, and **raxrafcare1y** are coded as 0 if no one with the specified relationship provided informal care to the respondent for difficulty with ADLs. **raxrascare1y**, **raxraccare1y**, **raxrarcare1y**, and **raxrafcare1y** are coded as 1 if somebody with the specified relationship provided informal care to the respondent for difficulty with ADLs. **raxrascare1y**, **raxraccare1y**, **raxrarcare1y**, and **raxrafcare1y** are coded as special missing (.x) if the respondent had no difficulty with any ADL in the last twelve months of life. **raxrascare1y**, **raxraccare1y**, **raxrarcare1y**, and **raxrafcare1y** are coded as special missing (.h) if the respondent had difficulty but did not receive any help with any ADL in the last twelve months of life. This question is not asked in Waves 2 through 4, and so **raxrascare1y**, **raxraccare1y**, **raxrarcare1y**, and **raxrafcare1y** are assigned special missing (.q) in these waves. This question is asked together with help received for IADLs in Wave 8 onwards, and so **raxrascare1y**, **raxraccare1y**, **raxrarcare1y**, and **raxrafcare1y** are assigned special missing (.q) in these waves. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Cross Wave Differences in SHARE EOL

In Waves 2, 3, 4, and 8 onwards, the proxy is asked who helped the respondent with a combination of ADLs and IADLs, so these variables are not available. In Waves 5 through 7, the proxy is asked who helped the respondent with ADLs separately from IADLs, so these variables become available.

Differences with the Harmonized HRS End of Life

In the HRS exit interview, the proxy is asked who helped the respondent with ADLs in the last three months of life, and the proxy is able to respond with up to 7 relationships. Whereas in the SHARE end of life interview, the proxy is asked who helped the respondent with ADLs in the last twelve months of life, and the proxy is able to respond with up to 3 relationships. Additionally, the HRS exit interview has more responses for the relationship between the respondent and the helper than the SHARE end of life interview, which are grouped into the same categories for informal caregivers (i.e. spouse, child/grandchild, relative, non-relatives) for the Harmonized variables.

In order to account for the difference in the reference time period, the variables in the Gateway Harmonized SHARE End of Life add "1Y" to the end of the variable names.

In the HRS exit interview, the proxy is only asked whether the respondent received help with the ADLs but not whether they had difficulty with the ADLs. In the SHARE end of life interview, the proxy is asked whether the respondent received help with the ADLs and whether they had difficulty with the ADLs. As such, the variables in the Harmonized HRS End of Life only employ a special missing (.h) indicating if they received no help, while the variables in the Gateway Harmonized SHARE End of Life employ a special missing (.h) as well as a special missing (.x) indicating if they had no difficulty.

Categorical Variable Frequencies

	raxracaany1y
0.no	4,289
1.yes	4,488
.d:dk	240
.m:missing	112
.q:not asked this wave	11,908
.r:refuse	61
Total	21,098
	raxrascare1y
0.no	3,066
1.yes	2,119
.d:dk	240
.h:no help recvd	347
.m:missing	112
.q:not asked this wave	11,908
.r:refuse	61
.x:no difficulty	3,245
Total	21,098
	raxraccare1y
0.no	2,506
1.yes	2,679
.d:dk	240
.h:no help recvd	347
.m:missing	112
.q:not asked this wave	11,908
.r:refuse	61
.x:no difficulty	3,245

Total	21,098
raxrarcare1y	
o.no	4,756
1.yes	429
.d:dk	240
.h:no help recvd	347
.m:missing	112
.q:not asked this wave	11,908
.r:refuse	61
.x:no difficulty	3,245
Total	21,098
raxrafcare1y	
o.no	4,664
1.yes	521
.d:dk	240
.h:no help recvd	347
.m:missing	112
.q:not asked this wave	11,908
.r:refuse	61
.x:no difficulty	3,245
Total	21,098

Original SHARE Variables Used

Wave 5 End of Life:

xto23d1	Who has helped with ADL: yourself (proxy respondent)
xto23d2	Who has helped with ADL: spouse/partner of deceased
xto23d3	Who has helped with ADL: mother/father of deceased
xto23d4	Who has helped with ADL: son of deceased
xto23d5	Who has helped with ADL: son-in-law of deceased
xto23d6	Who has helped with ADL: daughter of deceased
xto23d7	Who has helped with ADL: daughter-in-law of deceased
xto23d8	Who has helped with ADL: grandson of deceased
xto23d9	Who has helped with ADL: granddaughter of deceased
xto23d10	Who has helped with ADL: sister of deceased
xto23d11	Who has helped with ADL: brother of deceased
xto23d12	Who has helped with ADL: other relative
xto23d15	Who has helped with ADL: friend/neighbor of deceased
xto23dot	Who has helped with ADL: other person
xto02_	Relationship to the deceased
xto22_	Anyone helped with ADL
xto20dno	Difficulties doing activities: none of these

Wave 6 End of Life:

xto23d1	Who has helped with ADL: yourself (proxy respondent)
xto23d2	Who has helped with ADL: spouse/partner of deceased
xto23d3	Who has helped with ADL: mother/father of deceased
xto23d4	Who has helped with ADL: son of deceased
xto23d5	Who has helped with ADL: son-in-law of deceased
xto23d6	Who has helped with ADL: daughter of deceased
xto23d7	Who has helped with ADL: daughter-in-law of deceased
xto23d8	Who has helped with ADL: grandson of deceased

xt023d9	Who has helped with ADL: granddaughter of deceased
xt023d10	Who has helped with ADL: sister of deceased
xt023d11	Who has helped with ADL: brother of deceased
xt023d12	Who has helped with ADL: other relative
xt023d15	Who has helped with ADL: friend/neighbor of deceased
xt023dot	Who has helped with ADL: other person
xt002_	Relationship to the deceased
xt022_	Anyone helped with ADL
xt020dno	Difficulties doing activities: none of these

Wave 7 End of Life:

xt023d1	Who has helped with ADL: yourself (proxy respondent)
xt023d2	Who has helped with ADL: spouse/partner of deceased
xt023d3	Who has helped with ADL: mother/father of deceased
xt023d4	Who has helped with ADL: son of deceased
xt023d5	Who has helped with ADL: son-in-law of deceased
xt023d6	Who has helped with ADL: daughter of deceased
xt023d7	Who has helped with ADL: daughter-in-law of deceased
xt023d8	Who has helped with ADL: grandson of deceased
xt023d9	Who has helped with ADL: granddaughter of deceased
xt023d10	Who has helped with ADL: sister of deceased
xt023d11	Who has helped with ADL: brother of deceased
xt023d12	Who has helped with ADL: other relative
xt023d15	Who has helped with ADL: friend/neighbor of deceased
xt023dot	Who has helped with ADL: other person
xt002_	Relationship to the deceased
xt022_	Anyone helped with ADL
xt020dno	Difficulties doing activities: none of these

Activities of Daily Living: Formal Caregiving			
Variable	Waves	Label	Type
raxrafaany1y	All	raxrafaany1y: r recvd formal care for 6 adls final 12 months	Categ
raxrapfcare1y	All	raxrapfcare1y: r recvd formal care from paid caregiver for 6 adls final 12 month	Categ
raxraufcare1y	All	raxraufcare1y: r recvd formal care from unpaid caregiver for 6 adls final 12 mon	Categ

How Constructed

raxrafaany1y indicates whether the respondent received any formal care for difficulty with an activity of daily living (ADL) in the last twelve months of life, as reported by the proxy. In Waves 5, 6, and 7, if the proxy reports that the respondent had difficulty with at least one ADL and that someone helped the respondent with an activity of daily living, then the proxy is asked for the relationship of the top three people who helped the respondent. **raxrafaany1y** is coded as 1 if a professional helper or unpaid volunteer provided any formal care to the respondent. **raxrafaany1y** is coded as 0 if a professional helper or unpaid volunteer did not provide any formal care to the respondent, if nobody helped the respondent with any ADL in the last twelve months of life, or if the respondent had no difficulty with any ADL in the last twelve months of life. This question is asked together with help received for IADLs in Waves 2 through 4 and in Wave 8 onwards, and so **raxrafaany1y** is assigned special missing (.q) in these waves. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

raxrapfcare1y and **raxraufcare1y** indicate whether a person with the specified relationship provided formal care to the deceased respondent for difficulty with an activity of daily living (ADL) in the last twelve months of life, as reported by the proxy. In Waves 5, 6, and 7, if the proxy reports that the respondent had difficulty with at least one ADL and that someone helped the respondent with an activity of daily living, then the proxy is asked for the relationship of the top three people who helped the respondent. **raxrapfcare1y** indicates whether a professional helper (e.g. nurse) provided formal care to the respondent for difficulty with ADLs. **raxraufcare1y** indicates whether an unpaid volunteer provided formal care to the respondent for difficulty with ADLs. **raxrapfcare1y** and **raxraufcare1y** are coded as 0 if no one with the specified relationship provided formal care to the respondent for difficulty with ADLs. **raxrapfcare1y** and **raxraufcare1y** are coded as 1 if somebody with the specified relationship provided formal care to the respondent for difficulty with ADLs. **raxrapfcare1y** and **raxraufcare1y** are coded as special missing (.x) if the respondent had no difficulty with any ADL in the last twelve months of life. **raxrapfcare1y** and **raxraufcare1y** are coded as special missing (.h) if the respondent had difficulty but did not receive any help with any ADL in the last twelve months of life. This question is asked together with help received for IADLs in Waves 2 through 4 and in Wave 8 onwards, and so **raxrapfcare1y** and **raxraufcare1y** are assigned special missing (.q) in these waves. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Cross Wave Differences in SHARE EOL

In Waves 2, 3, 4, and 8 onwards, the proxy is asked who helped the respondent with a combination of ADLs and IADLs, so these variables are not available. In Waves 5 through 7, the proxy is asked who helped the respondent with ADLs separately from IADLs, so these variables become available.

Differences with the Harmonized HRS End of Life

In the HRS exit interview, the proxy is asked who helped the respondent with ADLs in the last three months of life, and the proxy is able to respond with up to 7 relationships. Whereas in the SHARE end of life interview, the proxy is asked who helped the respondent with ADLs in the last twelve months of life, and the proxy is able to respond with up to 3 relationships. Additionally, the HRS exit interview has more responses for the relationship between the respondent and the helper than the SHARE end of life interview, which are grouped into the same categories for formal caregivers (i.e. paid caregivers and unpaid caregivers) for the Harmonized variables.

In order to account for the difference in the reference time period, the variables in the Gateway Harmonized SHARE End of Life add "1Y" to the end of the variable names.

In the HRS exit interview, the proxy is only asked whether the respondent received help with the ADLs but not whether they had difficulty with the ADLs. In the SHARE end of life interview, the proxy is asked whether the respondent received help with the ADLs and whether they had difficulty with the ADLs. As such, the variables in the Harmonized HRS End of Life only employ a special missing (.h) indicating if they received no help, while the variables in the Gateway Harmonized SHARE End of Life employ a special missing (.h) as well as a special missing (.x) indicating if they had no difficulty.

Categorical Variable Frequencies

	raxrafaany1y
0.no	6,839
1.yes	1,938
.d:dk	240
.m:missing	112
.q:not asked this wave	11,908
.r:refuse	61
Total	21,098

	raxrapfcare1y
0.no	3,274
1.yes	1,911
.d:dk	240
.h:no help recvd	347
.m:missing	112
.q:not asked this wave	11,908
.r:refuse	61
.x:no difficulty	3,245
Total	21,098

	raxraufcare1y
0.no	5,138
1.yes	47
.d:dk	240
.h:no help recvd	347
.m:missing	112
.q:not asked this wave	11,908
.r:refuse	61
.x:no difficulty	3,245
Total	21,098

Original SHARE Variables Used

Wave 5 End of Life:

xt023d13	Who has helped with ADL: unpaid volunteer
xt023d14	Who has helped with ADL: professional helper
xt002_	Relationship to the deceased
xt022_	Anyone helped with ADL
xt020dno	Difficulties doing activities: none of these

Wave 6 End of Life:

xt023d13	Who has helped with ADL: unpaid volunteer
----------	---

Section L. Assistance and Caregiving

xt023d14	Who has helped with ADL: professional helper
xt002_	Relationship to the deceased
xt022_	Anyone helped with ADL
xt020dno	Difficulties doing activities: none of these
Wave 7 End of Life:	
xt023d13	Who has helped with ADL: unpaid volunteer
xt023d14	Who has helped with ADL: professional helper
xt002_	Relationship to the deceased
xt022_	Anyone helped with ADL
xt020dno	Difficulties doing activities: none of these

Activities of Daily Living: Frequency of Caregiving			
Variable	Waves	Label	Type
raxracared	All	raxracared: duration r recvd any care for 6 adls final 12 months	Categ
raxracarehr	All	raxracarehr: hrs/day r recvd any care for 6 adls final 12 months	Cont

How Constructed

raxracared indicates how long the deceased respondent received help with any activity of daily living (ADL) because of a physical, mental, emotional or memory problem in the last year of life. The activities of daily living include dressing, walking across a room, bathing, eating, getting in and out of bed, and using the toilet. The proxy is first asked whether the respondent had difficulty with each activity of daily living, and if the respondent had difficulty with at least one ADL, then the proxy is asked if someone helped the respondent, and if so how long the respondent received help during the last twelve months of life. **raxracared** is coded as follows: 1.less than 1 month, 2.1 month or more but less than 3 months, 3.3 months or more but less than 6 months, 4.6 months or more but less than 1 year, 5.a full year. **raxracared** is coded as special missing (.x) if the respondent did not need help with any ADL activity. **raxracared** is coded as special missing (.h) if the respondent had difficulty but did not receive any help with any ADL activity. This question is not asked of ADL activities separately in Waves 2, 3, 4, and 8 onwards, and so **raxracared** is assigned special missing (.q) in these waves. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

raxracarehr indicates the number of hours of help that were necessary during a typical day for help with any activity of daily living because of a physical, mental, emotional or memory problem in the last year of life. The activities of daily living include dressing, walking across a room, bathing, eating, getting in and out of bed, and using the toilet. The proxy is first asked whether the respondent had difficulty with each activity of daily living, and if the respondent had difficulty with at least one ADL, then the proxy is asked if someone helped the respondent, and if so about how many hours of help were necessary during a typical day. **raxracarehr** takes the number of hours reported by the proxy, ranging from 0 to 24. **raxracarehr** is coded as special missing (.x) if the respondent did not need help with any ADL activity. **raxracarehr** is coded as special missing (.h) if the respondent had difficulty but did not receive any help with any ADL activity. This question is not asked of ADL activities separately in Waves 2, 3, 4, and 8 onwards, and so **raxracarehr** is assigned special missing (.q) in these waves. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Cross Wave Differences in SHARE EOL

In Waves 2, 3, 4, and 8 onwards, how long the respondent received help is asked for a combination of ADL and IADL activities, so these variables are not available. In Waves 5, 6, and 7, how long the respondent received help is asked for ADL activities separately from IADL activities, so these variables become available.

Differences with the Harmonized HRS End of Life

In the HRS exit interview, the proxy is asked whether the respondent received help with each ADL activity in the final three months of life, and if so, how frequently the respondent received help from each individual helper, which can be provided as days in a typical month, days per week, or everyday. These values are transformed into Harmonized variables indicating the number of days per month for each category of helper for the Harmonized HRS End of Life. However, in the SHARE end of life interview, the proxy is asked whether the respondent received help with a list of ADL activities in the last twelve months of life, and if so, asked how long the respondent had received help with any of the activities in the last year, which has specific response categories with a maximum of one year, and are reported as is and as a total for all helpers in the Gateway Harmonized SHARE End of Life. As such, these harmonized variables would need adjustment before being comparable.

In the HRS exit interview, the proxy is only asked whether the respondent received help with the ADLs but not whether they had difficulty with the ADLs. In the SHARE end of life interview, the proxy is asked whether the respondent received help with the ADLs and whether they had difficulty with the ADLs. As such, the variables in the Harmonized HRS End of Life only employ a special missing (.h) indicating if they received no help, while the variables in the Gateway Harmonized SHARE End of Life employ a special missing (.h) as well as a special missing (.x) indicating if they had no difficulty.

Continuous Variable Distributions

Variable	N	Mean	Std.Dev	Minimum	Maximum	Total missing
raxracarehr	4,722	12.29	9.37	0.00	24.00	16,376

Categorical Variable Frequencies

	raxracared
1.less than 1 month	265
2.1 month+ but less than 3	487
3.3 months+ but less than 6	605
4.6 months+ but less than 1	758
5.a full year	3,051
.d:dk	250
.h:no help recvd	347
.m:missing	118
.q:not asked this wave	11,908
.r:refuse	64
.x:no difficulty	3,245
Total	21,098

Original SHARE Variables Used

Wave 5 End of Life:

xto22_	Anyone helped with ADL
xto24_	Time the deceased received help
xto25_	Hours of help necessary during typical day
xto2odno	Difficulties doing activities: none of these

Wave 6 End of Life:

xto22_	Anyone helped with ADL
xto24_	Time the deceased received help
xto25_	Hours of help necessary during typical day
xto2odno	Difficulties doing activities: none of these

Wave 7 End of Life:

xto22_	Anyone helped with ADL
xto24_	Time the deceased received help
xto25_	Hours of help necessary during typical day
xto2odno	Difficulties doing activities: none of these

Activities of Daily Living: Whether Began Receiving Help within Final Year

Variable	Waves	Label	Type
raxracareb	All	raxracareb: r began recv help with 6 adls final 12 months	Categ

How Constructed

raxracareb indicates whether the deceased respondent began receiving help with activities of daily living (ADLs) because of a physical, mental, emotional or memory problem in the last year of life. The activities of daily living include dressing, walking across a room, bathing, eating, getting in and out of bed, and using the toilet. The proxy is first asked whether the respondent had difficulty with each activity of daily living, and if the respondent had difficulty with at least one ADL, then the proxy is asked if someone helped the respondent, and if so how long the respondent received help during the last twelve months of life. **raxracareb** is coded as 0 if the respondent needed help with at least one ADL activity and the respondent required help for the full year. **raxracareb** is coded as 1 if the respondent needed help with at least one ADL activity and the respondent received help for less than one month, one month or more but less than three months, three months or more but less than six months, or six months or more but less than a year. **raxracareb** is coded as special missing (.x) if the respondent did not need help with any ADL activity. **raxracareb** is coded as special missing (.h) if the respondent had difficulty but did not receive any help with any ADL activity. This question is not asked of ADL activities separately in Waves 2 through 4, and Wave 8 onwards, and so **raxracareb** is assigned special missing (.q) in these waves. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Cross Wave Differences in SHARE EOL

In Waves 2, 3, 4, and 8 onwards, how long the respondent received help is asked for a combination of ADL and IADL activities, so this variable is not available. In Wave 5 through 7, how long the respondent received help is asked for ADL activities separately from IADL activities, and so becomes available.

Differences with the Harmonized HRS End of Life

In the HRS exit interview, the proxy is asked whether the respondent received help with each ADL activity in the final three months of life, and if so, asked how long the respondent had needed help with each ADL, which can be provided in months, years, since age, or since year. These responses have been converted to be equivalent to approximately 1 year in the Harmonized HRS End of Life. However, in the SHARE end of life interview, the proxy is asked whether the respondent received help with a list of ADL activities in the last twelve months of life, and if so, asked how long the respondent had received help with any of the activities in the last year, which has specific response categories with a maximum of one year. This response has been converted so that responses of a full year are coded as not having begun to need help in the last year of life in the Gateway Harmonized SHARE End of Life. As such, this variable in the Harmonized HRS End of Life captures a slightly different concept than the same variable in the Gateway Harmonized SHARE End of Life because we can ascertain with some degree of certainty when the respondent began needing help based on the questions in the HRS exit interview, whereas we are assuming that the respondent began needing help prior to the last twelve months of life if the respondent received help during their entire final year in the SHARE end of life interview.

In the HRS exit interview, the proxy is only asked whether the respondent received help with the ADLs but not whether they had difficulty with the ADLs. In the SHARE end of life interview, the proxy is asked whether the respondent received help with the ADLs and whether they had difficulty with the ADLs. As such, the variable in the Harmonized HRS End of Life only employs a special missing (.h) indicating if they received no help, while the variable in the Gateway Harmonized SHARE End of Life employs a special missing (.h) as well as a special missing (.x) indicating if they had no difficulty.

Categorical Variable Frequencies

	raxracareb
0.no	3,051
1.yes	2,115
.d:dk	250

.h:no help recvd	347
.m:missing	118
.q:not asked this wave	11,908
.r:refuse	64
.x:no difficulty	3,245
Total	21,098

Original SHARE Variables Used

Wave 5 End of Life:

xto22_	Anyone helped with ADL
xto24_	Time the deceased received help
xto2odno	Difficulties doing activities: none of these

Wave 6 End of Life:

xto22_	Anyone helped with ADL
xto24_	Time the deceased received help
xto2odno	Difficulties doing activities: none of these

Wave 7 End of Life:

xto22_	Anyone helped with ADL
xto24_	Time the deceased received help
xto2odno	Difficulties doing activities: none of these

Instrumental Activities of Daily Living: Whether Received Care for IADLs

Variable	Waves	Label	Type
raxricany1y_s	All	raxricany1y_s: r recvd any care for 9 iadls + cont final 12 months	Categ

How Constructed

raxricany1y_s indicates whether anyone provided care to the deceased respondent for difficulty with any instrumental activity of daily living (IADL) or with continence over urination or defecation in the last twelve months of life because of a physical, mental, emotional or memory problem, as reported by the proxy. In Waves 6 and 7, the proxy is first asked whether the deceased respondent had any difficulty with instrumental activities of daily living, including preparing hot meals, shopping for groceries, using the telephone, taking medication, using a map, doing work around the house or garden, managing money, leaving the house independently, and doing personal laundry, and any difficulty with continence over urination or defecation. If the respondent had difficulty with at least one IADL or with continence over urination or defecation, then the proxy is asked whether anyone regularly helped with these activities. **raxricany1y_s** is coded as 0 if nobody provided care to the respondent for difficulty with an IADL or with continence over urination or defecation, or if the respondent had no difficulty with IADLs or with continence over urination or defecation and the proxy is not asked if anyone helped. **raxricany1y_s** is coded as 1 if somebody provided care to the respondent for difficulty with IADLs or with continence over urination or defecation. This question was asked together with help received for ADLs in Waves 2 through 4 and in Wave 8 onwards, and not at all in Wave 5, so in these waves **raxricany1y_s** is assigned special missing (.q). Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Cross Wave Differences in SHARE EOL

Whether anyone helped the respondent with IADLs alone was not asked in Waves 2 through 4 or in Wave 8 onwards, instead the proxy is asked whether anyone helped the respondent with a combination of ADLs and IADLs in these waves, so this variable is not available. In Wave 5, the proxy is not asked about the respondent's difficulty with IADL activities at all. In Waves 6 and 7, IADLs are asked about separately from ADLs and this variable is available.

Differences with the Harmonized HRS End of Life

In the Harmonized HRS End of Life, this variable indicates whether anyone helped the respondent with at least one of four IADLs (preparing hot meals, shopping for groceries, using the telephone, taking medication) in the last three months of life. In the HRS exit interview, the proxy is asked whether anyone helped the respondent with each IADL and this variable summarizes these responses. In the Gateway Harmonized SHARE End of Life, this variable is taken from a single question asking whether anyone helped the respondent with 9 IADLs and continence over urination or defecation in the last 12 months of life, which is asked after questions ascertaining whether the respondent had any difficulty with these activities. Because this variable in the Harmonized HRS End of Life captures a similar concept but with four IADLs in the final three months, while the Gateway Harmonized SHARE End of Life has 9 IADLs and continence over urination or defecation in the final 12 months, the "1Y_S" is added to the SHARE variable to denote the difference in time period and additional activities.

Categorical Variable Frequencies

	raxricany1y_s
0.no	2,272
1.yes	4,500
.d:dk	174
.m:missing	17
.q:not asked this wave	14,095
.r:refuse	40
Total	21,098

Original SHARE Variables Used

Wave 6 End of Life:

xt622_

Anyone helped with ADLII

xt62odno

Difficulties doing activities II: none of these

Wave 7 End of Life:

xt622_

Anyone helped with ADLII

xt62odno

Difficulties doing activities II: none of these

Instrumental Activities of Daily Living: Informal Caregiving			
Variable	Waves	Label	Type
raxricaanyly_s	All	raxricaanyly_s: r recvd informal care for 9 iadls + cont final 12 months	Categ
raxriscarely_s	All	raxriscarely_s: r recvd informal care from spouse for 9 iadls + cont final 12 mo	Categ
raxriccarely_s	All	raxriccarely_s: r recvd informal care from children/grandchildren for 9 iadls +	Categ
raxrircarely_s	All	raxrircarely_s: r recvd informal care from relatives for 9 iadls + cont final 12	Categ
raxrifcarely_s	All	raxrifcarely_s: r recvd informal care from non-relatives for 9 iadls + cont fina	Categ

How Constructed

raxricaanyly_s indicates whether the respondent received any informal care for difficulty with an instrumental activity of daily living (IADL) or with continence over urination or defecation in the last twelve months of life, as reported by the proxy. In Waves 6 and 7, if the proxy reports that someone helped the respondent with any of 9 instrumental activities of daily living, including preparing hot meals, shopping for groceries, using the telephone, taking medication, using a map, doing work around the house or garden, managing money, leaving the house independently, and doing personal laundry, or with continence over urination or defecation, then the proxy is asked for the relationship of the top three people who helped the respondent. **raxricaanyly_s** is coded as 1 if the respondent's spouse, children, grandchildren, relatives, or non-relatives provided any informal care to the respondent. **raxricaanyly_s** is coded as 0 if the respondent's spouse, children, grandchildren, relatives, or non-relatives did not provide any informal care to the respondent. **raxricaanyly_s** is also coded as 0 if nobody helped the respondent with IADLs or with continence over urination or defecation in the last twelve months of life, or if the respondent had no difficulties with IADLs or with continence over urination or defecation in the last twelve months of life. This question was asked together with help received for ADLs in Waves 2 through 4 and in Wave 8 onwards and not at all in Wave 5, so **raxricaanyly_s** is assigned special missing (.q) for these waves. Don't know, refused, and other missing responses are assigned special missing (.d), (.r), (.m), respectively.

raxriscarely_s, **raxriccarely_s**, **raxrircarely_s**, and **raxrifcarely_s** indicate whether a specific person provided informal care to the deceased respondent for difficulty with any instrumental activity of daily living (IADL) or with continence over urination or defecation, as reported by the proxy. In Waves 6 and 7, if the proxy reports that someone helped the respondent with any of 9 instrumental activities of daily living, including preparing hot meals, shopping for groceries, using the telephone, taking medication, using a map, doing work around the house or garden, managing money, leaving the house independently, and doing personal laundry, or with continence over urination or defecation, then the proxy is asked for the relationship of the top three people who helped the respondent. **raxriscarely_s** indicates whether the respondent's spouse or partner, or if the proxy, who is the respondent's husband/wife/partner, provided informal care to the respondent for difficulty with IADLs or with continence over urination or defecation. **raxriccarely_s** indicates whether the respondent's son, son-in-law, daughter, daughter-in-law, grandson, granddaughter, or if the proxy, who is the respondent's son or daughter, son- or daughter-in-law, son or daughter of husband/wife/partner, or grandchild, provided informal care to the respondent for difficulty with IADLs or with continence over urination or defecation. **raxrircarely_s** indicates whether the respondent's mother or father, sister, brother, other relative, or if the proxy, who is the respondent's sibling or other relative, provided informal care to the respondent for difficulty with IADLs or with continence over urination or defecation. **raxrifcarely_s** indicates whether a friend or neighbor of the deceased, other person, or if the proxy, who is the respondent's other non-relative, provided care to the respondent for difficulty with IADLs or with continence over urination or defecation.

raxriscarely_s, **raxriccarely_s**, **raxrircarely_s**, and **raxrifcarely_s** are coded as 0 if no one with the specified relationship provided informal care to the respondent for difficulty with IADLs or with continence over urination or defecation. **raxriscarely_s**, **raxriccarely_s**, **raxrircarely_s**, and **raxrifcarely_s** are coded as 1 if somebody with the specified relationship provided informal care to the respondent for difficulty with IADLs or with continence over

urination or defecation. **raxriscarely_s**, **raxriccarely_s**, **raxrircarely_s**, and **raxrifcarely_s** are coded as special missing (.x) if the respondent had no difficulties with IADLs or with continence over urination or defecation in the last twelve months of life. **raxriscarely_s**, **raxriccarely_s**, **raxrircarely_s**, and **raxrifcarely_s** are coded as special missing (.h) if the respondent had difficulty but did not receive help with IADLs or with continence over urination or defecation in the last twelve months of life. This question was asked together with help received for ADLs in Waves 2 through 4 and in Wave 8 onwards and not at all in Wave 5, so **raxriscarely_s**, **raxriccarely_s**, **raxrircarely_s**, and **raxrifcarely_s** are assigned special missing (.q) for these waves. Don't know, refused, and other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Cross Wave Differences in SHARE EOL

Who helped the respondent with IADL activities is asked about in Waves 6 and 7. In Waves 2, 3, 4, and 8 onwards, who helped the respondent is asked for a combination of ADL and IADL activities. In Wave 5, IADL activities are not asked about at all.

Differences with the Harmonized HRS End of Life

In the HRS exit interview, the proxy is asked who helped the respondent with 4 IADLs in the last three months of life, and the proxy is able to respond with up to 6 relationships. Whereas in the SHARE end of life interview, the proxy is asked who helped the respondent with 9 IADLs and with continence over urination or defecation in the last twelve months of life, and the proxy is able to respond with up to 3 relationships. In order to account for the difference in the reference time period and number of activities, the variables in the Gateway Harmonized SHARE End of Life add "1Y_S" to the end of the variables. Additionally, the HRS exit interview has more responses for the relationship between the respondent and the helper than the SHARE end of life interview, which are grouped into the same categories for informal caregivers (i.e. spouse, child/grandchild, relative, non-relative) for the Harmonized variables.

In the HRS exit interview, the proxy is only asked whether the respondent received help with the IADLs but not whether they had difficulty with the IADLs. In the SHARE end of life interview, the proxy is asked whether the respondent received help with the IADLs and whether they had difficulty with the IADLs. As such, the variables in the Harmonized HRS End of Life only employ a special missing (.h) indicating if they received no help, while the variables in the Gateway Harmonized SHARE End of Life employ a special missing (.h) as well as a special missing (.x) in some variables indicating if they had no difficulty.

Categorical Variable Frequencies

	raxricaanyly_s
0.no	2,753
1.yes	4,011
.d:dk	181
.m:missing	17
.q:not asked this wave	14,095
.r:refuse	41
Total	21,098
	raxriscarely_s
0.no	2,745
1.yes	1,747
.d:dk	181
.h:no help recvd	264
.m:missing	17
.q:not asked this wave	14,095
.r:refuse	41
.x:no difficulty	2,008
Total	21,098

	raxriccarely_s
0.no	1,988
1.yes	2,504
.d:dk	181
.h:no help recvd	264
.m:missing	17
.q:not asked this wave	14,095
.r:refuse	41
.x:no difficulty	2,008
Total	21,098

	raxrircarely_s
0.no	4,125
1.yes	367
.d:dk	181
.h:no help recvd	264
.m:missing	17
.q:not asked this wave	14,095
.r:refuse	41
.x:no difficulty	2,008
Total	21,098

	raxrifcarely_s
0.no	4,006
1.yes	486
.d:dk	181
.h:no help recvd	264
.m:missing	17
.q:not asked this wave	14,095
.r:refuse	41
.x:no difficulty	2,008
Total	21,098

Original SHARE Variables Used

Wave 6 End of Life:

xt623d1	Who has helped with ADLII: yourself (proxy respondent)
xt623d2	Who has helped with ADLII: spouse/partner of deceased
xt623d3	Who has helped with ADLII: mother/father of deceased
xt623d4	Who has helped with ADLII: son of deceased
xt623d5	Who has helped with ADLII: son-in-law of deceased
xt623d6	Who has helped with ADLII: daughter of deceased
xt623d7	Who has helped with ADLII: daughter-in-law of deceased
xt623d8	Who has helped with ADLII: grandson of deceased
xt623d9	Who has helped with ADLII: granddaughter of deceased
xt623d10	Who has helped with ADLII: sister of deceased
xt623d11	Who has helped with ADLII: brother of deceased
xt623d12	Who has helped with ADLII: other relative
xt623d15	Who has helped with ADLII: friend/neighbor of deceased
xt623dot	Who has helped with ADLII: other person
xt002_	Relationship to the deceased
xt622_	Anyone helped with ADLII

xt62odno	Difficulties doing activities II: none of these
Wave 7 End of Life:	
xt623d1	Who has helped with ADLII: yourself (proxy respondent)
xt623d2	Who has helped with ADLII: spouse/partner of deceased
xt623d3	Who has helped with ADLII: mother/father of deceased
xt623d4	Who has helped with ADLII: son of deceased
xt623d5	Who has helped with ADLII: son-in-law of deceased
xt623d6	Who has helped with ADLII: daughter of deceased
xt623d7	Who has helped with ADLII: daughter-in-law of deceased
xt623d8	Who has helped with ADLII: grandson of deceased
xt623d9	Who has helped with ADLII: granddaughter of deceased
xt623d10	Who has helped with ADLII: sister of deceased
xt623d11	Who has helped with ADLII: brother of deceased
xt623d12	Who has helped with ADLII: other relative
xt623d15	Who has helped with ADLII: friend/neighbor of deceased
xt623dot	Who has helped with ADLII: other person
xt002_	Relationship to the deceased
xt622_	Anyone helped with ADLII
xt62odno	Difficulties doing activities II: none of these

Instrumental Activities of Daily Living: Formal Caregiving			
Variable	Waves	Label	Type
raxrifaany1y_s	All	raxrifaany1y_s: r recvd formal care for 9 iadls + cont final 12 months	Categ
raxripfcare1y_s	All	raxripfcare1y_s: r recvd formal care from paid caregiver for 9 iadls + cont fina	Categ
raxriufcare1y_s	All	raxriufcare1y_s: r recvd formal care from unpaid caregiver for 9 iadls + cont fi	Categ

How Constructed

raxrifaany1y_s indicates whether the respondent received any formal care for difficulty with an instrumental activity of daily living (IADL) or with continence over urination or defecation in the last twelve months of life, as reported by the proxy. In Waves 6 and 7, if the proxy reports that someone helped the respondent with any of 9 instrumental activities of daily living, including preparing hot meals, shopping for groceries, using the telephone, taking medication, using a map, doing work around the house or garden, managing money, leaving the house independently, and doing personal laundry, or with continence over urination or defecation, then the proxy is asked for the relationship of the top three people who helped the respondent. **raxrifaany1y_s** is coded as 1 if a professional helper or unpaid volunteer provided any formal care to the respondent. **raxrifaany1y_s** is coded as 0 if a professional helper or unpaid volunteer did not provide any formal care to the respondent, or if the respondent had no difficulties with IADLs or with continence over urination or defecation in the last twelve months of life. **raxrifaany1y_s** is also coded as 0 if nobody helped the respondent with IADLs or with continence over urination or defecation in the last twelve months of life. This question was asked together with help received for ADLs in Waves 2 through 4 and in Wave 8 onwards and not at all in Wave 5, so **raxrifaany1y_s** is assigned special missing (.q) for these waves. Don't know, refused, and other missing responses are assigned special missing (.d), (.r), (.m), respectively.

raxripfcare1y_s and **raxriufcare1y_s** indicate whether a specific person provided formal care to the deceased respondent for difficulty with any instrumental activity of daily living (IADL) or with continence over urination or defecation, as reported by the proxy. In Waves 6 and 7, if the proxy reports that someone helped the respondent with any of 9 instrumental activities of daily living, including preparing hot meals, shopping for groceries, using the telephone, taking medication, using a map, doing work around the house or garden, managing money, leaving the house independently, and doing personal laundry, or with continence over urination or defecation, then the proxy is asked for the relationship of the top three people who helped the respondent. **raxripfcare1y_s** indicates whether a professional helper (e.g. nurse) provided formal care to the respondent for difficulty with IADLs or with continence over urination or defecation. **raxriufcare1y_s** indicates whether an unpaid volunteer provided formal care to the respondent for difficulty with IADLs or with continence over urination or defecation. **raxripfcare1y_s** and **raxriufcare1y_s** are coded as 0 if no one with the specified relationship provided formal care to the respondent for difficulty with IADLs or with continence over urination or defecation. **raxripfcare1y_s** and **raxriufcare1y_s** are coded as 1 if somebody with the specified relationship provided care to the respondent for difficulty with IADLs or with continence over urination or defecation. **raxripfcare1y_s** and **raxriufcare1y_s** are coded as special missing (.x) if the respondent had no difficulties with IADLs or with continence over urination or defecation in the last twelve months of life. **raxripfcare1y_s** and **raxriufcare1y_s** are coded as special missing (.h) if the respondent had difficulty but did not receive help with IADLs or with continence over urination or defecation in the last twelve months of life. This question was asked together with help received for ADLs in Waves 2 through 4 and in Wave 8 onwards and not at all in Wave 5, so **raxripfcare1y_s** and **raxriufcare1y_s** are assigned special missing (.q) for these waves. Don't know, refused, and other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Cross Wave Differences in SHARE EOL

Who helped the respondent with IADL activities is asked about in Waves 6 and 7. In Waves 2, 3, 4, and 8 onwards, who helped the respondent is asked for a combination of ADL and IADL activities. In Wave 5, IADL activities are not asked about at all.

Differences with the Harmonized HRS End of Life

In the HRS exit interview, the proxy is asked who helped the respondent with 4 IADLs in the last three months of life, and the proxy is able to respond with up to 6 relationships. Whereas in the SHARE end of life interview, the proxy is asked who helped the respondent with 9 IADLs and with continence over urination or defecation in the last twelve months of life, and the proxy is able to respond with up to 3 relationships. In order to account for the difference in the reference time period and number of activities, the variables in the Gateway Harmonized SHARE End of Life add "1Y_S" to the end of the variables. Additionally, the HRS exit interview has more responses for the relationship between the respondent and the helper than the SHARE end of life interview, which are grouped into the same categories for formal caregivers (i.e. paid caregivers and unpaid caregivers) for the Harmonized variables.

In the HRS exit interview, the proxy is only asked whether the respondent received help with the IADLs but not whether they had difficulty with the IADLs. In the SHARE end of life interview, the proxy is asked whether the respondent received help with the IADLs and whether they had difficulty with the IADLs. As such, the variables in the Harmonized HRS End of Life only employ a special missing (.h) indicating if they received no help, while the variables in the Gateway Harmonized SHARE End of Life employ a special missing (.h) as well as a special missing (.x) indicating if they had no difficulty.

Categorical Variable Frequencies

	raxrifaanyly_s
0.no	5,306
1.yes	1,458
.d:dk	181
.m:missing	17
.q:not asked this wave	14,095
.r:refuse	41
Total	21,098
	raxripfcarely_s
0.no	3,057
1.yes	1,435
.d:dk	181
.h:no help recvd	264
.m:missing	17
.q:not asked this wave	14,095
.r:refuse	41
.x:no difficulty	2,008
Total	21,098
	raxriufcarely_s
0.no	4,453
1.yes	39
.d:dk	181
.h:no help recvd	264
.m:missing	17
.q:not asked this wave	14,095
.r:refuse	41
.x:no difficulty	2,008
Total	21,098

Original SHARE Variables Used

Wave 6 End of Life:

xt623d13	Who has helped with ADLII: unpaid volunteer
xt623d14	Who has helped with ADLII: professional helper

Section L. Assistance and Caregiving

xt002_	Relationship to the deceased
xt622_	Anyone helped with ADLII
xt62odno	Difficulties doing activities II: none of these
Wave 7 End of Life:	
xt623d13	Who has helped with ADLII: unpaid volunteer
xt623d14	Who has helped with ADLII: professional helper
xt002_	Relationship to the deceased
xt622_	Anyone helped with ADLII
xt62odno	Difficulties doing activities II: none of these

Instrumental Activities of Daily Living: Frequency of Caregiving			
Variable	Waves	Label	Type
raxricared_s	All	raxricared: duration r recvd any care for 9 iadls + cont final 12 months	Categ
raxricarehr_s	All	raxricarehr_s: hrs/day r recvd any care for 9 iadls + cont final 12 months	Cont

How Constructed

raxricared_s indicates how long the deceased respondent received help with instrumental activities of daily living or continence over urination or defecation because of a physical, mental, emotional or memory problem in the last year of life. The instrumental activities of daily living include preparing hot meals, shopping for groceries, using the phone, taking medications, using a map, doing work around the house or garden, managing money, leaving the house independently, and doing personal laundry. The proxy is first asked whether the respondent had difficulty with each IADL and difficulty with continence over urination or defecation, and if the respondent had difficulty with at least one activity, then the proxy is asked whether someone helped the respondent, and if so how long the respondent received help during the last twelve months of life. **raxricared_s** is coded as follows: 1.less than 1 month, 2.1 month or more but less than 3 months, 3.3 months or more but less than 6 months, 4.6 months or more but less than 1 year, 5.a full year. **raxricared_s** is coded as special missing (.x) if the respondent did not need help with any IADL activity or with continence over urination or defecation. **raxricared_s** is coded as special missing (.h) if the respondent had difficulty but did not receive any help with any IADL activity or with continence over urination or defecation. This question is combined with all ADLs in Waves 2, 3, 4, and 8 onwards and not asked at all in Wave 5, and so **raxricared_s** is assigned special missing (.q) in these waves. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

raxricarehr_s indicates the number of hours of help that were necessary during a typical day for help with any instrumental activity of daily living or continence over urination or defecation because of a physical, mental, emotional or memory problem in the last year of life. The instrumental activities of daily living include preparing hot meals, shopping for groceries, using the phone, taking medications, using a map, doing work around the house or garden, managing money, leaving the house independently, and doing personal laundry. The proxy is first asked whether the respondent had difficulty with each IADL and difficulty with continence over urination or defecation, and if the respondent had difficulty with at least one activity, then the proxy is asked whether someone helped the respondent, and if so about how many hours of help were necessary during a typical day. **raxricarehr_s** takes the number of hours reported by the proxy, ranging from 0 to 24. **raxricarehr_s** is coded as special missing (.x) if the respondent did not need help with any IADL activity or with continence over urination or defecation. **raxricarehr_s** is coded as special missing (.h) if the respondent had difficulty but did not receive any help with any IADL activity or with continence over urination or defecation. This question is combined with all ADLs in Waves 2, 3, 4, and 8 onwards and not asked at all in Wave 5, and so **raxricarehr_s** is assigned special missing (.q) in these waves. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Cross Wave Differences in SHARE EOL

In Waves 2, 3, 4, and 8 onwards, how long the respondent received help is asked for a combination of ADL and IADL activities, and so these variables are not available. In Wave 5, how long the respondent received help is asked for ADL activities only, and not at all for IADL activities, and so these variables are not available. In Waves 6 and 7, how long the respondent received help is asked separately for ADL activities and IADL activities, and so these variables are available.

Please note that 4 IADL activities (preparing hot meals, shopping for groceries, using the telephone, taking medication) are asked about in Waves 2 through 4, while 9 IADL activities and continence over urination or defecation are asked about starting in Wave 6. Difficulty with using a map, doing work around the house or garden, managing money, leaving the house independently, doing personal laundry, and continence over urination or defecation are added in Wave 6.

Differences with the Harmonized HRS End of Life

In the HRS exit interview, the proxy is asked whether the respondent received help with each of 4 IADL activities (preparing hot meals, shopping for groceries, using the telephone, taking medication) in the final three months of life, and if so, how frequently the respondent received help from each individual helper, which can be provided as days in a typical month, days per week, or everyday. These values are transformed into Harmonized variables indicating the number of days per month for each category of helper for the Harmonized HRS End of Life. However, in the SHARE end of life interview, the proxy is asked whether the respondent received help with a list of 9 IADL activities and continence over urination or defecation in the last twelve months of life, and if so, asked how long the respondent had received help with any of the activities in the last year, which has specific response categories with a maximum of one year, and are reported as is and as a total for all helpers in the Gateway Harmonized SHARE End of Life. As such, these harmonized variables would need adjustment before being comparable.

In the HRS exit interview, the proxy is only asked whether the respondent received help with the IADLs but not whether they had difficulty with the IADLs. In the SHARE end of life interview, the proxy is asked whether the respondent received help with the IADLs and whether they had difficulty with the IADLs. As such, the variables in the Harmonized HRS End of Life only employ a special missing (.h) indicating if they received no help, while the variables in the Gateway Harmonized SHARE End of Life employ a special missing (.h) as well as a special missing (.x) indicating if they had no difficulty.

Continuous Variable Distributions

Variable	N	Mean	Std.Dev	Minimum	Maximum	Total missing
raxricarehr_s	4,007	11.53	9.37	0.00	24.00	17,091

Categorical Variable Frequencies

	raxricared_s
1.less than 1 month	199
2.1 month+ but less than 3	422
3.3 months+ but less than 6	543
4.6 months+ but less than 1	612
5.a full year	2,701
.d:dk	196
.h:no help recvd	264
.m:missing	17
.q:not asked this wave	14,095
.r:refuse	41
.x:no difficulty	2,008
Total	21,098

Original SHARE Variables Used

Wave 6 End of Life:

xt622_	Anyone helped with ADLII
xt624_	Time the deceased received help with ADLII
xt625_	Hours of help with ADLII necessary during typical day
xt62odno	Difficulties doing activities II: none of these

Wave 7 End of Life:

xt622_	Anyone helped with ADLII
xt624_	Time the deceased received help with ADLII
xt625_	Hours of help with ADLII necessary during typical day
xt62odno	Difficulties doing activities II: none of these

Instrumental Activities of Daily Living: Whether Began Receiving Help within Final Year

Variable	Waves	Label	Type
raxricareb_s	All	raxricareb_s: r began recv help with 9 iadls + cont final 12 months	Categ

How Constructed

raxricareb_s indicates whether the deceased respondent began receiving help with instrumental activities of daily living or continence over urination or defecation because of a physical, mental, emotional or memory problem in the last year of life. The instrumental activities of daily living include preparing hot meals, shopping for groceries, using the phone, taking medications, using a map, doing work around the house or garden, managing money, leaving the house independently, and doing personal laundry. The proxy is first asked whether the respondent had difficulty with each IADL and difficulty with continence over urination or defecation, and if the respondent had difficulty with at least one activity, then the proxy is asked whether someone helped the respondent, and if so how long the respondent received help during the last twelve months of life. **raxricareb_s** is coded as 0 if the respondent received help for at least one IADL activity or with continence over urination or defecation and the respondent received help for the full year. **raxricareb_s** is coded as 1 if the respondent received help with at least one IADL activity or with continence over urination or defecation and the respondent received help for less than one month, one month or more but less than three months, three months or more but less than six months, or six months or more but less than a year. **raxricareb_s** is coded as special missing (.x) if the respondent did not need help with any IADL activity or with continence over urination or defecation. **raxricareb_s** is coded as special missing (.h) if the respondent had difficulty but did not receive any help with any IADL activity or with continence over urination or defecation. This question is combined with all ADLs in Waves 2, 3, 4, and 8 onwards, and not asked at all in Wave 5, and so **raxricareb_s** is assigned special missing (.q) in these waves. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Cross Wave Differences in SHARE EOL

In Waves 2, 3, 4, and 8 onwards, how long the respondent received help is asked for a combination of ADL and IADL activities, and so this variable is not available. In Wave 5, how long the respondent received help is asked for ADL activities only, and not at all for IADL activities, and so this variable is not available. In Waves 6 and 7, how long the respondent received help is asked separately for ADL activities and IADL activities, and so this variable is available.

Please note that 4 IADL activities (preparing hot meals, shopping for groceries, using the telephone, taking medication) are asked about in Waves 2 through 4, while 9 IADL activities and continence over urination or defecation are asked about starting in Wave 6. Difficulty with using a map, doing work around the house or garden, managing money, leaving the house independently, doing personal laundry, and continence over urination or defecation are added in Wave 6.

Differences with the Harmonized HRS End of Life

In the HRS exit interview, the proxy is asked whether the respondent received help with each of 4 IADL activities (preparing hot meals, shopping for groceries, using the telephone, taking medication) in the final three months of life, and if so, asked how long the respondent had needed help with each IADL, which can be provided in months, years, since age, or since year. These responses have been converted to be equivalent to approximately 1 year in this variable in the Harmonized HRS End of Life. However, in the SHARE end of life interview, the proxy is asked whether the respondent received help with a list of 9 IADL activities and continence over urination or defecation in the last twelve months of life, and if so, asked how long the respondent had received help with any of the activities in the last year, which has specific response categories with a maximum of one year. This response has been converted so that responses of a full year are coded as not having begun to need help in the last year of life in the Gateway Harmonized SHARE End of Life. As such, this variable in the Harmonized HRS End of Life captures a slightly different concept than this variable in the Gateway Harmonized SHARE End of Life because we can ascertain with some degree of certainty when the respondent began needing help based on the questions in the HRS exit interview, whereas we are assuming that the respondent began needing help prior to the last twelve months of life if the respondent received help during their entire final year in the SHARE end of life interview. Additionally, this variable in the Harmonized HRS End of Life involves requiring help with 4 instrumental activities of daily living, while this variable in the

Gateway Harmonized SHARE End of Life involves help with the same 4 instrumental activities of daily living plus 5 additional instrumental activities of daily living and continence over urination or defecation.

In the HRS exit interview, the proxy is only asked whether the respondent received help with the IADLs but not whether they had difficulty with the IADLs. In the SHARE end of life interview, the proxy is asked whether the respondent received help with the IADLs and whether they had difficulty with the IADLs. As such, the variables in the Harmonized HRS End of Life only employ a special missing (.h) indicating if they received no help, while the variables in the Gateway Harmonized SHARE End of Life employ a special missing (.h) as well as a special missing (.x) indicating if they had no difficulty.

Categorical Variable Frequencies

	raxricareb_s
0.no	2,701
1.yes	1,776
.d:dk	196
.h:no help recvd	264
.m:missing	17
.q:not asked this wave	14,095
.r:refuse	41
.x:no difficulty	2,008
Total	21,098

Original SHARE Variables Used

Wave 6 End of Life:

xt622_	Anyone helped with ADLII
xt624_	Time the deceased received help with ADLII
xt62odno	Difficulties doing activities II: none of these

Wave 7 End of Life:

xt622_	Anyone helped with ADLII
xt624_	Time the deceased received help with ADLII
xt62odno	Difficulties doing activities II: none of these

ADLs or IADLs: Received Care for ADLs or IADLs			
Variable	Waves	Label	Type
raxrcany1y	All	raxrcany1y: r recvd any care for 10 adls/iadls final 12 months	Categ
raxrcany1y_s	All	raxrcany1y_s: r recvd any care for 16 adls/iadls final 12 months	Categ

How Constructed

raxrcany1y indicates whether anyone regularly provided care to the deceased respondent for difficulty with any activities of daily living (ADLs) or with any of 4 instrumental activities of daily living (IADLs) in the last twelve months of life because of a physical, mental, emotional or memory problem, as reported by the proxy. In Waves 2 through 4, the proxy is first asked whether the respondent had difficulty with any ADLs or with any of the 4 IADLs, and if the proxy reported that the respondent had difficulty with at least one of these activities, then the proxy is asked whether anyone regularly helped the respondent with these activities in a single question. The activities of daily living include dressing, walking across a room, bathing, eating, getting in and out of bed, and using the toilet. The instrumental activities of daily living include preparing hot meals, shopping for groceries, using the telephone, and taking medication. **raxrcany1y** is coded as 0 if nobody provided care to the respondent for difficulty with any ADLs or any of the 4 IADLs, or if the respondent had no difficulty with ADLs or with any of the 4 IADLs and the proxy is not asked if anyone helped. **raxrcany1y** is coded as 1 if somebody provided care to the respondent for difficulty with ADLs or with any of the 4 IADLs. This question was not asked starting in Wave 5, so in Waves 5 and onward **raxrcany1y** is assigned special missing (.q). Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

raxrcany1y_s indicates whether anyone regularly provided care to the deceased respondent for difficulty with any ADLs or with any of 9 IADLs or with continence over urination or defecation, in the last twelve months of life because of a physical, mental, emotional or memory problem, as reported by the proxy. In Waves 6 and 7, the proxy is asked whether the respondent had difficulty with any ADLs, and if so, whether anyone regularly helped the deceased respondent with ADLs. The activities of daily living include dressing, walking across a room, bathing, eating, getting in and out of bed, and using the toilet. Then the proxy is asked whether the respondent had difficulty with any of 9 IADLs or with continence over urination or defecation, and if so, whether anyone regularly helped the deceased respondent with these activities. The instrumental activities of daily living include preparing hot meals, shopping for groceries, using the telephone, taking medication, using a map, doing work around the house or garden, managing money, leaving the house independently, and doing personal laundry. In Wave 8 onwards, instead the proxy is asked whether anyone helped the respondent with a combination of ADLs and 9 IADLs and with continence over urination or defecation. In Waves 6 and 7, **raxrcany1y_s** summarizes whether the respondent received care for difficulty with either ADLs (**raxracany1y_s**) or IADLs and continence over urination or defecation (**raxricany1y_s**). In Wave 8 onwards, **raxrcany1y_s** indicates whether the respondent received care for difficulty with ADLs or IADLs and continence over urination and defecation. **raxrcany1y_s** is coded as 0 if nobody provided care to the respondent for difficulty with any of the activities, or if the respondent had no difficulty with any of the activities and the proxy is not asked if anyone helped. **raxrcany1y_s** is coded as 1 if somebody provided care to the respondent for difficulty with any of the activities. This question was not asked in Waves 2 through 5, so in these waves **raxrcany1y_s** is assigned special missing (.q). Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Cross Wave Differences in SHARE EOL

In Waves 2 through 4, the proxy is asked about difficulty with 6 ADLs and 4 IADLs (preparing hot meals, shopping for groceries, using the telephone, taking medication) in the last 12 months of life in a single question. If the proxy reports that the respondent had difficulty with any one of the ADLs or IADLs, then the proxy is asked whether anyone regularly helped the respondent with these activities. This response is recorded in **raxrcany1y** in these waves.

In Wave 5, difficulty with ADLs is asked about, but any difficulty with IADLs is not asked about, so neither **raxrcany1y** nor **raxrcany1y_s** are available in this wave.

In Waves 6 and 7, difficulty with and help received for ADLs is asked about, and difficulty with and help received for 9 IADLs and continence over urination or defecation are asked about, but in separate questions. In Wave 8 onwards, difficulty with ADLs and difficulty with 9 IADLs and continence over urination or defecation are asked about separately, but help with ADLs

and IADLs are asked about together. Because **raxrcany1y** indicates any difficulty with 6 ADLs and 4 IADLs, **raxrcany1y** is assigned special missing (.q) starting in Wave 5. **raxrcany1y_s**, which indicates difficulty with 6 ADLs and 9 IADLs and continence over urination or defecation, is available starting in Wave 6 when difficulty with 9 IADLs and continence over urination or defecation are inquired about. As such, **raxrcany1y_s** is assigned special missing (.q) in Waves 2 through 5. Because ADLs and IADLs are asked about separately in Waves 6 and 7, **raxrcany1y_s** summarizes whether someone helped with either ADLs or IADLs. In Wave 8 onwards, when help with ADLs and IADLs are asked about together, **raxrcany1y_s** indicates whether someone help with either ADLs or IADLs.

Differences with the Harmonized HRS End of Life

In the Harmonized HRS End of Life, this variable indicates whether anyone helped the respondent with any ADLs or any of four IADLs (preparing hot meals, shopping for groceries, using the telephone, taking medication) in the last three months of life. In the HRS exit interview, the proxy is asked whether anyone helped the respondent with each ADL and IADL individually and this variable summarizes these responses. In Waves 2 through 4 of the Gateway Harmonized SHARE End of Life, this variable is taken from a single question asking whether anyone helped the respondent with 6 ADLs or 4 IADLs in the last 12 months of life, which is asked after questions ascertaining whether the respondent had any difficulty with these activities. Because this variable in the Harmonized HRS End of Life captures a similar concept but within the final three months, while the Gateway Harmonized SHARE End of Life is within the final 12 months, the "1Y" is added to the SHARE variable names in Waves 2 through 4 to denote the difference in time period.

In Wave 5, no IADLs are asked about, and so neither **raxrcany1y** nor **raxrcany1y_s** are available in this wave. Whereas, this variable is available in all waves of the Harmonized HRS End of Life.

In Waves 6 and 7 of the Gateway Harmonized SHARE End of Life, **raxrcany1y_s** is taken from two questions, the first asking whether anyone helped the respondent with 6 ADLs, and the second asking whether anyone helped the respondent with 9 IADLs or with continence over urination or defecation in the last 12 months of life, which are asked after questions ascertaining whether the respondent had any difficulty with the respective activities. In Wave 8 onwards, help received for difficulty with ADLs or IADLs is asked in a single question but uses the same ADLs and IADLs as in Waves 6 and 7. Because this variable in the Harmonized HRS End of Life captures a similar concept but with 4 IADLs in the final three months, while the Gateway Harmonized SHARE End of Life has 9 IADLs and continence over urination or defecation in the final 12 months, the "1Y_S" is added to the SHARE variable names starting in Wave 6 to denote the difference in time period and number of activities.

In the HRS exit interview, the proxy is only asked whether the respondent received help with the ADLs or IADLs but not whether they had difficulty with the ADLs or IADLs. In the SHARE end of life interview, the proxy is asked whether the respondent received help with the ADLs or IADLs and whether they had difficulty with the ADLs or IADLs. As such, the Gateway Harmonized SHARE End of Life variables are coded as 0 if they had no difficulty.

Categorical Variable Frequencies

	raxrcany1y
0.no	1,073
1.yes	1,901
.d:dk	52
.m:missing	32
.q:not asked this wave	18,031
.r:refuse	9
Total	21,098
	raxrcany1y_s
0.no	4,758
1.yes	10,690
.d:dk	271
.m:missing	64

.q:not asked this wave	5,254
.r:refuse	61
Total	21,098

Original SHARE Variables Used

Wave 2 End of Life:	
xto2odno	Difficulties doing activities: none of these
xto22_	Anyone helped with ADL
Wave 3 End of Life:	
sl_xto2odno	Difficulties: none of these
sl_xto22_	Anyone helped with adl
Wave 4 End of Life:	
xto2odno	Difficulties doing activities: none of these
xto22_	Anyone helped with ADL
Wave 6 End of Life:	
xto22_	Anyone helped with ADL
xto2odno	Difficulties doing activities: none of these
xt622_	Anyone helped with ADLII
xt62odno	Difficulties doing activities II: none of these
Wave 7 End of Life:	
xto22_	Anyone helped with ADL
xto2odno	Difficulties doing activities: none of these
xt622_	Anyone helped with ADLII
xt62odno	Difficulties doing activities II: none of these
Wave 8 End of Life:	
xto22_	Anyone helped with ADL
xto2odno	Difficulties doing activities: none of these
xt62odno	Difficulties doing activities II: none of these
Wave 9 End of Life:	
xto22_	Anyone helped with ADL
xto2odno	Difficulties doing activities: none of these
xt62odno	Difficulties doing activities II: none of these
Wave 9 COVID End of Life:	
xto22_	Anyone helped with ADL
xto2odno	Difficulties doing activities: none of these
xt62odno	Difficulties doing activities II: none of these

ADLs or IADLs: Informal Caregiving			
Variable	Waves	Label	Type
raxrcaany1y	All	raxrcaany1y: r recvd informal care for 10 adls/iadls final 12 months	Categ
raxrscare1y	All	raxrscare1y: r recvd informal care from spouse for 10 adls/iadls final 12 months	Categ
raxrccare1y	All	raxrccare1y: r recvd informal care from children/grandchildren for 10 adls/iadls	Categ
raxrrcare1y	All	raxrrcare1y: r recvd informal care from relatives for 10 adls/iadls final 12 mon	Categ
raxrfcare1y	All	raxrfcare1y: r recvd informal care from non-relatives for 10 adls/iadls final 12	Categ
raxrcaany1y_s	All	raxrcaany1y_s: r recvd informal care for 16 adls/iadls final 12 months	Categ
raxrscare1y_s	All	raxrscare1y_s: r recvd informal care from spouse for 16 adls/iadls final 12 mont	Categ
raxrccare1y_s	All	raxrccare1y_s: r recvd informal care from children/grandchildren for 16 adls/iad	Categ
raxrrcare1y_s	All	raxrrcare1y_s: r recvd informal care from relatives for 16 adls/iadls final 12 m	Categ
raxrfcare1y_s	All	raxrfcare1y_s: r recvd informal care from non-relatives for 16 adls/iadls final	Categ

How Constructed

raxrcaany1y indicates whether the respondent received any informal care for difficulty with any activities of daily living (ADLs) or 4 instrumental activities of daily living (IADLs) in the last twelve months of life, as reported by the proxy. In Waves 2 through 4, if the proxy reports that someone helped the respondent with any ADLs or 4 IADLs, then the proxy is asked for the relationship of the top three people who helped the respondent with any of these activities in a single question. The activities of daily living include dressing, walking across a room, bathing, eating, getting in and out of bed, and using the toilet. The instrumental activities of daily living include preparing hot meals, shopping for groceries, using the telephone, and taking medication. **raxrcaany1y** is coded as 1 if the respondent's spouse, children, grandchildren, relatives, or non-relatives provided any informal care to the respondent. **raxrcaany1y** is coded as 0 if the respondent's spouse, children, grandchildren, relatives, or non-relatives did not provide any informal care to the respondent. **raxrcaany1y** is also coded as 0 if nobody helped the respondent with any activity in the last twelve months of life, or if the respondent had no difficulty with these activities and the proxy is not asked if anyone helped. In Waves 2 through 4, these variables are taken from a single question, but starting in Wave 5 the question is no longer asked the same way and **raxrcaany1y** is assigned special missing (.q). Don't know, refused, and other missing responses are assigned special missing (.d), (.r), (.m), respectively.

raxrscare1y, **raxrccare1y**, **raxrrcare1y**, and **raxrfcare1y** indicate whether a specific person provided informal care to the deceased respondent for difficulty with any activities of daily living (ADLs) or 4 instrumental activities of daily living (IADLs), as reported by the proxy. In Waves 2 through 4, if the proxy reports that someone helped the respondent with any ADLs or 4 IADLs, then the proxy is asked for the relationship of the top three people who helped the respondent with any of these activities in a single question. The activities of daily living include dressing, walking across a room, bathing, eating, getting in and out of bed, and using the toilet. The instrumental activities of daily living include preparing hot meals, shopping for groceries, using the telephone, and taking medication. **raxrscare1y** indicates whether the respondent's spouse or partner, or if the proxy, who is the respondent's husband/wife/partner, provided informal care to the respondent

for difficulty with ADLs or IADLs. **raxrccarely** indicates whether the respondent's son, son-in-law, daughter, daughter-in-law, grandson, granddaughter, or if the proxy, who is the respondent's son or daughter, son- or daughter-in-law, son or daughter of husband/wife/partner, or grandchild, provided informal care to the respondent for difficulty with ADLs or IADLs. **raxrrcarely** indicates whether the respondent's mother or father, sister, brother, other relative, or if the proxy, who is the respondent's sibling or other relative, provided informal care to the respondent for difficulty with ADLs or IADLs. **raxrfcarely** indicates whether a friend or neighbor of the deceased, other person, or if the proxy, who is the respondent's other non-relative, provided informal care to the respondent for difficulty with ADLs or IADLs.

raxrscarely, **raxrccarely**, **raxrrcarely**, and **raxrfcarely** are coded as 0 if no one with the specified relationship provided informal care to the respondent for difficulty with ADLs or IADLs. **raxrscarely**, **raxrccarely**, **raxrrcarely**, and **raxrfcarely** are coded as 1 if somebody with the specified relationship provided informal care to the respondent for difficulty with ADLs or IADLs. If the respondent had no difficulty with these activities and the proxy is not asked if anyone helped, then these variables are coded as special missing (.x). If the respondent had difficulty but did not receive help with these activities, then these variables are coded as special missing (.h). In Waves 2 through 4, these variables are taken from a single question, but starting in Wave 5 the question is no longer asked the same way and **raxrscarely**, **raxrccarely**, **raxrrcarely**, and **raxrfcarely** are assigned special missing (.q). Don't know, refused, and other missing responses are assigned special missing (.d), (.r), (.m), respectively.

raxrcaanyly_s indicates whether the respondent received any informal care for any activities of daily living (ADLs), 9 instrumental activities of daily living (IADLs), or with continence over urination or defecation in the last twelve months of life, as reported by the proxy. In Waves 6 and 7, if the proxy reports that someone helped the respondent with any ADLs, then the proxy is asked for the relationship of the top three people who helped the respondent with the ADLs in one question. The activities of daily living include dressing, walking across a room, bathing, eating, getting in and out of bed, and using the toilet. Also in Waves 6 and 7, if the proxy reports that someone helped the respondent with any of 9 IADLs or with continence over urination or defecation, then the proxy is asked for the relationship of the top three people who helped the respondent with those activities in a second question. The instrumental activities of daily living include preparing hot meals, shopping for groceries, using the telephone, taking medication, using a map, doing work around the house or garden, managing money, leaving the house independently, and doing personal laundry. **raxrcaanyly_s** summarizes whether the respondent received informal care for difficulty with either ADLs (**raxracaanyly_s**) or IADLs and continence over urination or defecation (**raxricaanyly_s**). In Wave 8 onwards, if the proxy reports that someone helped the respondent with any ADLs, IADLs, or with continence over urination or defecation, then the proxy is asked for the relationship of the top three people who helped the respondent with any of these activities in one question. **raxrcaanyly_s** directly reports whether the respondent received informal care for difficulty with either ADLs or IADLs and continence over urination or defecation. **raxrcaanyly_s** is coded as 1 if the respondent's spouse, children, grandchildren, relatives, or non-relatives provided any informal care to the respondent. **raxrcaanyly_s** is coded as 0 if the respondent's spouse, children, grandchildren, relatives, or non-relatives did not provide any informal care to the respondent. **raxrcaanyly_s** is also coded as 0 if nobody helped the respondent with any activity in the last twelve months of life, or if the respondent had no difficulty with these activities and the proxy is not asked if anyone helped. In Waves 2 through 5, these questions are not asked separately and inquire about only 4 IADL activities, so **raxrcaanyly_s** is assigned special missing (.q). Don't know, refused, and other missing responses are assigned special missing (.d), (.r), (.m), respectively.

raxrscarely_s, **raxrccarely_s**, **raxrrcarely_s**, and **raxrfcarely_s** indicate whether a specific person provided informal care to the deceased respondent for difficulty with any ADLs or 9 IADLs, or with continence over urination or defecation, as reported by the proxy. In Waves 6 and 7, if the proxy reports that someone helped the respondent with any ADLs, then the proxy is asked for the relationship of the top three people who helped the respondent with the ADLs in one question. The activities of daily living include dressing, walking across a room, bathing, eating, getting in and out of bed, and using the toilet. Also in Waves 6 and 7, if the proxy reports that someone helped the respondent with any of 9 IADLs or with continence over urination or defecation, then the proxy is asked for the relationship of the top three people who helped the respondent with those activities in a second question. The instrumental activities of daily living include preparing hot meals, shopping for groceries, using the telephone, taking medication, using a map, doing work around the house or garden, managing money, leaving the house independently, and doing personal laundry. In Wave 8 onwards, if the proxy reports that someone helped the respondent with any ADLs, IADLs, or with continence over urination or defecation, then the proxy is asked for the relationship of the top three people who helped the respondent with any of these activities in one question. These variables either summarize or directly report whether a person of the specified relationship provided informal care for difficulty with either set of activities. **raxrscarely_s** indicates whether the respondent's spouse or partner, or if the proxy,

who is the respondent's husband/wife/partner, provided informal care to the respondent for difficulty with ADLs or IADLs. **raxrccarely_s** indicates whether the respondent's son, son-in-law, daughter, daughter-in-law, grandson, granddaughter, or if the proxy, who is the respondent's son or daughter, son- or daughter-in-law, son or daughter of husband/wife/partner, or grandchild, provided informal care to the respondent for difficulty with ADLs or IADLs. **raxrrcarely_s** indicates whether the respondent's mother or father, sister, brother, other relative, or if the proxy, who is the respondent's sibling or other relative, provided informal care to the respondent for difficulty with ADLs or IADLs. **raxrfcarely_s** indicates whether a friend or neighbor of the deceased, other person, or if the proxy, who is the respondent's other non-relative, provided informal care to the respondent with ADLs or IADLs.

raxrscarely_s, **raxrccarely_s**, **raxrrcarely_s**, and **raxrfcarely_s** are coded as 0 if no one with the specified relationship provided care to the respondent for difficulty with ADLs or IADLs. **raxrscarely_s**, **raxrccarely_s**, **raxrrcarely_s**, and **raxrfcarely_s** are coded as 1 if somebody with the specified relationship provided informal care to the respondent for difficulty with ADLs or IADLs. If the respondent had no difficulty with these activities and the proxy is not asked if anyone helped, then these variables are coded as special missing (.x). If the respondent had difficulty but did not receive any help with these activities, then these variables are coded as special missing (.h). In Waves 2 through 5, these questions are not asked because they inquire about only 4 IADL activities and these variables are assigned special missing (.q). Don't know, refused, and other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Cross Wave Differences in SHARE EOL

In Waves 2 through 4, the proxy is asked about difficulty with 6 ADLs and 4 IADLs (preparing hot meals, shopping for groceries, using the telephone, taking medication) in the last 12 months of life in a single question. If the proxy reports that the respondent had difficulty with any one of the ADLs or IADLs, then the proxy is asked whether anyone regularly helped the respondent with these activities. If someone regularly helped with these activities, the proxy is asked for the respondent's relationship to the top three helpers. This response is recorded in **raxrcaany1y**, **raxrscarely**, **raxrccarely**, **raxrrcarely**, and **raxrfcarely** in these waves.

In Wave 5, difficulty with ADLs is asked about, but difficulty with IADLs is not asked about. Because no IADLs are asked about in Wave 5, neither **raxrcaany1y**, **raxrscarely**, **raxrccarely**, **raxrrcarely**, and **raxrfcarely**, nor **raxrscarely_s**, **raxrccarely_s**, **raxrrcarely_s**, and **raxrfcarely_s** are available in this wave.

In Waves 6 and 7, difficulty with and help received for ADLs is asked about, and difficulty with and help received for 9 IADLs and continence over urination or defecation are asked about, but in separate questions. In Wave 8 and onwards, difficulty with ADLs and difficulty with 9 IADLs and continence over urination or defecation are asked about separately, but help received for either ADLs or IADLs are asked about together. Because **raxrcaany1y**, **raxrscarely**, **raxrccarely**, **raxrrcarely**, and **raxrfcarely** indicate who helped the respondent with 6 ADLs and 4 IADLs, these variables are assigned special missing (.q) starting in Wave 5. **raxrcaany1y_s**, **raxrscarely_s**, **raxrccarely_s**, **raxrrcarely_s**, and **raxrfcarely_s**, which indicate who helped the respondent with 6 ADLs and 9 IADLs and continence over urination or defecation, is available starting in Wave 6 when difficulty with 9 IADLs or with continence over urination or defecation are inquired about. As such, **raxrcaany1y_s**, **raxrscarely_s**, **raxrccarely_s**, **raxrrcarely_s**, and **raxrfcarely_s** are assigned special missing (.q) in Waves 2 through 5. Because help for ADLs and IADLs are asked about separately in Waves 6 and 7, **raxrcaany1y_s**, **raxrscarely_s**, **raxrccarely_s**, **raxrrcarely_s**, and **raxrfcarely_s** summarize whether someone helped with either ADLs or IADLs. Because help for ADLs and IADLs are asked about together in Wave 8 onwards, **raxrcaany1y_s**, **raxrscarely_s**, **raxrccarely_s**, **raxrrcarely_s**, and **raxrfcarely_s** directly report whether someone helped with either ADLs or IADLs.

Differences with the Harmonized HRS End of Life

In the Harmonized HRS End of Life, these variables indicate whether anyone helped the respondent with any ADLs or any of four IADLs (preparing hot meals, shopping for groceries, using the telephone, taking medication) in the last three months of life. In the HRS exit interview, the proxy is asked whether anyone helped the respondent with each ADL and IADL individually, and then asked for the relationship of the person who helped with ADLs separately from IADLs, and these variables summarize these responses. In the HRS exit interview, proxies can respond with the relationship of up to 7 people who help with ADLs and 6 people who help with IADLs. In Waves 2 through 4 of the Gateway Harmonized SHARE End of Life, **raxrcaany1y**,

raxrscare1y, **raxrccare1y**, **raxrrcare1y**, and **raxrfcare1y** are taken from a single question asking for the relationship of the top three people who helped the respondent with 6 ADLs or 4 IADLs in the last 12 months of life, which is asked after questions ascertaining whether the respondent had any difficulty with these activities and whether someone regularly helped them with the activities. Because these variables in the Harmonized HRS End of Life capture similar concepts but within the final three months, while the Gateway Harmonized SHARE End of Life is within the final 12 months, the "1Y" is added to the SHARE variable names in Waves 2 through 4 to denote the difference in time period.

In Wave 5, no IADLs are asked about, and so neither **raxrcaany1y**, **raxrscare1y**, **raxrccare1y**, **raxrrcare1y**, and **raxrfcare1y**, nor **raxrcaany1y_s**, **raxrscare1y_s**, **raxrccare1y_s**, **raxrrcare1y_s**, and **raxrfcare1y_s** are available in this wave for SHARE. These variables are available in all harmonized waves in the Harmonized HRS End of Life.

In Waves 6 and 7 of the Gateway Harmonized SHARE End of Life, **raxrcaany1y_s**, **raxrscare1y_s**, **raxrccare1y_s**, **raxrrcare1y_s**, and **raxrfcare1y_s** are taken from two questions, the first asking the relationship of the top three people who helped the respondent with 6 ADLs, and the second asking the relationship of the top three people who helped the respondent with 9 IADLs or with continence over urination or defecation in the last 12 months of life, which are asked after questions ascertaining whether the respondent had any difficulty with the respective activities and whether someone regularly helped them with the activities. In Wave 8 onwards, help received for difficulty with ADLs or IADLs is asked in a single question but uses the same ADLs and IADLs as in Waves 6 and 7. Because these variables in the Harmonized HRS End of Life capture a similar concept but with 4 IADLs in the final three months, while the Gateway Harmonized SHARE End of Life has 9 IADLs and continence over urination or defecation in the final 12 months, the "1Y_S" is added to the SHARE variable names starting in Wave 6 to denote the difference in time period and number of activities.

In the HRS exit interview, the proxy is only asked whether the respondent received help with the ADLs or IADLs but not whether they had difficulty with the ADLs or IADLs. In the SHARE end of life interview, the proxy is asked whether the respondent received help with the ADLs or IADLs and whether they had difficulty with the ADLs or IADLs. As such, the variables in the Harmonized HRS End of Life only employ a special missing (.h) indicating if they received no help, while the variables in the Gateway Harmonized SHARE End of Life employ a special missing (.h) as well as a special missing (.x) in some variables indicating if they had no difficulty.

Categorical Variable Frequencies

	raxrcaany1y
0.no	1,314
1.yes	1,659
.d:dk	53
.m:missing	32
.q:not asked this wave	18,031
.r:refuse	9
Total	21,098
	raxrscare1y
0.no	1,120
1.yes	780
.d:dk	53
.h:no help recvd	82
.m:missing	32
.q:not asked this wave	18,031
.r:refuse	9
.x:no difficulty	991
Total	21,098
	raxrccare1y
0.no	870
1.yes	1,030

.d:dk	53
.h:no help recvd	82
.m:missing	32
.q:not asked this wave	18,031
.r:refuse	9
.x:no difficulty	991
Total	21,098
raxrrcare1y	
o.no	1,745
1.yes	155
.d:dk	53
.h:no help recvd	82
.m:missing	32
.q:not asked this wave	18,031
.r:refuse	9
.x:no difficulty	991
Total	21,098
raxrfcare1y	
o.no	1,711
1.yes	189
.d:dk	53
.h:no help recvd	82
.m:missing	32
.q:not asked this wave	18,031
.r:refuse	9
.x:no difficulty	991
Total	21,098
raxrcaany1y_s	
o.no	5,760
1.yes	9,671
.d:dk	272
.m:missing	77
.q:not asked this wave	5,254
.r:refuse	64
Total	21,098
raxrscare1y_s	
o.no	6,246
1.yes	4,427
.d:dk	272
.h:no help recvd	766
.m:missing	77
.q:not asked this wave	5,254
.r:refuse	64
.x:no difficulty	3,992
Total	21,098
raxrccare1y_s	
o.no	4,539
1.yes	6,134

.d:dk	272
.h:no help recvd	766
.m:missing	77
.q:not asked this wave	5,254
.r:refuse	64
.x:no difficulty	3,992
Total	21,098
raxrrcarely_s	
o.no	9,748
1.yes	925
.d:dk	272
.h:no help recvd	766
.m:missing	77
.q:not asked this wave	5,254
.r:refuse	64
.x:no difficulty	3,992
Total	21,098
raxrfcarely_s	
o.no	9,496
1.yes	1,177
.d:dk	272
.h:no help recvd	766
.m:missing	77
.q:not asked this wave	5,254
.r:refuse	64
.x:no difficulty	3,992
Total	21,098

Original SHARE Variables Used

Wave 2 End of Life:

xto2odno	Difficulties doing activities: none of these
xto22_	Anyone helped with ADL
xto02_	Relationship to the deceased
xto23d1	Who has helped with ADL: yourself (proxy respondent)
xto23d2	Who has helped with ADL: spouse/partner of deceased
xto23d3	Who has helped with ADL: mother/father of deceased
xto23d4	Who has helped with ADL: son of deceased
xto23d5	Who has helped with ADL: son-in-law of deceased
xto23d6	Who has helped with ADL: daughter of deceased
xto23d7	Who has helped with ADL: daughter-in-law of deceased
xto23d8	Who has helped with ADL: grandson of deceased
xto23d9	Who has helped with ADL: granddaughter of deceased
xto23d10	Who has helped with ADL: sister of deceased
xto23d11	Who has helped with ADL: brother of deceased
xto23d12	Who has helped with ADL: other relative
xto23d15	Who has helped with ADL: friend/neighbor of deceased
xto23dot	Who has helped with ADL: other person

Wave 3 End of Life:

sl_xto2odno	Difficulties: none of these
-------------	-----------------------------

sl_xt022_	Anyone helped with adl
sl_xt002_	Relationship to the deceased
sl_xt023d1	Who has helped with adl: proxy respondent
sl_xt023d2	Who has helped with adl: husband/wife/partner of the deceased
sl_xt023d3	Who has helped with adl: mother/father of the deceased
sl_xt023d4	Who has helped with adl: son of the deceased
sl_xt023d5	Who has helped with adl: son-in-law of the deceased
sl_xt023d6	Who has helped with adl: daughter of the deceased
sl_xt023d7	Who has helped with adl: daughter-in-law of the deceased
sl_xt023d8	Who has helped with adl: grandson of the deceased
sl_xt023d9	Who has helped with adl: granddaughter of the deceased
sl_xt023d10	Who has helped with adl: sister of the deceased
sl_xt023d11	Who has helped with adl: brother of the deceased
sl_xt023d12	Who has helped with adl: other relative
sl_xt023d15	Who has helped with adl: friend/neighbor of the deceased
sl_xt023dot	Who has helped with adl: other person
Wave 4 End of Life:	
xt020dno	Difficulties doing activities: none of these
xt022_	Anyone helped with ADL
xt002_	Relationship to the deceased
xt023d1	Who has helped with ADL: yourself (proxy respondent)
xt023d2	Who has helped with ADL: spouse/partner of deceased
xt023d3	Who has helped with ADL: mother/father of deceased
xt023d4	Who has helped with ADL: son of deceased
xt023d5	Who has helped with ADL: son-in-law of deceased
xt023d6	Who has helped with ADL: daughter of deceased
xt023d7	Who has helped with ADL: daughter-in-law of deceased
xt023d8	Who has helped with ADL: grandson of deceased
xt023d9	Who has helped with ADL: granddaughter of deceased
xt023d10	Who has helped with ADL: sister of deceased
xt023d11	Who has helped with ADL: brother of deceased
xt023d12	Who has helped with ADL: other relative
xt023d15	Who has helped with ADL: friend/neighbor of deceased
xt023dot	Who has helped with ADL: other person
Wave 6 End of Life:	
xt023d1	Who has helped with ADL: yourself (proxy respondent)
xt023d2	Who has helped with ADL: spouse/partner of deceased
xt023d3	Who has helped with ADL: mother/father of deceased
xt023d4	Who has helped with ADL: son of deceased
xt023d5	Who has helped with ADL: son-in-law of deceased
xt023d6	Who has helped with ADL: daughter of deceased
xt023d7	Who has helped with ADL: daughter-in-law of deceased
xt023d8	Who has helped with ADL: grandson of deceased
xt023d9	Who has helped with ADL: granddaughter of deceased
xt023d10	Who has helped with ADL: sister of deceased
xt023d11	Who has helped with ADL: brother of deceased
xt023d12	Who has helped with ADL: other relative
xt023d15	Who has helped with ADL: friend/neighbor of deceased
xt023dot	Who has helped with ADL: other person
xt002_	Relationship to the deceased
xt022_	Anyone helped with ADL
xt020dno	Difficulties doing activities: none of these
xt623d1	Who has helped with ADLII: yourself (proxy respondent)

xt623d2	Who has helped with ADLII: spouse/partner of deceased
xt623d3	Who has helped with ADLII: mother/father of deceased
xt623d4	Who has helped with ADLII: son of deceased
xt623d5	Who has helped with ADLII: son-in-law of deceased
xt623d6	Who has helped with ADLII: daughter of deceased
xt623d7	Who has helped with ADLII: daughter-in-law of deceased
xt623d8	Who has helped with ADLII: grandson of deceased
xt623d9	Who has helped with ADLII: granddaughter of deceased
xt623d10	Who has helped with ADLII: sister of deceased
xt623d11	Who has helped with ADLII: brother of deceased
xt623d12	Who has helped with ADLII: other relative
xt623d15	Who has helped with ADLII: friend/neighbor of deceased
xt623dot	Who has helped with ADLII: other person
xt622_	Anyone helped with ADLII
xt62odno	Difficulties doing activities II: none of these

Wave 7 End of Life:

xt023d1	Who has helped with ADL: yourself (proxy respondent)
xt023d2	Who has helped with ADL: spouse/partner of deceased
xt023d3	Who has helped with ADL: mother/father of deceased
xt023d4	Who has helped with ADL: son of deceased
xt023d5	Who has helped with ADL: son-in-law of deceased
xt023d6	Who has helped with ADL: daughter of deceased
xt023d7	Who has helped with ADL: daughter-in-law of deceased
xt023d8	Who has helped with ADL: grandson of deceased
xt023d9	Who has helped with ADL: granddaughter of deceased
xt023d10	Who has helped with ADL: sister of deceased
xt023d11	Who has helped with ADL: brother of deceased
xt023d12	Who has helped with ADL: other relative
xt023d15	Who has helped with ADL: friend/neighbor of deceased
xt023dot	Who has helped with ADL: other person
xt002_	Relationship to the deceased
xt022_	Anyone helped with ADL
xt02odno	Difficulties doing activities: none of these
xt623d1	Who has helped with ADLII: yourself (proxy respondent)
xt623d2	Who has helped with ADLII: spouse/partner of deceased
xt623d3	Who has helped with ADLII: mother/father of deceased
xt623d4	Who has helped with ADLII: son of deceased
xt623d5	Who has helped with ADLII: son-in-law of deceased
xt623d6	Who has helped with ADLII: daughter of deceased
xt623d7	Who has helped with ADLII: daughter-in-law of deceased
xt623d8	Who has helped with ADLII: grandson of deceased
xt623d9	Who has helped with ADLII: granddaughter of deceased
xt623d10	Who has helped with ADLII: sister of deceased
xt623d11	Who has helped with ADLII: brother of deceased
xt623d12	Who has helped with ADLII: other relative
xt623d15	Who has helped with ADLII: friend/neighbor of deceased
xt623dot	Who has helped with ADLII: other person
xt622_	Anyone helped with ADLII
xt62odno	Difficulties doing activities II: none of these

Wave 8 End of Life:

xt023d1	Who has helped with ADL: yourself (proxy respondent)
xt023d2	Who has helped with ADL: spouse/partner of deceased
xt023d3	Who has helped with ADL: mother/father of deceased

xto23d4	Who has helped with ADL: son of deceased
xto23d5	Who has helped with ADL: son-in-law of deceased
xto23d6	Who has helped with ADL: daughter of deceased
xto23d7	Who has helped with ADL: daughter-in-law of deceased
xto23d8	Who has helped with ADL: grandson of deceased
xto23d9	Who has helped with ADL: granddaughter of deceased
xto23d10	Who has helped with ADL: sister of deceased
xto23d11	Who has helped with ADL: brother of deceased
xto23d12	Who has helped with ADL: other relative
xto23d15	Who has helped with ADL: friend/neighbor of deceased
xto23dot	Who has helped with ADL: other person
xto02_	Relationship to the deceased
xto22_	Anyone helped with ADL
xto2odno	Difficulties doing activities: none of these
xt62odno	Difficulties doing activities II: none of these

Wave 9 End of Life:

xto23d1	Who has helped with ADL: yourself (proxy respondent)
xto23d2	Who has helped with ADL: spouse/partner of deceased
xto23d3	Who has helped with ADL: mother/father of deceased
xto23d4	Who has helped with ADL: son of deceased
xto23d5	Who has helped with ADL: son-in-law of deceased
xto23d6	Who has helped with ADL: daughter of deceased
xto23d7	Who has helped with ADL: daughter-in-law of deceased
xto23d8	Who has helped with ADL: grandson of deceased
xto23d9	Who has helped with ADL: granddaughter of deceased
xto23d10	Who has helped with ADL: sister of deceased
xto23d11	Who has helped with ADL: brother of deceased
xto23d12	Who has helped with ADL: other relative
xto23d15	Who has helped with ADL: friend/neighbor of deceased
xto23dot	Who has helped with ADL: other person
xto02_	Relationship to the deceased
xto22_	Anyone helped with ADL
xto2odno	Difficulties doing activities: none of these
xt62odno	Difficulties doing activities II: none of these

Wave 9 COVID End of Life:

xto23d1	Who has helped with ADL: yourself (proxy respondent)
xto23d2	Who has helped with ADL: spouse/partner of deceased
xto23d3	Who has helped with ADL: mother/father of deceased
xto23d4	Who has helped with ADL: son of deceased
xto23d5	Who has helped with ADL: son-in-law of deceased
xto23d6	Who has helped with ADL: daughter of deceased
xto23d7	Who has helped with ADL: daughter-in-law of deceased
xto23d8	Who has helped with ADL: grandson of deceased
xto23d9	Who has helped with ADL: granddaughter of deceased
xto23d10	Who has helped with ADL: sister of deceased
xto23d11	Who has helped with ADL: brother of deceased
xto23d12	Who has helped with ADL: other relative
xto23d15	Who has helped with ADL: friend/neighbor of deceased
xto23dot	Who has helped with ADL: other person
xto02_	Relationship to the deceased
xto22_	Anyone helped with ADL
xto2odno	Difficulties doing activities: none of these
xt62odno	Difficulties doing activities II: none of these

ADLs or IADLs: Formal Caregiving			
Variable	Waves	Label	Type
raxrfaany1y	All	raxrfaany1y: r recvd formal care for 10 adls/iadls final 12 months	Categ
raxrpfcare1y	All	raxrpfcare1y: r recvd formal care from paid caregiver for 10 adls/iadls final 12	Categ
raxrufcare1y	All	raxrufcare1y: r recvd formal care from unpaid caregiver for 10 adls/iadls final	Categ
raxrfaany1y_s	All	raxrfaany1y_s: r recvd formal care for 16 adls/iadls final 12 months	Categ
raxrpfcare1y_s	All	raxrpfcare1y_s: r recvd formal care from paid caregiver for 16 adls/iadls final	Categ
raxrufcare1y_s	All	raxrufcare1y_s: r recvd formal care from unpaid caregiver for 16 adls/iadls fina	Categ

How Constructed

raxrfaany1y indicates whether the respondent received any formal care for any activities of daily living (ADLs) or 4 instrumental activities of daily living (IADLs) in the last twelve months of life, as reported by the proxy. In Waves 2 through 4, if the proxy reports that someone helped the respondent with any ADLs or 4 IADLs, then the proxy is asked for the relationship of the top three people who helped the respondent with any of these activities in a single question. The activities of daily living include dressing, walking across a room, bathing, eating, getting in and out of bed, and using the toilet. The instrumental activities of daily living include preparing hot meals, shopping for groceries, using the telephone, and taking medication. **raxrfaany1y** is coded as 1 if a professional helper or unpaid volunteer provided any formal care to the respondent. **raxrfaany1y** is coded as 0 if a professional helper or unpaid volunteer did not provide any formal care to the respondent. **raxrfaany1y** is also coded as 0 if nobody helped the respondent with any activity in the last twelve months of life, or if the respondent had no difficulty with these activities and the proxy is not asked if anyone helped. In Waves 2 through 4, these variables are taken from a single question, but starting in Wave 5 the question is no longer asked the same way and **raxrfaany1y** is assigned special missing (.q). Don't know, refused, and other missing responses are assigned special missing (.d), (.r), (.m), respectively.

raxrpfcare1y and **raxrufcare1y** indicate whether a specific person provided formal care to the deceased respondent for difficulty with any activities of daily living (ADLs) or 4 instrumental activities of daily living (IADLs), as reported by the proxy. In Waves 2 through 4, if the proxy reports that someone helped the respondent with any ADLs or 4 IADLs, then the proxy is asked for the relationship of the top three people who helped the respondent with any of these activities in a single question. The activities of daily living include dressing, walking across a room, bathing, eating, getting in and out of bed, and using the toilet. The instrumental activities of daily living include preparing hot meals, shopping for groceries, using the telephone, and taking medication. **raxrpfcare1y** indicates whether a professional helper (e.g. nurse) provided formal care to the respondent for difficulty with ADLs or IADLs. **raxrufcare1y** indicates whether an unpaid volunteer provided formal care to the respondent for difficulty with ADLs or IADLs. **raxrpfcare1y** and **raxrufcare1y** are coded as 0 if no one with the specified relationship provided formal care to the respondent for difficulty with ADLs or IADLs. **raxrpfcare1y** and **raxrufcare1y** are coded as 1 if somebody with the specified relationship helped the respondent with ADLs or IADLs. If the respondent had no difficulty with these activities and the proxy is not asked if anyone helped, then these variables are coded as special missing (.x). If the respondent had difficulty but did not receive help with these activities, then these variables are coded as special missing (.h). In Waves 2 through 4, these variables are taken from a single question, but starting in Wave 5 the question is no longer asked the same way and **raxrpfcare1y** and **raxrufcare1y** are assigned special missing (.q). Don't know, refused, and other missing responses are assigned special missing (.d), (.r), (.m), respectively.

raxrfaany1y_s indicates whether the respondent received any formal care for any activities of daily living (ADLs), 9 instrumental activities of daily living (IADLs), or with continence over urination or defecation in the last twelve months of life, as reported by the proxy. In Waves 6 and 7, if the proxy reports that someone helped the respondent with any ADLs, then the proxy is asked for the relationship of the top three people who helped the respondent with the ADLs in one question.

The activities of daily living include dressing, walking across a room, bathing, eating, getting in and out of bed, and using the toilet. Also in Waves 6 and 7, if the proxy reports that someone helped the respondent with any of 9 IADLs or with continence over urination or defecation, then the proxy is asked for the relationship of the top three people who helped the respondent with those activities in a second question. The instrumental activities of daily living include preparing hot meals, shopping for groceries, using the telephone, taking medication, using a map, doing work around the house or garden, managing money, leaving the house independently, and doing personal laundry. **raxrfaany1y_s** summarizes whether the respondent received formal care for difficulty with either ADLs (**raxrafaany1y_s**) or IADLs and continence over urination or defecation (**raxrifaany1y_s**). In Wave 8 onwards, if the proxy reports that someone helped the respondent with any ADLs or with 9 IADLs or continence over urination or defecation, then the proxy is asked for the relationship of the top three people who helped the respondent with any of those activities in a single question. **raxrfaany1y_s** directly reports whether the respondent received formal care for difficulty with either ADLs or IADLs and continence over urination or defecation. **raxrfaany1y_s** is coded as 1 if a professional helper or unpaid volunteer provided any formal care to the respondent. **raxrfaany1y_s** is coded as 0 if a professional helper or unpaid volunteer did not provide any formal care to the respondent. **raxrfaany1y_s** is also coded as 0 if nobody helped the respondent with any activity in the last twelve months of life, or if the respondent had no difficulty with these activities and the proxy is not asked if anyone helped. In Waves 2 through 5, these questions are not asked separately and inquire about only 4 IADL activities, so **raxrfaany1y_s** is assigned special missing (.q). Don't know, refused, and other missing responses are assigned special missing (.d), (.r), (.m), respectively.

raxrpfcare1y_s and **raxrufcare1y_s** indicate whether a specific person provided formal care to the deceased respondent for difficulty with any ADLs or 9 IADLs, or with continence over urination or defecation, as reported by the proxy. In Waves 6 and 7, if the proxy reports that someone helped the respondent with any ADLs, then the proxy is asked for the relationship of the top three people who helped the respondent with the ADLs in one question. The activities of daily living include dressing, walking across a room, bathing, eating, getting in and out of bed, and using the toilet. Also in Waves 6 and 7, if the proxy reports that someone helped the respondent with any of 9 IADLs or with continence over urination or defecation, then the proxy is asked for the relationship of the top three people who helped the respondent with those activities in a second question. The instrumental activities of daily living include preparing hot meals, shopping for groceries, using the telephone, taking medication, using a map, doing work around the house or garden, managing money, leaving the house independently, and doing personal laundry. In Wave 8 onwards, if the proxy reports that someone helped the respondent with any ADLs or with 9 IADLs or continence over urination or defecation, then the proxy is asked for the relationship of the top three people who helped the respondent with those activities in a single question. These variables either summarize or directly report whether a person of the specified relationship provided formal care for difficulty with either set of activities. **raxrpfcare1y_s** indicates whether a professional helper (e.g. nurse) provided formal care to the respondent for difficulty with ADLs or IADLs. **raxrufcare1y_s** indicates whether an unpaid volunteer provided formal care to the respondent for difficulty with ADLs or IADLs. **raxrpfcare1y_s** and **raxrufcare1y_s** are coded as 0 if no one with the specified relationship provided formal care to the respondent for difficulty with ADLs or IADLs. **raxrpfcare1y_s** and **raxrufcare1y_s** are coded as 1 if somebody with the specified relationship provided care to the respondent for difficulty with ADLs or IADLs. If the respondent had no difficulty with these activities and the proxy is not asked if anyone helped, then these variables are coded as special missing (.x). If the respondent had difficulty but did not receive help with these activities, then these variables are coded as special missing (.h). In Waves 2 through 5, these questions are not asked separately and inquire about only 4 IADL activities and these variables are assigned special missing (.q). Don't know, refused, and other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Cross Wave Differences in SHARE EOL

In Waves 2 through 4, the proxy is asked about difficulty with 6 ADLs and 4 IADLs (preparing hot meals, shopping for groceries, using the telephone, taking medication) in the last 12 months of life in a single question. If the proxy reports that the respondent had difficulty with any one of the ADLs or IADLs, then the proxy is asked whether anyone regularly helped the respondent with these activities. If someone regularly helped with these activities, the proxy is asked for the respondent's relationship to the top three helpers. This response is recorded in **raxrfaany1y**, **raxrpfcare1y**, and **raxrufcare1y** in these waves.

In Wave 5, difficulty with ADLs is asked about, but difficulty with IADLs is not asked about. Because no IADLs are asked about in Wave 5, neither **raxrfaany1y**, **raxrpfcare1y**, and **raxrufcare1y**, nor **raxrfaany1y_s**, **raxrpfcare1y_s**, and

raxrufcare1y_s are available in this wave.

In Waves 6 and 7, difficulty with and help received for ADLs is asked about, and difficulty with and help received for 9 IADLs and continence over urination or defecation are asked about, but in separate questions. In Wave 8 onwards, difficulty with ADLs is asked about, and difficulty with 9 IADLs and continence over urination or defecation are asked about separately, but help received for ADLs or IADLs are asked about together. Because **raxrfaany1y**, **raxrpfcare1y**, and **raxrufcare1y** indicate who helped the respondent with 6 ADLs and 4 IADLs, these variables are assigned special missing (.q) starting in Wave 5. **raxrfaany1y_s**, **raxrpfcare1y_s**, and **raxrufcare1y_s**, which indicate who helped the respondent with 6 ADLs and 9 IADLs and continence over urination or defecation, are available starting in Wave 6 when difficulty with 9 IADLs or with continence over urination or defecation are inquired about. As such, **raxrfaany1y_s**, **raxrpfcare1y_s**, and **raxrufcare1y_s** are assigned special missing (.q) in Waves 2 through 5. Because help received for ADLs and IADLs are asked about separately in Waves 6 and 7, **raxrfaany1y_s**, **raxrpfcare1y_s**, and **raxrufcare1y_s** summarize whether someone helped with either ADLs or IADLs. Because help received for ADLs and IADLs are asked about together in Wave 8 onwards, **raxrfaany1y_s**, **raxrpfcare1y_s**, and **raxrufcare1y_s** directly report whether someone helped with either ADLs or IADLs.

Differences with the Harmonized HRS End of Life

In the Harmonized HRS End of Life, these variables indicate whether anyone helped the respondent with any ADLs or any of four IADLs (preparing hot meals, shopping for groceries, using the telephone, taking medication) in the last three months of life. In the HRS exit interview, the proxy is asked whether anyone helped the respondent with each ADL and IADL individually, and then asked for the relationship of the person who helped with ADLs separately from IADLs, and these variables summarize these responses. In the HRS exit interview, proxies can respond with the relationship of up to 7 people who help with ADLs and 6 people who help with IADLs. In Waves 2 through 4 of the Gateway Harmonized SHARE End of Life, **raxrfaany1y**, **raxrpfcare1y**, and **raxrufcare1y** are taken from a single question asking for the relationship of the top three people who helped the respondent with 6 ADLs or 4 IADLs in the last 12 months of life, which is asked after questions ascertaining whether the respondent had any difficulty with these activities and whether someone regularly helped them with the activities. Because these variables in the Harmonized HRS End of Life capture similar concepts but within the final three months, while the Gateway Harmonized SHARE End of Life is within the final 12 months, the "1Y" is added to the SHARE variable names in Waves 2 through 4 to denote the difference in time period.

In Wave 5, no IADLs are asked about, and so neither **raxrfaany1y**, **raxrpfcare1y**, and **raxrufcare1y**, nor **raxrfaany1y_s**, **raxrpfcare1y_s**, and **raxrufcare1y_s** are available in this wave for SHARE. These variables are available in all harmonized waves in the Harmonized HRS End of Life.

In Waves 6 and 7 of the Gateway Harmonized SHARE End of Life, **raxrfaany1y_s**, **raxrpfcare1y_s**, and **raxrufcare1y_s** are taken from two questions, the first asking the relationship of the top three people who helped the respondent with 6 ADLs, and the second asking the relationship of the top three people who helped the respondent with 9 IADLs or with continence over urination or defecation in the last 12 months of life, which are asked after questions ascertaining whether the respondent had any difficulty with the respective activities and whether someone regularly helped them with the activities. In Wave 8 onwards, help received for difficulty with ADLs or IADLs is asked in a single question but uses the same ADLs and IADLs as in Waves 6 and 7. Because these variables in the Harmonized HRS End of Life capture a similar concept but with 4 IADLs in the final three months, while the Gateway Harmonized SHARE End of Life has 9 IADLs and continence over urination or defecation in the final 12 months, the "1Y_S" is added to the SHARE variable names starting in Wave 6 to denote the difference in time period and number of activities.

In the HRS exit interview, the proxy is only asked whether the respondent received help with the ADLs or IADLs but not whether they had difficulty with the ADLs or IADLs. In the SHARE end of life interview, the proxy is asked whether the respondent received help with the ADLs or IADLs and whether they had difficulty with the ADLs or IADLs. As such, the variables in the Harmonized HRS End of Life only employ a special missing (.h) indicating if they received no help, while the variables in the Gateway Harmonized SHARE End of Life employ a special missing (.h) as well as a special missing (.x) in some variables indicating if they had no difficulty.

Categorical Variable Frequencies

	raxrfaany1y	
0.no	2,217	
1.yes	756	
.d:dk	53	
.m:missing	32	
.q:not asked this wave	18,031	
.r:refuse	9	
Total	21,098	
	raxrpfcare1y	
0.no	1,153	
1.yes	747	
.d:dk	53	
.h:no help recvd	82	
.m:missing	32	
.q:not asked this wave	18,031	
.r:refuse	9	
.x:no difficulty	991	
Total	21,098	
	raxrufcare1y	
0.no	1,879	
1.yes	21	
.d:dk	53	
.h:no help recvd	82	
.m:missing	32	
.q:not asked this wave	18,031	
.r:refuse	9	
.x:no difficulty	991	
Total	21,098	
	raxrfaany1y_s	
0.no	11,708	
1.yes	3,723	
.d:dk	271	
.m:missing	78	
.q:not asked this wave	5,254	
.r:refuse	64	
Total	21,098	
	raxrpfcarely_s	
0.no	7,026	
1.yes	3,647	
.d:dk	271	
.h:no help recvd	766	
.m:missing	78	
.q:not asked this wave	5,254	
.r:refuse	64	
.x:no difficulty	3,992	
Total	21,098	
	raxrufcarely_s	
0.no	10,540	

1.yes	128
.d:dk	271
.h:no help recvd	766
.m:missing	83
.q:not asked this wave	5,254
.r:refuse	64
.x:no difficulty	3,992
Total	21,098

Original SHARE Variables Used

Wave 2 End of Life:

xt02odno	Difficulties doing activities: none of these
xt022_	Anyone helped with ADL
xt002_	Relationship to the deceased
xt023d13	Who has helped with ADL: unpaid volunteer
xt023d14	Who has helped with ADL: professional helper

Wave 3 End of Life:

sl_xt02odno	Difficulties: none of these
sl_xt022_	Anyone helped with adl
sl_xt002_	Relationship to the deceased
sl_xt023d13	Who has helped with adl: unpaid volunteer
sl_xt023d14	Who has helped with adl: professional helper (e.g. nurse)

Wave 4 End of Life:

xt02odno	Difficulties doing activities: none of these
xt022_	Anyone helped with ADL
xt002_	Relationship to the deceased
xt023d13	Who has helped with ADL: unpaid volunteer
xt023d14	Who has helped with ADL: professional helper

Wave 6 End of Life:

xt023d13	Who has helped with ADL: unpaid volunteer
xt023d14	Who has helped with ADL: professional helper
xt002_	Relationship to the deceased
xt022_	Anyone helped with ADL
xt02odno	Difficulties doing activities: none of these
xt623d13	Who has helped with ADLII: unpaid volunteer
xt623d14	Who has helped with ADLII: professional helper
xt622_	Anyone helped with ADLII
xt62odno	Difficulties doing activities II: none of these

Wave 7 End of Life:

xt023d13	Who has helped with ADL: unpaid volunteer
xt023d14	Who has helped with ADL: professional helper
xt002_	Relationship to the deceased
xt022_	Anyone helped with ADL
xt02odno	Difficulties doing activities: none of these
xt623d13	Who has helped with ADLII: unpaid volunteer
xt623d14	Who has helped with ADLII: professional helper
xt622_	Anyone helped with ADLII
xt62odno	Difficulties doing activities II: none of these

Wave 8 End of Life:

xt023d13	Who has helped with ADL: unpaid volunteer
----------	---

xt023d14	Who has helped with ADL: professional helper
xt002_	Relationship to the deceased
xt022_	Anyone helped with ADL
xt020dno	Difficulties doing activities: none of these
xt620dno	Difficulties doing activities II: none of these
Wave 9 End of Life:	
xt023d13	Who has helped with ADL: unpaid volunteer
xt023d14	Who has helped with ADL: professional helper
xt002_	Relationship to the deceased
xt022_	Anyone helped with ADL
xt020dno	Difficulties doing activities: none of these
xt620dno	Difficulties doing activities II: none of these
Wave 9 COVID End of Life:	
xt023d13	Who has helped with ADL: unpaid volunteer
xt023d14	Who has helped with ADL: professional helper
xt002_	Relationship to the deceased
xt022_	Anyone helped with ADL
xt020dno	Difficulties doing activities: none of these
xt620dno	Difficulties doing activities II: none of these

ADLs or IADLs: Frequency of Caregiving			
Variable	Waves	Label	Type
raxrcared	All	raxrcared: duration r recvd any care for 10 adls/iadls final 12 months	Categ
raxrcared_s	All	raxrcared_s: duration r recvd any care for 16 adls/iadls final 12 months	Categ
raxrcarehr	All	raxrcarehr: hrs/day r recvd any care for 10 adls/iadls final 12 months	Cont
raxrcarehr_s	All	raxrcarehr_s: hrs/day r recvd any care for 16 adls/iadls final 12 months	Cont

How Constructed

raxrcared indicates how long the deceased respondent received help with any of 6 activities of daily living (ADLs) or 4 instrumental activities of daily living (IADLs) in the last twelve months of life overall, as reported by the proxy. In Waves 2 through 4, the activities of daily living include dressing, walking across a room, bathing, eating, getting in and out of bed, and using the toilet, and the instrumental activities of daily living include preparing hot meals, shopping for groceries, using the telephone, and taking medication. In these waves, the 6 ADL activities and the 4 IADL activities are asked about in a single question. The proxy is first asked whether the respondent had difficulty with each ADL and IADL, and if the respondent had difficulty with at least one ADL or IADL, then the proxy is asked whether someone helped the respondent, and if so how long the respondent received help during the last twelve months of life. **raxrcared** is coded as follows: 1.less than 1 month, 2.1 month or more but less than 3 months, 3.3 months or more but less than 6 months, 4.6 months or more but less than 1 year, 5.a full year. **raxrcared** is coded as special missing (.x) if the respondent did not need help with any ADL or IADL activity. **raxrcared** is coded as special missing (.h) if the respondent had difficulty but did not receive any help with any ADL or IADL activity. This question is no longer asked after Wave 4, and so **raxrcared** is assigned special missing (.q) starting in Wave 5. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

raxrcared_s indicates how long the deceased respondent received help with any of 6 activities of daily living (ADLs) or 9 instrumental activities of daily living (IADLs) or continence over urination and defecation in the last twelve months of life overall, as reported by the proxy. Starting in Wave 6, the activities of daily living include dressing, walking across a room, bathing, eating, getting in and out of bed, and using the toilet, and the instrumental activities of daily living include preparing hot meals, shopping for groceries, using the telephone, taking medication, using a map, doing work around the house or garden, managing money, leaving the house independently, and doing personal laundry. In Waves 6 and 7, help received for the 6 ADL activities and the 9 IADL activities and continence over urination or defecation are asked about in separate questions, so **raxrcared_s** is assigned the maximum amount of time the respondent received help for ADL activities (**raxracared**) and for IADL activities and continence (**raxricared_s**). In Wave 8 onwards, help received for the 6 ADL activities and the 9 IADL activities and continence over urination or defecation are asked about together, so **raxrcared_s** is the direct report of the amount of time the respondent received help for ADL activities and for IADL activities and continence. The proxy is first asked whether the respondent had difficulty with each ADL and IADL, and if the respondent had difficulty with at least one ADL or IADL, then the proxy is asked whether someone helped the respondent (separately for ADLs and IADLs in Waves 6 and 7 or for either ADLs and IADLS in Wave 8 onwards), and if so how long the respondent received help during the last twelve months of life. **raxrcared_s** is coded as follows: 1.less than 1 month, 2.1 month or more but less than 3 months, 3.3 months or more but less than 6 months, 4.6 months or more but less than 1 year, 5.a full year. **raxrcared_s** is coded as special missing (.x) if the respondent did not need help with any ADL or IADL activity. **raxrcared_s** is coded as special missing (.h) if the respondent had difficulty but did not receive any help with any ADL or IADL activity. This question is asked starting in Wave 6, and so **raxrcared_s** is assigned special missing (.q) in Waves 2 through 5. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

raxrcarehr indicates the number of hours of help that were necessary during a typical day for help with any of 6 activities of daily living (ADLs) or 4 instrumental activities of daily living (IADLs) in the last twelve months of life overall, as reported by the proxy. In Waves 2 through 4, the activities of daily living include dressing, walking across a room, bathing, eating, getting in and out of bed, and using the toilet, and the instrumental activities of daily living include preparing hot meals, shopping

for groceries, using the telephone, and taking medication. In these waves, the 6 ADL activities and the 4 IADL activities are asked about in a single question. The proxy is first asked whether the respondent had difficulty with each ADL and IADL, and if the respondent had difficulty with at least one ADL or IADL, then the proxy is asked whether someone helped the respondent, and if so about how many hours of help were necessary during a typical day. **raxrcarehr** takes the number of hours reported by the proxy, ranging from 0 to 24. **raxrcarehr** is coded as 0 if the respondent needed help with an ADL or IADL activity but nobody provided help. **raxrcarehr** is coded as special missing (.x) if the respondent did not need help with any ADL or IADL activity. **raxrcarehr** is coded as special missing (.h) if the respondent had difficulty but did not receive any help with any ADL or IADL activity. This question is no longer asked after Wave 4, and so **raxrcarehr** is assigned special missing (.q) starting in Wave 5. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

raxrcarehr_s indicates the number of hours of help that were necessary during a typical day for help with any of 6 activities of daily living (ADLs) or 9 instrumental activities of daily living (IADLs) or with continence over urination or defecation in the last twelve months of life overall, as reported by the proxy. Starting in Wave 6, the activities of daily living include dressing, walking across a room, bathing, eating, getting in and out of bed, and using the toilet, and the instrumental activities of daily living include preparing hot meals, shopping for groceries, using the telephone, taking medication, using a map, doing work around the house or garden, managing money, leaving the house independently, and doing personal laundry. In Waves 6 and 7, help received for the 6 ADL activities and the 9 IADL activities and continence over urination or defecation are asked about in separate questions, so **raxrcarehr_s** sums the number of hours of help received for ADL activities (**raxracarehr**) and for IADL activities and continence (**raxricarehr_s**). In Wave 8 onwards, help received for the 6 ADL activities and the 9 IADL activities and continence over urination or defecation are asked about together, so **raxrcarehr_s** is the direct report of the number of hours of help received for ADL activities and for IADL activities and continence. **raxrcarehr_s** is calculated as long as at least one of these variables is not missing, and values can range from 0 to 48. **raxrcarehr_s** is coded as special missing (.x) if the respondent did not need help with any ADL or IADL activity. **raxrcarehr_s** is coded as special missing (.h) if the respondent had difficulty but did not receive any help with any ADL or IADL activity. This question is asked starting in Wave 6, and so **raxrcarehr_s** is assigned special missing (.q) in Waves 2 through 5. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Cross Wave Differences in SHARE EOL

In Waves 2, 3, 4, and 8 onwards, how long and the number of hours the respondent received help is asked for a combination of ADL and IADL activities. In Wave 5, how long and the number of hours the respondent received help is asked for ADL activities only, and not at all for IADL activities. In Waves 6 and 7, how long and the number of hours the respondent received help is asked separately for ADL activities and IADL activities.

Please note that 4 IADL activities (preparing hot meals, shopping for groceries, using the telephone, taking medication) are asked about in Waves 2 through 4, while 9 IADL activities and continence over urination or defecation are asked about starting in Wave 6.

Differences with the Harmonized HRS End of Life

In the HRS exit interview, the proxy is asked whether the respondent received help with each ADL activity and each of 4 IADL activities in the final three months of life, and if so, asked how frequently the respondent received help from each individual helper, which can be provided as days in a typical month, days per week, or everyday. These values are transformed into Harmonized variables indicating the number of days per week for each category of helper for the Harmonized HRS End of Life. However, in the SHARE end of life interview in Waves 2 through 4, the proxy is asked whether the respondent received help with a single list of ADL and 4 IADL activities in the last twelve months of life, and if so, asked how long the respondent had received help with any of the activities in the last year. In these waves, the question asking how long the respondent had needed help has specific response categories with a maximum of one year, and are reported as is and as a total for all helpers in the Gateway Harmonized SHARE End of Life. As such, these harmonized variables would need adjustment before being comparable.

In the SHARE end of life interview, in Waves 6 and 7, the proxy is asked whether the respondent received help with two separate lists of ADL and 9 IADL activities and continence over urination or defecation, and if so, asked how long the respondent

had needed help with the activities separately in the last year, and in Wave 8 onwards, the proxy is asked whether the respondent received help with ADL and 9 IADL activities and continence over urination or defecation, and if so, asked how long the respondent had needed help with the activities together in the last year. The response categories for the length of time the respondent has received help with ADLS and with IADLS and continence cannot be combined, so has not been created.

The variables in Harmonized HRS End of Life involve requiring help with 4 instrumental activities of daily living, which is comparable to the 4 IADLs in the Gateway Harmonized SHARE End of Life, while **raxrcarehr_s** in the Gateway Harmonized SHARE End of Life involves help with the same 4 instrumental activities of daily living plus 5 additional instrumental activities of daily living and help with continence over urination or defecation.

Also note that the amount of time each helper assisted the respondent are asked about individually in the HRS exit interview, which is summarized for each relationship category in the Harmonized HRS End of Life. However, the amount of time helpers assisted the respondent are asked about together in the SHARE end of life interview in Waves 2, 3, 4, and 8 onwards, and are asked about separately for ADL helpers and IADL helpers in Waves 6 and 7.

In the HRS exit interview, the proxy is only asked whether the respondent received help with the ADLs or IADLS but not whether they had difficulty with the ADLs or IADLS. In the SHARE end of life interview, the proxy is asked whether the respondent received help with the ADLs or IADLS and whether they had difficulty with the ADLs or IADLS. As such, the variables in the Harmonized HRS End of Life only employ a special missing (.h) indicating if they received no help, while the variables in the Gateway Harmonized SHARE End of Life employ a special missing (.h) as well as a special missing (.x) indicating if they had no difficulty.

Continuous Variable Distributions

Variable	N	Mean	Std.Dev	Minimum	Maximum	Total missing
raxrcarehr	1,747	10.89	9.18	0.00	24.00	19,351
raxrcarehr_s	9,704	15.40	14.61	0.00	48.00	11,394

Categorical Variable Frequencies

	raxrcared
1.less than 1 month	75
2.1 month+ but less than 3	177
3.3 months+ but less than 6	232
4.6 months+ but less than 1	264
5.a full year	1,145
.d:dk	60
.h:no help recvd	82
.m:missing	32
.q:not asked this wave	18,031
.r:refuse	9
.x:no difficulty	991
Total	21,098

	raxrcared_s
1.less than 1 month	576
2.1 month+ but less than 3	1,118
3.3 months+ but less than 6	1,250
4.6 months+ but less than 1	1,452
5.a full year	6,167
.d:dk	361
.h:no help recvd	766
.m:missing	82

.q:not asked this wave	5,254
.r:refuse	80
.x:no difficulty	3,992
Total	21,098

Original SHARE Variables Used

Wave 2 End of Life:

xt02odno	Difficulties doing activities: none of these
xt024_	Time the deceased received help
xt025_	Hours of help necessary during typical day

Wave 3 End of Life:

sl_xt02odno	Difficulties: none of these
sl_xt024_	Time the deceased received help
sl_xt025_	Hours of help necessary during typical day

Wave 4 End of Life:

xt02odno	Difficulties doing activities: none of these
xt024_	Time the deceased received help
xt025_	Hours of help necessary during typical day

Wave 6 End of Life:

xt022_	Anyone helped with ADL
xt024_	Time the deceased received help
xt025_	Hours of help necessary during typical day
xt02odno	Difficulties doing activities: none of these
xt622_	Anyone helped with ADLII
xt624_	Time the deceased received help with ADLII
xt625_	Hours of help with ADLII necessary during typical day
xt62odno	Difficulties doing activities II: none of these

Wave 7 End of Life:

xt022_	Anyone helped with ADL
xt024_	Time the deceased received help
xt025_	Hours of help necessary during typical day
xt02odno	Difficulties doing activities: none of these
xt622_	Anyone helped with ADLII
xt624_	Time the deceased received help with ADLII
xt625_	Hours of help with ADLII necessary during typical day
xt62odno	Difficulties doing activities II: none of these

Wave 8 End of Life:

xt022_	Anyone helped with ADL
xt024_	Time the deceased received help
xt025_	Hours of help necessary during typical day
xt02odno	Difficulties doing activities: none of these
xt62odno	Difficulties doing activities II: none of these

Wave 9 End of Life:

xt022_	Anyone helped with ADL
xt024_	Time the deceased received help
xt025_	Hours of help necessary during typical day
xt02odno	Difficulties doing activities: none of these
xt62odno	Difficulties doing activities II: none of these

Wave 9 COVID End of Life:

xt022_	Anyone helped with ADL
--------	------------------------

Section L. Assistance and Caregiving

xt024_	Time the deceased received help
xt025_	Hours of help necessary during typical day
xt02odno	Difficulties doing activities: none of these
xt62odno	Difficulties doing activities II: none of these

ADLs or IADLs: Whether Began Receiving Help within Final Year			
Variable	Waves	Label	Type
raxrcareb	All	raxrcareb: r began recv help with 10 adls/iadls final 12 months	Categ
raxrcareb_s	All	raxrcareb_s: r began recv help with 16 adls/iadls final 12 months	Categ

How Constructed

raxrcareb indicates whether the deceased respondent began receiving help with any of 6 activities of daily living (ADLs) or 4 instrumental activities of daily living (IADLs) in the last twelve months of life overall, as reported by the proxy. In Waves 2 through 4, the activities of daily living include dressing, walking across a room, bathing, eating, getting in and out of bed, and using the toilet, and the instrumental activities of daily living include preparing hot meals, shopping for groceries, using the telephone, and taking medication. In these waves, the 6 ADL activities and the 4 IADL activities are asked about in a single question. The proxy is first asked whether the respondent had difficulty with each ADL and IADL, and if the respondent had difficulty with at least one ADL or IADL, then the proxy is asked whether someone helped the respondent, and if so how long the respondent received help during the last twelve months of life. **raxrcareb** is coded as 0 if the respondent received help with at least one of 6 ADL or 4 IADL activities and the respondent required help for the full year. **raxrcareb** is coded as 1 if the respondent received help with at least one of 6 ADL or 4 IADL activities and the respondent received help for less than one month, one month or more but less than three months, three months or more but less than six months, or six months or more but less than a year. **raxrcareb** is coded as special missing (.x) if the respondent did not need help with any ADL or IADL activity. **raxrcareb** is coded as special missing (.h) if the respondent had difficulty but did not receive any help with any ADL or IADL activity. This question is no longer asked with the same combination of activities after Wave 4, and so **raxrcareb** is assigned special missing (.q) starting in Wave 5. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

raxrcareb_s indicates whether the deceased respondent began receiving help with any of 6 activities of daily living (ADLs) or 9 instrumental activities of daily living (IADLs) or with continence over urination or defecation in the last twelve months of life overall, as reported by the proxy. Starting in Wave 6, the activities of daily living include dressing, walking across a room, bathing, eating, getting in and out of bed, and using the toilet, and the instrumental activities of daily living include preparing hot meals, shopping for groceries, using the telephone, taking medication, using a map, doing work around the house or garden, managing money, leaving the house independently, and doing personal laundry. In Waves 6 and 7, the 6 ADL activities and the 9 IADL activities and continence over urination or defecation are asked about in separate questions, so **raxrcareb_s** summarizes the information for when the respondent began needing help with ADLs (**raxracareb**) and IADLs and continence (**raxricareb_s**). In Wave 8 onwards, the 6 ADL activities and the 9 IADL activities and continence over urination or defecation are asked about together, so **raxrcareb_s** directly reports the information for when the respondent began needing help with ADLs and IADLs and continence. **raxrcareb_s** is coded as 0 if the respondent received help with at least one of 16 ADL or IADL activities and the respondent required help for the full year. **raxrcareb_s** is coded as 1 if the respondent received help with at least one of 16 ADL or IADL activities and the respondent received help for less than one month, one month or more but less than three months, three months or more but less than six months, or six months or more but less than a year. **raxrcareb_s** is coded as special missing (.x) if the respondent did not need help with any ADL or IADL activity. **raxrcareb_s** is coded as special missing (.h) if the respondent had difficulty but did not receive any help with any ADL or IADL activity. This question is asked starting in Wave 6, and so **raxrcareb_s** is assigned special missing (.q) in Waves 2 through 5. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Cross Wave Differences in SHARE EOL

In Waves 2 through 4, how long the respondent received help is asked for a combination of ADL and IADL activities. In Wave 5, how long the respondent received help is asked for ADL activities only, and not at all for IADL activities. In Waves 6 and 7, how long the respondent received help is asked separately for ADL activities and IADL activities. In Wave 8 onwards, how long the respondent received help is asked jointly for ADL activities and IADL activities.

Please note that 4 IADL activities (preparing hot meals, shopping for groceries, using the telephone, taking medication) are asked about in Waves 2 through 4, while 9 IADL activities and continence over urination or defecation are asked about starting

in Wave 6.

Differences with the Harmonized HRS End of Life

In the HRS exit interview, the proxy is asked whether the respondent received help with each ADL activity and each of 4 IADL activities in the final three months of life, and if so, asked how long the respondent had needed help with each ADL and IADL, which can be provided in months, years, since age, or since year. These responses have been converted to be equivalent to approximately 1 year in this variable in the Harmonized HRS End of Life. However, in the SHARE end of life interview in Waves 2 through 4, the proxy is asked whether the respondent received help with a single list of ADL and 4 IADL activities in the last twelve months of life, and if so, asked how long the respondent had received help with any of the activities in the last year. In the SHARE end of life interview in Waves 6 and 7, the proxy is asked whether the respondent received help with two separate lists of ADL and 9 IADL activities and continence over urination or defecation, and if so, asked how long the respondent had received help with the activities separately in the last year. In Wave 8 onwards, instead the proxy is asked whether the respondent received help with two lists of ADL and 9 IADL activities and continence over urination or defecation in a single question, and if so, asked how long the respondent had received help with the activities jointly in the last year. Each question asking how long the respondent had needed help has specific response categories with a maximum of one year. These responses have been converted so that responses of a full year are coded as not having begun to need help in the last year of life in the Gateway Harmonized SHARE End of Life.

As such, this variable in the Harmonized HRS End of Life captures a slightly different concept than **raxrcareb** and **raxrcareb_s** in the Gateway Harmonized SHARE End of Life because we can ascertain with some degree of certainty when the respondent began needing help based on the questions in the HRS exit interview, whereas we are assuming that the respondent began needing help prior to the last twelve months of life if the respondent received help during their entire final year in the SHARE end of life interview.

Additionally, this variable in the Harmonized HRS End of Life involves requiring help with 4 instrumental activities of daily living, which is comparable to the 4 IADLs in **raxrcareb** in the Gateway Harmonized SHARE End of Life, while **raxrcareb_s** in the Gateway Harmonized SHARE End of Life involves help with the same 4 instrumental activities of daily living plus 5 additional instrumental activities of daily living and help with continence over urination or defecation.

Also note that ADL and IADL activities are asked about individually in the HRS exit interview, so this variable summarizes the information from each activity in the Harmonized HRS End of Life. However, ADL and IADL activities are asked about together in the SHARE end of life interview in Waves 2, 3, 4, and 8 onwards, and are asked about separately as groups of ADLs and IADLs in Waves 6 and 7, so **raxrcareb** is taken directly from a single question in Waves 2 through 4 of the Gateway Harmonized SHARE End of Life, and **raxrcareb_s** summarizes the information from 2 separate questions in Wave 6 and 7 while **raxrcareb_s** is taken directly from a single question in Wave 8 onwards.

In the HRS exit interview, the proxy is only asked whether the respondent received help with the ADLs or IADLs but not whether they had difficulty with the ADLs or IADLs. In the SHARE end of life interview, the proxy is asked whether the respondent received help with the ADLs or IADLs and whether they had difficulty with the ADLs or IADLs. As such, the variables in the Harmonized HRS End of Life only employ a special missing (.h) indicating if they received no help, while the variables in the Gateway Harmonized SHARE End of Life employ a special missing (.h) as well as a special missing (.x) indicating if they had no difficulty.

Categorical Variable Frequencies

	raxrcareb
0.no	1,145
1.yes	748
.d:dk	60
.h:no help recvd	82
.m:missing	32
.q:not asked this wave	18,031
.r:refuse	9

.x:no difficulty	991
Total	21,098
raxrcareb_s	
0.no	5,977
1.yes	4,665
.d:dk	299
.h:no help recvd	766
.m:missing	78
.q:not asked this wave	5,254
.r:refuse	67
.x:no difficulty	3,992
Total	21,098

Original SHARE Variables Used

Wave 2 End of Life:

xt02odno	Difficulties doing activities: none of these
xt024_	Time the deceased received help

Wave 3 End of Life:

sl_xt02odno	Difficulties: none of these
sl_xt024_	Time the deceased received help

Wave 4 End of Life:

xt02odno	Difficulties doing activities: none of these
xt024_	Time the deceased received help

Wave 6 End of Life:

xt022_	Anyone helped with ADL
xt024_	Time the deceased received help
xt02odno	Difficulties doing activities: none of these
xt622_	Anyone helped with ADLII
xt624_	Time the deceased received help with ADLII
xt62odno	Difficulties doing activities II: none of these

Wave 7 End of Life:

xt022_	Anyone helped with ADL
xt024_	Time the deceased received help
xt02odno	Difficulties doing activities: none of these
xt622_	Anyone helped with ADLII
xt624_	Time the deceased received help with ADLII
xt62odno	Difficulties doing activities II: none of these

Wave 8 End of Life:

xt022_	Anyone helped with ADL
xt024_	Time the deceased received help
xt02odno	Difficulties doing activities: none of these
xt62odno	Difficulties doing activities II: none of these

Wave 9 End of Life:

xt022_	Anyone helped with ADL
xt024_	Time the deceased received help
xt02odno	Difficulties doing activities: none of these
xt62odno	Difficulties doing activities II: none of these

Wave 9 COVID End of Life:

xt022_	Anyone helped with ADL
--------	------------------------

Section L. Assistance and Caregiving

xt024_	Time the deceased received help
xt02odno	Difficulties doing activities: none of these
xt62odno	Difficulties doing activities II: none of these

O. End of Life Planning

Wills: Whether Had a Will

Variable	Waves	Label	Type
rawitwill	All	rawitwill: r had a will	Categ

How Constructed

rawitwill indicates whether the deceased respondent had a will. **rawitwill** is coded as 0 if the respondent did not have a will. **rawitwill** is coded as 1 if the respondent did have a will. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Cross Wave Differences in SHARE EOL

No differences known.

Differences with the Harmonized HRS End of Life

The HRS exit interview inquires as to whether the deceased respondent had a written and witnessed will, whereas the SHARE end of life interview inquires as to whether the deceased respondent had a will. The responses from both of these questions are recorded in **rawitwill** in the Harmonized HRS End of Life and the Gateway Harmonized SHARE End of Life, respectively.

Categorical Variable Frequencies

	rawitwill
0.no	14,545
1.yes	4,741
.d:dk	1,319
.m:missing	214
.r:refuse	279
Total	21,098

Original SHARE Variables Used

Wave 2 End of Life:	
xto26_	The deceased had a will
Wave 3 End of Life:	
sl_xto26_	The deceased had a will
Wave 4 End of Life:	
xto26_	The deceased had a will
Wave 5 End of Life:	
xto26_	The deceased had a will
Wave 6 End of Life:	
xto26_	The deceased had a will
Wave 7 End of Life:	
xto26_	The deceased had a will
Wave 8 End of Life:	
xto26_	The deceased had a will
Wave 9 End of Life:	
xto26_	The deceased had a will
Wave 9 COVID End of Life:	

xt026_

The deceased had a will

Wills: Estate Beneficiaries			
Variable	Waves	Label	Type
rabnfcersp	All	rabnfcersp: spouse was beneficiary of r's estate	Categ
rabnfcrcg	All	rabnfcrcg: child/grandchild was beneficiary of r's estate	Categ
rabnfcrrl	All	rabnfcrrl: relative was beneficiary of r's estate	Categ
rabnfcrot	All	rabnfcrot: other was beneficiary of r's estate	Categ
rabnfcrin	All	rabnfcrin: charity was beneficiary of r's estate	Categ

How Constructed

rabnfcersp, **rabnfcrcg**, **rabnfcrrl**, **rabnfcrot**, and **rabnfcrin** indicate whether a person with the specified relationship was a beneficiary of the respondent's estate, as reported by the proxy. The proxy can select as many relationships as applicable from a list of relationships which includes the proxy, and if the proxy is selected, then the relationship of the proxy to the respondent is used to assign values to the correct variables. **rabnfcersp** indicates whether the respondent's husband/wife/partner was a beneficiary of the respondent's estate. **rabnfcrcg** indicates whether the respondent's children or grandchildren were a beneficiary of the respondent's estate. **rabnfcrrl** indicates whether the respondent's siblings or other relatives were a beneficiary of the respondent's estate. **rabnfcrot** indicates whether other non-relatives were a beneficiary of the respondent's estate. **rabnfcrin** indicates whether a church, foundation, or charitable organization was a beneficiary of the respondent's estate. These variables are assigned a 0 if a person of the specified relationship was not mentioned as a beneficiary of the respondent's estate or if the proxy responded that the deceased did not leave anything at all, and are assigned a 1 if a person of the specified relationship was mentioned as a beneficiary of the respondent's estate. Starting in Wave 7, special missing code (.n) is assigned if the proxy responded that the beneficiaries of the estate have not been decided yet. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Cross Wave Differences in SHARE EOL

Starting in Wave 7, when asked about the beneficiaries of the estate, the questionnaire allows the proxy to respond that the beneficiaries have not been decided yet.

Differences with the Harmonized HRS End of Life

In the HRS exit interview, if the respondent has a will or trust, then the proxy is directly asked whether the respondent made provisions in the will or trust for their spouse, children/grandchildren/great-grandchildren, charities, siblings, other relatives, friends, or anyone else or anything else in separate questions. If the respondent does not have a will or trust, then the proxy is asked whether these groups were left any possessions or assets by the respondent. The proxy is not asked if the respondent left anything to the spouse or children/grandchildren/great-grandchildren if the respondent did not have a spouse or any children.

The SHARE end of life interview inquires who the beneficiaries of the respondent's estate were using the same question wording whether the respondent had a will or not. The questions asked are not limited by the absence of a living spouse or children.

SHARE allows the proxy to state that the beneficiaries have not yet been decided, while the HRS allows the proxy to state that the estate has not yet been distributed. Both of these require separate special missing values in these variables in the respective Harmonized End of Life datasets.

The categories used in most waves of the HRS exit interview for this question include: husband/wife/partner, children/grandchildren/great-grandchildren, charities, brothers or sisters, any other relatives, any friends, anyone else or anything else. The categories used in the SHARE end of life interview for this question include: proxy, husband/wife/partner, children, grandchildren, siblings, other relatives, other non-relatives, church or foundation or charitable organization. In the Harmonized HRS End of Life, **rabnfcrcg** includes children and grandchildren taken from a single question, while in the Gateway Harmonized SHARE End of Life, **rabnfcrcg** includes children and grandchildren taken from separate categories.

In the Harmonized HRS End of Life, **rabnfcrot** includes any friends and anyone else or anything else taken from two questions, while in the Gateway Harmonized SHARE End of Life, **rabnfcrot** includes other non-relatives taken from a single category.

Categorical Variable Frequencies

	rabnfcrsp
0.no	10,763
1.yes	8,293
.d:dk	979
.m:missing	194
.n:not decided yet	480
.r:refuse	389
Total	21,098

	rabnfcrcg
0.no	7,159
1.yes	11,900
.d:dk	979
.m:missing	194
.n:not decided yet	477
.r:refuse	389
Total	21,098

	rabnfcrrl
0.no	17,802
1.yes	1,250
.d:dk	979
.m:missing	194
.n:not decided yet	484
.r:refuse	389
Total	21,098

	rabnfcrot
0.no	18,870
1.yes	182
.d:dk	979
.m:missing	194
.n:not decided yet	484
.r:refuse	389
Total	21,098

	rabnfcrin
0.no	18,975
1.yes	75
.d:dk	979
.m:missing	194
.n:not decided yet	486
.r:refuse	389
Total	21,098

Original SHARE Variables Used

Wave 2 End of Life:

xto02_	Relationship to the deceased
xto27d1	Beneficiaries of the estate: yourself (proxy)
xto27d2	Beneficiaries of the estate: spouse/partner of the deceased
xto27d3	Beneficiaries of the estate: children of the deceased
xto27d4	Beneficiaries of the estate: grandchildren of the deceased
xto27d5	Beneficiaries of the estate: siblings of the deceased
xto27d6	Beneficiaries of the estate: other relatives of the deceased
xto27d7	Beneficiaries of the estate: other non-relatives
xto27d8	Beneficiaries of the estate: church, foundation or charitable organization
xto27d9	Beneficiaries of the estate: deceased did not leave anything at all (SPONTANEOUS)

Wave 3 End of Life:

sl_xto02_	Relationship to the deceased
sl_xto27d1	Beneficiaries of estate: proxy respondent
sl_xto27d2	Beneficiaries of estate: husband/wife/partner of the deceased
sl_xto27d3	Beneficiaries of estate: children of the deceased
sl_xto27d4	Beneficiaries of estate: grandchildren of the deceased
sl_xto27d5	Beneficiaries of estate: siblings of the deceased
sl_xto27d6	Beneficiaries of estate: other relatives of the deceased
sl_xto27d7	Beneficiaries of estate: other non-relatives
sl_xto27d8	Beneficiaries of estate: church, foundation or charitable organization
sl_xto27d9	Beneficiaries of estate: deceased did not leave anything at all (spontaneous)

Wave 4 End of Life:

xto02_	Relationship to the deceased
xto27d1	Beneficiaries of the estate: yourself (proxy)
xto27d2	Beneficiaries of the estate: spouse/partner of the deceased
xto27d3	Beneficiaries of the estate: children of the deceased
xto27d4	Beneficiaries of the estate: grandchildren of the deceased
xto27d5	Beneficiaries of the estate: siblings of the deceased
xto27d6	Beneficiaries of the estate: other relatives of the deceased
xto27d7	Beneficiaries of the estate: other non-relatives
xto27d8	Beneficiaries of the estate: church, foundation or charitable organization
xto27d9	Beneficiaries of the estate: deceased did not leave anything at all (SPONTANEOUS)

Wave 5 End of Life:

xto02_	Relationship to the deceased
xto27d1	Beneficiaries of the estate: yourself (proxy)
xto27d2	Beneficiaries of the estate: spouse/partner of the deceased
xto27d3	Beneficiaries of the estate: children of the deceased
xto27d4	Beneficiaries of the estate: grandchildren of the deceased
xto27d5	Beneficiaries of the estate: siblings of the deceased
xto27d6	Beneficiaries of the estate: other relatives of the deceased
xto27d7	Beneficiaries of the estate: other non-relatives
xto27d8	Beneficiaries of the estate: church, foundation or charitable organization
xto27d9	Beneficiaries of the estate: deceased did not leave anything at all (SPONTANEOUS)

Wave 6 End of Life:

xto02_	Relationship to the deceased
xto27d1	Beneficiaries of the estate: yourself (proxy)
xto27d2	Beneficiaries of the estate: spouse/partner of the deceased
xto27d3	Beneficiaries of the estate: children of the deceased
xto27d4	Beneficiaries of the estate: grandchildren of the deceased
xto27d5	Beneficiaries of the estate: siblings of the deceased

xto27d6	Beneficiaries of the estate: other relatives of the deceased
xto27d7	Beneficiaries of the estate: other non-relatives
xto27d8	Beneficiaries of the estate: church, foundation or charitable organization
xto27d9	Beneficiaries of the estate: deceased did not leave anything at all (SPONTANEOUS)
Wave 7 End of Life:	
xto02_	Relationship to the deceased
xto27d1	Beneficiaries of the estate: yourself (proxy)
xto27d2	Beneficiaries of the estate: spouse/partner of the deceased
xto27d3	Beneficiaries of the estate: children of the deceased
xto27d4	Beneficiaries of the estate: grandchildren of the deceased
xto27d5	Beneficiaries of the estate: siblings of the deceased
xto27d6	Beneficiaries of the estate: other relatives of the deceased
xto27d7	Beneficiaries of the estate: other non-relatives
xto27d8	Beneficiaries of the estate: church, foundation or charitable organization
xto27d9	Beneficiaries of the estate: deceased did not leave anything at all (SPONTANEOUS)
xto27d10	Beneficiaries of the estate: not decided yet (SPONTANEOUS)
Wave 8 End of Life:	
xto02_	Relationship to the deceased
xto27d1	Beneficiaries of the estate: yourself (proxy)
xto27d2	Beneficiaries of the estate: spouse/partner of the deceased
xto27d3	Beneficiaries of the estate: children of the deceased
xto27d4	Beneficiaries of the estate: grandchildren of the deceased
xto27d5	Beneficiaries of the estate: siblings of the deceased
xto27d6	Beneficiaries of the estate: other relatives of the deceased
xto27d7	Beneficiaries of the estate: other non-relatives
xto27d8	Beneficiaries of the estate: church, foundation or charitable organization
xto27d9	Beneficiaries of the estate: deceased did not leave anything at all (SPONTANEOUS)
xto27d10	Beneficiaries of the estate: not decided yet (SPONTANEOUS)
Wave 9 End of Life:	
xto02_	Relationship to the deceased
xto27d1	Beneficiaries of the estate: yourself (proxy)
xto27d2	Beneficiaries of the estate: spouse/partner of the deceased
xto27d3	Beneficiaries of the estate: children of the deceased
xto27d4	Beneficiaries of the estate: grandchildren of the deceased
xto27d5	Beneficiaries of the estate: siblings of the deceased
xto27d6	Beneficiaries of the estate: other relatives of the deceased
xto27d7	Beneficiaries of the estate: other non-relatives
xto27d8	Beneficiaries of the estate: church, foundation or charitable organization
xto27d9	Beneficiaries of the estate: deceased did not leave anything at all (SPONTANEOUS)
xto27d10	Beneficiaries of the estate: not decided yet (SPONTANEOUS)
Wave 9 COVID End of Life:	
xto02_	Relationship to the deceased
xto27d1	Beneficiaries of the estate: yourself (proxy)
xto27d2	Beneficiaries of the estate: spouse/partner of the deceased
xto27d3	Beneficiaries of the estate: children of the deceased
xto27d4	Beneficiaries of the estate: grandchildren of the deceased
xto27d5	Beneficiaries of the estate: siblings of the deceased
xto27d6	Beneficiaries of the estate: other relatives of the deceased
xto27d7	Beneficiaries of the estate: other non-relatives
xto27d8	Beneficiaries of the estate: church, foundation or charitable organization
xto27d9	Beneficiaries of the estate: deceased did not leave anything at all (SPONTANEOUS)
xto27d10	Beneficiaries of the estate: not decided yet (SPONTANEOUS)

Life Insurance Policies: Whether Had Life Insurance and Value

Variable	Waves	Label	Type
raxlifein	All	raxlifein: r any life insurance policies	Categ
raxlifeinv	All	raxlifeinv: r life insurance: total value of policies	Cont
raxlifeinvf	All	raxlifeinvf: r life insurance flag: total value of policies	Categ

How Constructed

raxlifein indicates whether the deceased respondent owned any life insurance policies. **raxlifein** is coded as 0 if the respondent did not own any life insurance policies. **raxlifein** is coded as 1 if the respondent did own a life insurance policy. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

raxlifeinv has been adjusted to 2010 Euros based on the consumer price index for the year and country of death.

raxlifeinv indicates the total value of all life insurance policies owned by the respondent and includes imputed values. If the respondent owned any life insurance policies, then the proxy is asked the approximate total value of all life insurance policies owned by the respondent. The value of the life insurance policies is elicited in Euros in all SHARE countries except in Denmark, Sweden, Switzerland, Poland, Czech Republic, and Israel where local currencies are used and which have been converted to Euros by the SHARE team; however, all reported or imputed values in **raxlifeinv** have been converted and adjusted to 2010 Euros based on the consumer price index for the year and country of death. **raxlifeinv** is assigned a value of 0 if the respondent did not own any life insurance policies. If the proxy answers don't know or refuses to answer the value of the life insurance policies, the proxy is not asked any additional questions to obtain threshold values; instead values are imputed with no bracket information. **raxlifeinv** is assigned special missing (.m) if the values are not able to be imputed due to a lack of reference observations in the donor pool or missing covariates necessary for the imputation procedure.

raxlifeinvf is a flag variable indicating the level of imputation for the total value of life insurance policies. A code of 1 indicates the proxy reported a continuous value and no imputation was necessary. A code of 5 indicates that the value was imputed without any bracket information. A code of 6 indicates the proxy reported the respondent not having any life insurance policies and the value is 0. A code of 7 indicates that the proxy was not sure whether the respondent had any life insurance policies. A code of -1 indicates that the value could not be imputed due to a lack of reference observations in the donor pool. A code of -2 indicates that the expense could not be imputed due to one or more missing values in the covariates used in the imputation procedure.

Cross Wave Differences in SHARE EOL

No differences known.

Differences with the Harmonized HRS End of Life

raxlifeins indicates whether anyone received a settlement from the deceased respondent's life insurance in the Harmonized HRS End of Life, whereas **raxlifein** indicates whether the respondent owned any life insurance policies in the Gateway Harmonized SHARE End of Life. While the two concepts are very similar, it is possible to own a life insurance policy and to ultimately not receive a settlement from it, as such we have provided different variable names for these two concepts.

In the HRS exit interview, the proxy is asked what amounts or percentages that each beneficiary received from the deceased's life insurance policy. Depending on the wave, proxies might also be asked the total amount of life insurance received altogether. In the SHARE end of life interview, on the other hand, proxies are asked a single direct question asking for the total of the deceased's owned life insurance policies.

In the HRS exit interview, if the proxy responds that they don't know or refuse to answer the value of the life insurance settlement, then a series of unfolding bracket questions is asked to obtain a minimum and maximum value of the settlement.

However, in the SHARE end of life interview, no unfolding bracket questions are asked to the proxy if they respond that they don't know or refuse to obtain the approximate value of the life insurance policies.

The monetary values in the Harmonized HRS End of Life are provided in 2010 U.S. dollars, whereas monetary values in the Gateway Harmonized SHARE End of Life are provided in 2010 Euro. As such, adjustment must be made before any comparison of these variables.

Continuous Variable Distributions

Variable	N	Mean	Std.Dev	Minimum	Maximum	Total missing
raxlifeinv	21,026	2,576.06	39,686.41	0.00	2,042,900.88	72

Categorical Variable Frequencies

	raxlifein
0.no	17,368
1.yes	1,917
.d:dk	1,335
.m:missing	219
.r:refuse	259
Total	21,098
	raxlifeinvf
-2.not imputed, missing cov	67
-1.not imputed, missing nei	5
1.continuous value	1,235
5.no bracket info	677
6.no asset	17,368
7.ownership unknown	1,746
Total	21,098

Original SHARE Variables Used

Wave 2 End of Life:

xt033_ Deceased owned any life insurance policies
xt034e Value of all life insurance policies

Wave 3 End of Life:

sl_xt033_ The deceased owned any life insurance policies
sl_xt034e Value of all life insurance policies

Wave 4 End of Life:

xt033_ Deceased owned any life insurance policies
xt034e Value of all life insurance policies

Wave 5 End of Life:

xt033_ Deceased owned any life insurance policies
xt034e Value of all life insurance policies

Wave 6 End of Life:

xt033_ Deceased owned any life insurance policies
xt034e Value of all life insurance policies

Wave 7 End of Life:

xt033_ Deceased owned any life insurance policies
xt034e Value of all life insurance policies

Wave 8 End of Life:

xt033_	Deceased owned any life insurance policies
xt034e	Value of all life insurance policies

Wave 9 End of Life:

xt033_	Deceased owned any life insurance policies
xt034e	Value of all life insurance policies

Wave 9 COVID End of Life:

xt033_	Deceased owned any life insurance policies
xt034e	Value of all life insurance policies

Life Insurance Policies: Beneficiaries

Variable	Waves	Label	Type
ralfinssp	All	ralfinssp: r spouse beneficiary of life ins	Categ
ralfinsch	All	ralfinsch: r child beneficiary of life ins	Categ
ralfinsgk	All	ralfinsgk: r grandchild beneficiary of life ins	Categ
ralfinsrl	All	ralfinsrl: r relative beneficiary of life ins	Categ
ralfinsot	All	ralfinsot: r other non-relative beneficiary of life ins	Categ

How Constructed

ralfinssp, **ralfinsch**, **ralfinsgk**, **ralfinsrl**, and **ralfinsot** indicate whether someone with the specified relationship were beneficiaries of the respondent's life insurance policies. The proxy is first asked whether the respondent owned any life insurance policies, and if so, the proxy can report the relationship of as many people as applicable who were beneficiaries of the life insurance policies. **ralfinssp** indicates whether the deceased respondent's spouse/partner was a beneficiary of the respondent's life insurance policies. **ralfinsch** indicates whether the deceased respondent's natural, adopted, foster, or step-children were beneficiaries of the respondent's life insurance policies. **ralfinsgk** indicates whether the deceased respondent's grandchildren were beneficiaries of the respondent's life insurance policies. **ralfinsrl** indicates whether other relatives of the deceased respondent were beneficiaries of the respondent's life insurance policies. Other relatives include natural parents, adopted parents, foster parents, step-parents, parents-in-law, natural siblings, half-siblings, step-siblings, adopted siblings, foster siblings, siblings-in-law, grandparents, or other relatives. **ralfinsot** indicates whether other non-relatives of the deceased respondent were beneficiaries of the respondent's life insurance policies. **ralfinssp**, **ralfinsch**, **ralfinsgk**, **ralfinsrl**, and **ralfinsot** are coded as 0 if that relationship is not mentioned as being a beneficiary of the respondent's life insurance policies, and coded as 1 if that relationship is mentioned as being a beneficiary of the respondent's life insurance policies. **ralfinssp**, **ralfinsch**, **ralfinsgk**, **ralfinsrl**, and **ralfinsot** are assigned special missing (.n) if the respondent had no life insurance policies. Don't know, refused, or other missing responses are assigned special missing (.d), (.r), (.m), respectively.

Cross Wave Differences in SHARE EOL

No differences known.

Differences with the Harmonized HRS End of Life

In the HRS exit interview, the proxy is asked who received a life insurance settlement, and between 2 and 4 relationships can be specified as recipients of life insurance settlements depending on the wave. In the SHARE end of life interview, the proxy is asked who was a beneficiary of the respondent's life insurance policies, and can select all the relationships that apply.

The categories used in most waves of the HRS exit interview for this question include: spouse/partner, children, grandchildren or great-grandchildren, other relatives, and other. The categories used in the SHARE end of life interview for this question include: proxy, husband/wife/partner, sons or daughters, grandchildren, siblings, other relatives, and other. In the Harmonized HRS End of Life, **ralfinsgk** includes the respondent's grandchildren or great-grandchildren from a single category, whereas **ralfinsgk** in the Gateway Harmonized SHARE End of Life includes the respondent's grandchildren only. In the Harmonized HRS End of Life, **ralfinsrl** includes the respondent's other relatives, whereas **ralfinsrl** in the Gateway Harmonized SHARE End of Life includes the respondent's siblings or other relatives taken from two separate categories.

Categorical Variable Frequencies

	ralfinssp
0.no	693
1.yes	1,131
.d:dk	1,395

.m:missing	222
.n:no life insurance policy	17,368
.r:refuse	289
Total	21,098
ralfinsch	
o.no	1,122
1.yes	702
.d:dk	1,391
.m:missing	230
.n:no life insurance policy	17,368
.r:refuse	285
Total	21,098
ralfinsgk	
o.no	1,767
1.yes	57
.d:dk	1,395
.m:missing	222
.n:no life insurance policy	17,368
.r:refuse	289
Total	21,098
ralfinsrl	
o.no	1,720
1.yes	104
.d:dk	1,395
.m:missing	222
.n:no life insurance policy	17,368
.r:refuse	289
Total	21,098
ralfinsot	
o.no	1,796
1.yes	28
.d:dk	1,395
.m:missing	222
.n:no life insurance policy	17,368
.r:refuse	289
Total	21,098

Original SHARE Variables Used

Wave 2 End of Life:

xto33_	Deceased owned any life insurance policies
xto35d1	Beneficiaries of the life insurance policies: yourself (proxy)
xto35d2	Beneficiaries of the life insurance policies: spouse/partner of the deceased
xto35d3	Beneficiaries of the life insurance policies: sons/daughters
xto35d4	Beneficiaries of the life insurance policies: grandchildren of the deceased
xto35d5	Beneficiaries of the life insurance policies: siblings of the deceased
xto35d6	Beneficiaries of the life insurance policies: other relatives
xto35d7	Beneficiaries of the life insurance policies: other non-relatives

Wave 3 End of Life:

sl_xt033_	The deceased owned any life insurance policies
sl_xt035d1	Beneficiaries of life insurance policies: proxy respondent
sl_xt035d2	Beneficiaries of life insurance policies: husband/wife/partner of the deceased
sl_xt035d3	Beneficiaries of life insurance policies: sons or daughters
sl_xt035d4	Beneficiaries of life insurance policies: grandchildren
sl_xt035d5	Beneficiaries of life insurance policies: siblings
sl_xt035d6	Beneficiaries of life insurance policies: other relatives
sl_xt035d7	Beneficiaries of life insurance policies: other non-relatives

Wave 4 End of Life:

xt033_	Deceased owned any life insurance policies
xt035d1	Beneficiaries of the life insurance policies: yourself (proxy)
xt035d2	Beneficiaries of the life insurance policies: spouse/partner of the deceased
xt035d3	Beneficiaries of the life insurance policies: sons/daughters
xt035d4	Beneficiaries of the life insurance policies: grandchildren of the deceased
xt035d5	Beneficiaries of the life insurance policies: siblings of the deceased
xt035d6	Beneficiaries of the life insurance policies: other relatives
xt035d7	Beneficiaries of the life insurance policies: other non-relatives

Wave 5 End of Life:

xt033_	Deceased owned any life insurance policies
xt035d1	Beneficiaries of the life insurance policies: yourself (proxy)
xt035d2	Beneficiaries of the life insurance policies: spouse/partner of the deceased
xt035d3	Beneficiaries of the life insurance policies: sons/daughters
xt035d4	Beneficiaries of the life insurance policies: grandchildren of the deceased
xt035d5	Beneficiaries of the life insurance policies: siblings of the deceased
xt035d6	Beneficiaries of the life insurance policies: other relatives
xt035d7	Beneficiaries of the life insurance policies: other non-relatives

Wave 6 End of Life:

xt033_	Deceased owned any life insurance policies
xt035d1	Beneficiaries of the life insurance policies: yourself (proxy)
xt035d2	Beneficiaries of the life insurance policies: spouse/partner of the deceased
xt035d3	Beneficiaries of the life insurance policies: sons/daughters
xt035d4	Beneficiaries of the life insurance policies: grandchildren of the deceased
xt035d5	Beneficiaries of the life insurance policies: siblings of the deceased
xt035d6	Beneficiaries of the life insurance policies: other relatives
xt035d7	Beneficiaries of the life insurance policies: other non-relatives

Wave 7 End of Life:

xt033_	Deceased owned any life insurance policies
xt035d1	Beneficiaries of the life insurance policies: yourself (proxy)
xt035d2	Beneficiaries of the life insurance policies: spouse/partner of the deceased
xt035d3	Beneficiaries of the life insurance policies: sons/daughters
xt035d4	Beneficiaries of the life insurance policies: grandchildren of the deceased
xt035d5	Beneficiaries of the life insurance policies: siblings of the deceased
xt035d6	Beneficiaries of the life insurance policies: other relatives
xt035d7	Beneficiaries of the life insurance policies: other non-relatives

Wave 8 End of Life:

xt033_	Deceased owned any life insurance policies
xt035d1	Beneficiaries of the life insurance policies: yourself (proxy)
xt035d2	Beneficiaries of the life insurance policies: spouse/partner of the deceased
xt035d3	Beneficiaries of the life insurance policies: sons/daughters
xt035d4	Beneficiaries of the life insurance policies: grandchildren of the deceased
xt035d5	Beneficiaries of the life insurance policies: siblings of the deceased
xt035d6	Beneficiaries of the life insurance policies: other relatives

xt035d7	Beneficiaries of the life insurance policies: other non-relatives
Wave 9 End of Life:	
xt033_	Deceased owned any life insurance policies
xt035d1	Beneficiaries of the life insurance policies: yourself (proxy)
xt035d2	Beneficiaries of the life insurance policies: spouse/partner of the deceased
xt035d3	Beneficiaries of the life insurance policies: sons/daughters
xt035d4	Beneficiaries of the life insurance policies: grandchildren of the deceased
xt035d5	Beneficiaries of the life insurance policies: siblings of the deceased
xt035d6	Beneficiaries of the life insurance policies: other relatives
xt035d7	Beneficiaries of the life insurance policies: other non-relatives
Wave 9 COVID End of Life:	
xt033_	Deceased owned any life insurance policies
xt035d1	Beneficiaries of the life insurance policies: yourself (proxy)
xt035d2	Beneficiaries of the life insurance policies: spouse/partner of the deceased
xt035d3	Beneficiaries of the life insurance policies: sons/daughters
xt035d4	Beneficiaries of the life insurance policies: grandchildren of the deceased
xt035d5	Beneficiaries of the life insurance policies: siblings of the deceased
xt035d6	Beneficiaries of the life insurance policies: other relatives
xt035d7	Beneficiaries of the life insurance policies: other non-relatives

6. References

Ailshire J, Chien S, Phillips D, Wilkens J, Lee J. 2019. Harmonized HRS End of Life Documentation, Version A (1992-2014). University of Southern California, Center for Economic and Social Research, Los Angeles, CA.
https://eol.g2aging.org/hrd/get_data

Bollinger CR, Hirsch BT. 2006. Match bias from earnings imputation in the Current Population Survey: the case of imperfect matching. *Journal of Labor Economics*, Volume 24, 483-519.
https://www.researchgate.net/profile/Barry_Hirsch2/publication/24099977_Match_Bias_from_Earnings_Imputation_in_the_Current_Population_Survey-The-Case-of-Imperfect-Matching.pdf

Börsch-Supan A, Hank K, Jürges H, Schraeder M. 2008. Longitudinal Data Collection in Continental Europe: Experiences from the Survey of Health, Ageing and Retirement in Europe (SHARE). SHARE Working Paper 162/2008.
<https://ideas.repec.org/p/mea/meawpa/o8162.html>

Börsch-Supan A, Jürges H (eds). 2005. The Survey of Health, Aging, and Retirement in Europe - Methodology. Mannheim Research Institute for the Economics of Aging (MEA): Mannheim, Germany.
https://share-eric.eu/fileadmin/user_upload/Methodology_Volumes/Methodology_2005.pdf

Hirsch BT, Schumacher EJ. 2004. Match bias in wage gap estimates due to earnings imputation. *Journal of Labor Economics*, Volume 22, 689-722.
<https://www.journals.uchicago.edu/doi/abs/10.1086/383112?journalCode=jole>

Lee J, Meijer E, Phillips D. 2015. The effect of using different imputation methods for economic variables in aging surveys (Working Paper No. 2015-019). University of Southern California, Center for Economic and Social Research, Los Angeles, CA. https://papers.ssrn.com/sol3/papers.cfm?abstract_id=2650214

Lee J, Phillips D, Wilkens J. 2019. Gateway to Global Aging Data, in: Gu D., Dupre M. (eds) *Encyclopedia of Gerontology and Population Aging*. 2019. Springer, Cham. https://link.springer.com/referenceworkentry/10.1007/978-3-319-69892-2_1105-1

Little RJA. 1988. Missing-data adjustments in large surveys. *Journal of Business & Economic Statistics*, Volume 6, 287-296.
<https://www.jstor.org/stable/1391878>

SHARE. 2024. SHARE Release Guide 9.0.0. SHARE BERLIN Institute, Berlin, Germany.
https://share-eric.eu/fileadmin/user_upload/Release_Guides/SHARE_release_guide_9-0-0.pdf

Wallace, RB, Herzog AR. 1995. Overview of the Health Measures in the Health and Retirement Study. *The Journal of Human Resources* Volume 30 (Supplement 1995):S84-S107.