



This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the authors institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

<http://www.elsevier.com/copyright>



Contents lists available at SciVerse ScienceDirect

Journal of Differential Equations

www.elsevier.com/locate/jde



Blow-up rate and uniqueness of singular radial solutions for a class of quasi-linear elliptic equations

Zhifu Xie^{a,*}, Chunshan Zhao^b

^a Department of Mathematics and Computer Science, Virginia State University, Petersburg, VA 23806, USA

^b Department of Mathematical Sciences, Georgia Southern University, Statesboro, GA 30460, USA

ARTICLE INFO

Article history:

Received 28 March 2011

Revised 16 August 2011

Available online 13 September 2011

Keywords:

Blow-up rate

Large positive solution

Quasi-linear elliptic problem

Uniqueness

ABSTRACT

We establish the uniqueness and the blow-up rate of the large positive solution of the quasi-linear elliptic problem $-\Delta_p u = \lambda u^{p-1} - b(x)h(u)$ in $B_R(x_0)$ with boundary condition $u = +\infty$ on $\partial B_R(x_0)$, where $B_R(x_0)$ is a ball centered at $x_0 \in \mathbb{R}^N$ with radius R , $N \geq 3$, $2 \leq p < \infty$, $\lambda > 0$ are constants and the weight function b is a positive radially symmetrical function. We only require $h(u)$ to be a locally Lipschitz function with $h(u)/u^{p-1}$ increasing on $(0, \infty)$ and $h(u) \sim u^{q-1}$ for large u with $q > p - 1$. Our results extend the previous work [Z. Xie, Uniqueness and blow-up rate of large solutions for elliptic equation $-\Delta u = \lambda u - b(x)h(u)$, J. Differential Equations 247 (2009) 344–363] from case $p = 2$ to case $2 \leq p < \infty$.

© 2011 Elsevier Inc. All rights reserved.

1. Introduction

Let $\Omega \subseteq \mathbb{R}^N$ ($N \geq 2$) be a smooth bounded domain. We consider the uniqueness and the blow-up rate of the large solutions of the quasi-linear elliptic problem with singular boundary value condition as follows:

$$\begin{cases} -\Delta_p u = \lambda u^{p-1} - b(x)h(u) & \text{in } \Omega, & (a) \\ u \geq 0 & \text{in } \Omega, & (b) \\ u = \infty & \text{on } \partial\Omega, & (c) \end{cases} \quad (1.1)$$

* Corresponding author.

E-mail addresses: zxie@vsu.edu (Z. Xie), czhao@GeorgiaSouthern.edu (C. Zhao).

where $\Delta_p u = \operatorname{div}(|\nabla u|^{p-2} \nabla u)$ with $Du(x) = (\partial_{x_1} u, \partial_{x_2} u, \dots, \partial_{x_N} u)$, $p \in [2, \infty)$ and $\lambda > 0$ are constants, $b(x) \geq 0$ is a weighted function which we will explain later, and the boundary condition (1.1c) is understood as $u(x) \rightarrow +\infty$ when $d(x) := \operatorname{dist}(x, \partial\Omega) \rightarrow 0^+$. The solutions of (1.1a–c) are called *large* (or *blow-up*) *solutions*.

The problem (1.1a–c) appears in the study of non-Newtonian flows, chemotaxis, and biological pattern formation etc. For example, in the study of non-Newtonian flows, the constant p in (1.1a) is a characteristic of medium. Media with $p > 2$ are called dilatant fluids and those with $p < 2$ are called pseudo-plastics. If $p = 2$ they are Newtonian fluids (see [8] and the references therein). Especially when $p = 2$, the problem (1.1a–c) becomes as follows:

$$\begin{cases} -\Delta u = \lambda u - b(x)h(u) & \text{in } \Omega, & (a) \\ u \geq 0 & \text{in } \Omega, & (b) \\ u = \infty & \text{on } \partial\Omega, & (c) \end{cases} \quad (1.2)$$

and it has been studied extensively. Next let us recollect some related results. In 1916, Bieberbach [3] studied the large solutions for the particular case $-\Delta u = -\exp(u)$ with conditions (1.2b–c) in smooth bounded two-dimensional domains, and showed that there exists a unique solution such that $u(x) - \log(d(x)^{-2})$ is bounded as $x \rightarrow \partial\Omega$. Problems of this type also arise in Riemannian geometry. For example, if a Riemannian metric of the form $|ds|^2 = \exp(2u(x))|dx|^2$ has constant Gaussian curvature $-c^2$, then $-\Delta u = -c^2 \exp(2u)$. Motivated by a problem in mathematical physics, Rademacher [24] continued the study of the large solutions for the particular case $-\Delta u = -\exp(u)$ on smooth bounded domains in \mathbb{R}^3 . Bandle and Essén [1] and Lazer and McKenna [15] extended Bieberbach's and Rademacher's results to general case $-\Delta u = -b(x)\exp(u)$ in bounded domains of \mathbb{R}^N satisfying a uniform external sphere condition, where the function $b(x)$ is continuous and strictly positive on $\overline{\Omega}$, the closure of Ω . It was shown that the problem has a unique solution together with an estimate of the form $u = \log d^{-2} + o(d)$, see [1] for case $b \equiv 1$ and [15] for case $b(x) \geq b_0 > 0$ as $d \rightarrow 0$.

Recently, the uniqueness of solutions for (1.2a–c) with $h(u) = u^q$ ($q > 1$) on bounded domains or the whole space \mathbb{R}^N was discussed in many papers (see e.g., [10–26]). The results can be summarized as follows: under the assumption

$$\lim_{d(x, \partial\Omega) \rightarrow 0} \frac{b(x)}{d(x, \partial\Omega)^\gamma} = \zeta$$

with $\gamma > 0$ and $\zeta > 0$, an explicit expression for the blow-up rates of solutions of (1.2a–c) was obtained in [10] and [12] as $u = (\alpha(\alpha + 1)/\zeta)^{1/(q-1)} d^{-\alpha} (1 + o(d))$, $\alpha = (\gamma + 2)/(q - 1)$. By using the localization method of [17], it was shown in [17,22] that (1.2a–c) with $h(u) = u^q$, $q > 1$ has at most one blow-up solution for the case when γ and ζ vary along $\partial\Omega$. Further improvements of these results can be found in [5,6,18,21,23,25,26] and the references therein.

The radial case of the problem (1.2a–c) on a ball domain $B_R(x_0)$ with $h(u) = u^q$ was studied by López-Gómez [18], and the author obtained the existence and uniqueness of a solution and also established the exact boundary blow-up rate for less restrictions on the weight function b , which is a positive non-decreasing function with $b(0) = 0$, $b'(r) \geq 0$. The author also extended the results to a general domain by adopting the localization method [17]. Later on, Cano-Casanova and López-Gómez improved the results in [18] for $h(u) = u^q$ to a general function $h(u)$ which satisfies the Keller–Osserman condition [14,20] and $h(u) \sim Hu^q$ ($H > 0$ is a constant and $q > 1$) for sufficiently large u [5,6]. In [21], the authors also considered the problem (1.2a–c) with $h(u) = u^q$ on a ball domain $B_R(x_0)$, but the decay rate of the weight function $b(x)$ was not assumed to be approximated by a distance function near the boundary $\partial\Omega$, i.e., no assumption as $b \sim C_0 d^\gamma + o(d^\gamma)$ or $b(r)$ is a non-decreasing function was needed. They only assumed that $b(x) = b(\|x - x_0\|)$ was a radially symmetric continuous function on a ball $B_R(x_0)$ such that $B(r)/b(r) \in C^1([0, R])$ and $\lim_{r \rightarrow R} B(r)/b(r) = 0$, where $B(r) = \int_r^R b(s) ds$. Uniqueness and blow-up rates of solution of (1.2a–c) in general domains was also

obtained in [22] by combining the localization method with the results in [21]. Also see [25,26] for more results in the direction of general function $h(u)$ in (1.2a–c).

It is often important to know what properties are retained when linear diffusion ($p = 2$) which corresponds to the Laplace operator is replaced by nonlinear diffusion ($p \neq 2$) which corresponds to the degenerate p -Laplace operator. We want to point out that it is not always possible to extend the results from Laplace operator to the degenerate p -Laplace operator (as many examples have already demonstrated); and even if such an extension is possible, one usually has to overcome many non-trivial technical difficulties since many nice properties inherent to the Laplace operator seem lost or difficult to verify once $p \neq 2$. We refer the readers to [9] for the existence of large positive solutions of the problem (1.1a–c). In this paper, we are interested in the uniqueness and the blow-up rate of solutions to the problem (1.1a–c). Our main theorem extends the results obtained in [25] for the problem (1.2a–c) to the case $2 \leq p < \infty$ and can be stated as follows.

Theorem 1.1. *Consider the radially symmetric quasi-linear elliptic equation*

$$\begin{cases} -\Delta_p u = \lambda u^{p-1} - b(\|x - x_0\|)h(u) & \text{in } B_R(x_0), & (a) \\ u \geq 0 & \text{in } B_R(x_0), & (b) \\ u = \infty & \text{on } \partial B_R(x_0), & (c) \end{cases} \quad (1.3)$$

where $B_R(x_0)$ is the ball of radius R centered at $x_0 \in \mathbb{R}^N$, $N \geq 2$, $2 \leq p < \infty$ and $\lambda > 0$. The weight function $b(r)$ satisfies:

(A1) $b \in C([0, R], (0, \infty))$ satisfies $B(r)/b(r) \in C^1([0, R])$, and

$$C_0 := \lim_{r \rightarrow R} \frac{(B(r))^2}{b^*(r)b(r)} \geq 1, \quad (1.4)$$

where $B(r) = \int_r^R b(s) ds$ and $b^*(r) = \int_r^R B(s) ds$. Denote $b_0 = b(R) > 0$.

The nonlinear function $h(u)$ satisfies:

(A2) $h(u) \geq 0$ is locally Lipschitz continuous on $[0, \infty)$ and $h(u)/u^{p-1}$ is increasing on $(0, \infty)$; and, for some $q > p - 1 > 1$,

$$H := \lim_{u \rightarrow \infty} \frac{h(u)}{u^q} > 0. \quad (1.5)$$

Then for any solution $u(x)$ of (1.3a–c),

$$\lim_{d(x) \rightarrow 0} \frac{u(x)}{KH^{-2\beta/p}(b^*(\|x - x_0\|))^{-\beta}} = 1, \quad (1.6)$$

where $d(x) = \text{dist}(x, \partial B_R(x_0))$ and K is a constant defined by

$$K = [(p-1)\beta^{p-1}((\beta+1)C_0 - 1)(b_0C_0)^{p/2-1}]^{2\beta/p}, \quad \beta = \frac{p}{2(q-p+1)}.$$

Therefore problem (1.3a–c) possesses a unique positive large solution $u(x)$ in $B_R(x_0)$.

2. Some preliminary results

At first let us present some lemmas which will be used in the proof of Theorem 1.1. The following lemmas are mainly from [9] with some notation modifications. Similar results for $p = 2$ can also be found in [7,10,9,17,19].

Consider the problem

$$\begin{cases} -\Delta_p u = \lambda u^{p-1} - b(x)h(u) & \text{in } \Omega, \\ u = \phi \geq 0 & \text{on } \partial\Omega, \end{cases} \quad (2.1)$$

where Ω is a smooth bounded domain in \mathbb{R}^N with $N \geq 2$, $\phi \in C^1(\partial\Omega)$, h satisfies (A2) and $b \in C(\Omega, \mathbb{R}^+)$.

Lemma 2.1. *If $\underline{u}, \bar{u} \in C^1(\Omega)$ are both positive in Ω such that*

$$\begin{cases} -\Delta_p \underline{u} \leq \lambda \underline{u}^{p-1} - b(x)h(\underline{u}) & \text{in } \Omega, \\ -\Delta_p \bar{u} \geq \lambda \bar{u}^{p-1} - b(x)h(\bar{u}) & \text{in } \Omega \end{cases} \quad (2.2)$$

hold in the sense of distribution, and $\underline{u} \leq \phi \leq \bar{u}$ on $\partial\Omega$, then $\underline{u}(x) \leq \bar{u}(x)$ on $\bar{\Omega}$.

Remark. We refer the readers to [13] and [16] for maximum and comparison principles for elliptic equations involving p -Laplacian. This lemma can be proved similarly to the proof of Lemma 1.1 in [19] (see also [9] and [4]), that goes back to Benguria, Brezis and Lieb [2]. For convenience, next we give a proof.

Proof of Lemma 2.1. By the hypotheses in (2.2), we have

$$\Delta_p \underline{u} + \lambda \underline{u}^{p-1} - b(x)h(\underline{u}) \geq 0 \quad \text{in } \Omega \quad (2.3)$$

and

$$\Delta_p \bar{u} + \lambda \bar{u}^{p-1} - b(x)h(\bar{u}) \leq 0 \quad \text{in } \Omega. \quad (2.4)$$

Let w_1 and w_2 be nonnegative C^1 functions on Ω and both of them vanish near $\partial\Omega$. Multiplying (2.3) by w_1 and (2.4) by w_2 , and applying integration by parts, and subtraction, we easily obtain

$$\begin{aligned} & - \int_{\Omega} [|\nabla \underline{u}|^{p-2} \nabla \underline{u} \nabla w_1 - |\nabla \bar{u}|^{p-2} \nabla \bar{u} \nabla w_2] dx + \int_{\partial\Omega} \left[|\nabla \underline{u}|^{p-2} \frac{\partial \underline{u}}{\partial \nu} w_1 - |\nabla \bar{u}|^{p-2} \frac{\partial \bar{u}}{\partial \nu} w_2 \right] dS \\ & \geq \int_{\Omega} [b(x)(h(\underline{u})w_1 - h(\bar{u})w_2)] dx - \lambda \int_{\Omega} [\underline{u}^{p-1} w_1 - \bar{u}^{p-1} w_2] dx. \end{aligned} \quad (2.5)$$

Because w_1 and w_2 vanish near $\partial\Omega$,

$$\begin{aligned} & - \int_{\Omega} [|\nabla \underline{u}|^{p-2} \nabla \underline{u} \nabla w_1 - |\nabla \bar{u}|^{p-2} \nabla \bar{u} \nabla w_2] dx \\ & \geq \int_{\Omega} [b(x)(h(\underline{u})w_1 - h(\bar{u})w_2)] dx - \lambda \int_{\Omega} [\underline{u}^{p-1} w_1 - \bar{u}^{p-1} w_2] dx. \end{aligned} \quad (2.6)$$

Let $\epsilon_2 > \epsilon_1 > 0$ and denote

$$v_1 = \left[\frac{(\underline{u} + \epsilon_1)^{2p-2} - (\bar{u} + \epsilon_2)^{2p-2}}{(\underline{u} + \epsilon_1)^{p-1}} \right]^+, \quad (2.7)$$

$$v_2 = \left[\frac{(\underline{u} + \epsilon_1)^{2p-2} - (\bar{u} + \epsilon_2)^{2p-2}}{(\bar{u} + \epsilon_2)^{p-1}} \right]^+. \quad (2.8)$$

Since v_i can be approximated arbitrarily closely by C^1 functions vanishing near $\partial\Omega$, we see that (2.6) holds when w_i is replaced by v_i . Denote

$$\Omega_+(\epsilon_1, \epsilon_2) = \{x \in \Omega \mid \underline{u} + \epsilon_1 > \bar{u} + \epsilon_2\}.$$

We note that the integrands in (2.6) (with $w_i = v_i$) vanish outside this set. Then the left-hand side of (2.6) is

$$\begin{aligned} & - \int_{\Omega_+(\epsilon_1, \epsilon_2)} [|\nabla \underline{u}|^{p-2} \nabla \underline{u} \nabla v_1 - |\nabla \bar{u}|^{p-2} \nabla \bar{u} \nabla v_2] dx \\ &= -(p-1) \int_{\Omega_+(\epsilon_1, \epsilon_2)} \left[|\nabla \underline{u}|^p (\underline{u} + \epsilon_1)^{p-2} \left(1 + \left(\frac{\bar{u} + \epsilon_2}{\underline{u} + \epsilon_1} \right)^{2p-2} \right) \right] dx \\ & \quad + 2(p-1) \int_{\Omega_+(\epsilon_1, \epsilon_2)} \left[\nabla \bar{u} \nabla \underline{u} \left((\bar{u} + \epsilon_2)^{p-2} \left(\frac{\bar{u} + \epsilon_2}{\underline{u} + \epsilon_1} \right)^{p-1} |\nabla \underline{u}|^{p-2} \right) \right] \\ & \quad - (p-1) \int_{\Omega_+(\epsilon_1, \epsilon_2)} \left[|\nabla \bar{u}|^p (\bar{u} + \epsilon_2)^{p-2} \left(1 + \left(\frac{\underline{u} + \epsilon_1}{\bar{u} + \epsilon_2} \right)^{2p-2} \right) \right] dx \\ & \quad + 2(p-1) \int_{\Omega_+(\epsilon_1, \epsilon_2)} \left[\nabla \bar{u} \nabla \underline{u} \left((\underline{u} + \epsilon_1)^{p-2} \left(\frac{\underline{u} + \epsilon_1}{\bar{u} + \epsilon_2} \right)^{p-1} |\nabla \bar{u}|^{p-2} \right) \right] \\ &= -(p-1) \int_{\Omega_+(\epsilon_1, \epsilon_2)} (\underline{u} + \epsilon_1)^{p-2} \left(\frac{\bar{u} + \epsilon_2}{\underline{u} + \epsilon_1} \right)^{2p-2} |\nabla \underline{u}|^{p-2} \left| \nabla \underline{u} - \nabla \bar{u} \left(\frac{\underline{u} + \epsilon_1}{\bar{u} + \epsilon_2} \right) \right|^2 dx \\ & \quad - (p-1) \int_{\Omega_+(\epsilon_1, \epsilon_2)} (\bar{u} + \epsilon_2)^{p-2} \left(\frac{\underline{u} + \epsilon_1}{\bar{u} + \epsilon_2} \right)^{2p-2} |\nabla \bar{u}|^{p-2} \left| \nabla \bar{u} - \nabla \underline{u} \left(\frac{\bar{u} + \epsilon_2}{\underline{u} + \epsilon_1} \right) \right|^2 dx \\ & \quad - (p-1) \int_{\Omega_+(\epsilon_1, \epsilon_2)} (\underline{u} + \epsilon_1)^{p-2} \left(|\nabla \underline{u}|^{p-2} - |\nabla \bar{u}|^{p-2} \left(\frac{\underline{u} + \epsilon_1}{\bar{u} + \epsilon_2} \right)^{p-2} \right) \\ & \quad \times \left(|\nabla \underline{u}|^2 - |\nabla \bar{u}|^2 \left(\frac{\bar{u} + \epsilon_2}{\underline{u} + \epsilon_1} \right)^{2p-4} \right) dx. \end{aligned}$$

Let the above equality be

$$- \int_{\Omega_+(\epsilon_1, \epsilon_2)} [|\nabla \underline{u}|^{p-2} \nabla \underline{u} \nabla v_1 - |\nabla \bar{u}|^{p-2} \nabla \bar{u} \nabla v_2] dx = J_1 + J_2 + J_3.$$

Then the inequality (2.6) becomes

$$J_1 + J_2 + J_3 \geq \int_{\Omega_+(\epsilon_1, \epsilon_2)} \left[b(x) \left(\frac{h(\underline{u})}{(\underline{u} + \epsilon_1)^{p-1}} - \frac{h(\bar{u})}{(\bar{u} + \epsilon_2)^{p-1}} \right) ((\underline{u} + \epsilon_1)^{2p-2} - (\bar{u} + \epsilon_2)^{2p-2}) \right] dx \\ - \lambda \int_{\Omega_+(\epsilon_1, \epsilon_2)} [\underline{u}^{p-1} v_1 - \bar{u}^{p-1} v_2] dx. \quad (2.9)$$

Note that J_1 and J_2 are non-positive. As $\epsilon_2 > \epsilon_1 \rightarrow 0$, the second term on the right-hand side of (2.9) converges to 0 and the first term on the right-hand side of (2.9) converges to

$$\int_{\Omega_+(0,0)} \left[b(x) \left(\frac{h(\underline{u})}{(\underline{u})^{p-1}} - \frac{h(\bar{u})}{(\bar{u})^{p-1}} \right) ((\underline{u})^{2p-2} - (\bar{u})^{2p-2}) \right] dx$$

which is positive unless $\Omega_+(0, 0)$ is empty. The proof of Lemma 2.1 is complete. \square

Next we present the definition of sub-solution and super-solution as follows.

Definition 1. If \underline{u} (resp. \bar{u}) satisfies the conditions in Lemma 2.1 and $\underline{u} \leq \phi$ (resp. $\bar{u} \geq \phi$) on $\partial\Omega$, then \underline{u} (resp. \bar{u}) is called a sub-solution (resp. super-solution) of (2.1).

Lemma 2.2. If $\underline{u}, \bar{u} \in C^1(\Omega)$ are both positive in Ω such that

$$\begin{aligned} -\Delta_p \underline{u} &\leq \lambda \underline{u}^{p-1} - b(x)h(\underline{u}) \quad \text{in } \Omega, \\ -\Delta_p \bar{u} &\geq \lambda \bar{u}^{p-1} - b(x)h(\bar{u}) \quad \text{in } \Omega, \\ \lim_{d(x) \rightarrow 0} \underline{u}(x) &= +\infty, \quad \lim_{d(x) \rightarrow 0} \bar{u}(x) = +\infty, \end{aligned}$$

hold in the sense of distribution, and $\underline{u}(x) \leq \bar{u}(x)$ in Ω , then there exists at least one solution $u \in C^1(\Omega)$ to (2.1) satisfying $\underline{u}(x) \leq u \leq \bar{u}(x)$ in Ω .

We note that $\underline{u} = 0$ is a sub-solution of (2.1), while $\bar{u} = L$ is a super-solution of (2.1) if L is large enough, so (2.1) has at least a solution u_ϕ . Thus the proof of Lemma 2.2 follows exactly the same arguments as in the proof of Theorem 3.2 in López-Gómez [18] and it is omitted for brevity: consider (2.1) in domains $\Omega_n := \{x \in \Omega \mid d(x, \partial\Omega) > \frac{1}{n}\}$ with $u = (\underline{u} + \bar{u})/2$ on $\partial\Omega_n$ and we make $n \rightarrow \infty$ through a diagonal process. The limit of the diagonal sequence provides us with a solution satisfying all the required conditions.

3. Proof of Theorem 1.1

Next we consider the corresponding singular problem in one dimension

$$\begin{cases} -\left[(|\psi'|^{p-2} \psi')' + \frac{N-1}{r} |\psi'|^{p-2} \psi' \right] = \lambda \psi^{p-1} - b(r)h(\psi) & \text{in } (0, R), \\ \lim_{r \rightarrow R} \psi(r) = \infty, \\ \psi'(0) = 0. \end{cases} \quad (3.1)$$

We first show that for each $\epsilon > 0$, problem (3.1) has a positive large solution ψ_ϵ such that

$$1 - \epsilon \leq \liminf_{r \rightarrow R} \frac{\psi_\epsilon(r)}{KH^{-2\beta/p}(b^*(r))^{-\beta}} \leq \limsup_{r \rightarrow R} \frac{\psi_\epsilon(r)}{KH^{-2\beta/p}(b^*(r))^{-\beta}} \leq 1 + \epsilon. \quad (3.2)$$

Therefore, for each $x_0 \in \mathbb{R}^N$, the function

$$u_\epsilon(x) := \psi_\epsilon(r); \quad r := \|x - x_0\|$$

provides us with a radially symmetric positive large solution of (1.3a–c) with the assumptions in Theorem 1.1 and the solution satisfies

$$1 - \epsilon \leq \liminf_{d(x) \rightarrow 0} \frac{u_\epsilon(x)}{KH^{-2\beta/p}(b^*(\|x - x_0\|))^{-\beta}} \leq \limsup_{d(x) \rightarrow 0} \frac{u_\epsilon(x)}{KH^{-2\beta/p}(b^*(\|x - x_0\|))^{-\beta}} \leq 1 + \epsilon. \quad (3.3)$$

In order to prove (3.2), we construct a super-solution and a sub-solution of (3.1). For each $\epsilon > 0$ we claim that

$$\bar{\psi}_\epsilon(r) = A + B_+ \left(\frac{r}{R} \right)^2 (b^*(r))^{-\beta} \quad (3.4)$$

provides us a super-solution, where $A > 0$ and $B_+ > 0$ have to be determined later. Then

$$\begin{aligned} \bar{\psi}'_\epsilon(r) &= 2B_+ \frac{r}{R^2} (b^*(r))^{-\beta} - \beta B_+ \left(\frac{r}{R} \right)^2 (b^*(r))^{-\beta-1} (b^*(r))', \\ \bar{\psi}''_\epsilon(r) &= 2B_+ \frac{1}{R^2} (b^*(r))^{-\beta} - 4\beta B_+ \frac{r}{R^2} (b^*(r))^{-\beta-1} (b^*(r))' \\ &\quad + \beta(\beta + 1)B_+ \left(\frac{r}{R} \right)^2 (b^*(r))^{-\beta-2} [(b^*(r))']^2 - \beta B_+ \left(\frac{r}{R} \right)^2 (b^*(r))^{-\beta-1} (b^*(r))''. \end{aligned}$$

We have $\bar{\psi}_\epsilon(r) \rightarrow \infty$ as $r \rightarrow R$ because $b^*(r) \rightarrow 0$ as $r \rightarrow R$ and $\beta > 0$. Also $\bar{\psi}'_\epsilon(r) \geq 0$ for $r \in [0, R]$ and $\bar{\psi}'_\epsilon(r) \rightarrow 0$ as $r \rightarrow 0$. Then $\bar{\psi}_\epsilon(r)$ is a super-solution if and only if

$$-\left[(|\bar{\psi}'_\epsilon(r)|^{p-2} \bar{\psi}'_\epsilon(r))' + \frac{N-1}{r} |\bar{\psi}'_\epsilon(r)|^{p-2} \bar{\psi}'_\epsilon(r) \right] \geq \lambda \bar{\psi}_\epsilon^{p-1}(r) - b(r)h(\bar{\psi}_\epsilon(r)), \quad (3.5)$$

i.e.

$$-(p-1)(\bar{\psi}'_\epsilon(r))^{p-2} \bar{\psi}''_\epsilon(r) - \frac{N-1}{r} (\bar{\psi}'_\epsilon(r))^{p-1} \geq \lambda \bar{\psi}_\epsilon^{p-1}(r) - b(r)h(\bar{\psi}_\epsilon(r)). \quad (3.6)$$

By the assumption (A2) on h , it is easy to see that for the same $\epsilon > 0$,

$$(1 - \epsilon)H\bar{\psi}_\epsilon^q(r) \leq h(\bar{\psi}_\epsilon(r)) \leq (1 + \epsilon)H\bar{\psi}_\epsilon^q(r) \quad (3.7)$$

for all $r \in [0, R]$ by choosing A sufficiently large, say $A \geq A_0$. The inequality (3.6) holds if

$$-(p-1)(\bar{\psi}'_\epsilon(r))^{p-2} \bar{\psi}''_\epsilon(r) - \frac{N-1}{r} (\bar{\psi}'_\epsilon(r))^{p-1} \geq \lambda \bar{\psi}_\epsilon^{p-1}(r) - b(r)(1 - \epsilon)H\bar{\psi}_\epsilon^q(r). \quad (3.8)$$

Note that $(b^*(r))' = -B(r)$, $(b^*(r))'' = b(r)$. The inequality (3.8) is equivalent to

$$\begin{aligned}
 & -(p-1) \left[2B_+ \frac{r}{R^2} (b^*(r))^{-\beta} + \beta B_+ \left(\frac{r}{R} \right)^2 (b^*(r))^{-\beta-1} B(r) \right]^{p-2} \cdot \left[2B_+ \frac{1}{R^2} (b^*(r))^{-\beta} \right. \\
 & \quad \left. + 4\beta B_+ \frac{r}{R^2} (b^*(r))^{-\beta-1} B(r) + \beta(\beta+1) B_+ \left(\frac{r}{R} \right)^2 (b^*(r))^{-\beta-2} (B(r))^2 \right. \\
 & \quad \left. - \beta B_+ \left(\frac{r}{R} \right)^2 (b^*(r))^{-\beta-1} b(r) \right] - \frac{N-1}{r} \left[2B_+ \frac{r}{R^2} (b^*(r))^{-\beta} + \beta B_+ \left(\frac{r}{R} \right)^2 (b^*(r))^{-\beta-1} B(r) \right]^{p-1} \\
 & \geq \lambda \left[A + B_+ \left(\frac{r}{R} \right)^2 (b^*(r))^{-\beta} \right]^{p-1} - b(r)(1-\epsilon)H \left[A + B_+ \left(\frac{r}{R} \right)^2 (b^*(r))^{-\beta} \right]^q.
 \end{aligned}$$

Multiplying both sides of this inequality by $\frac{(b^*(r))^{\beta+1}(p-1)+1}{(B(r))^p}$ and taking into consideration that $\beta = \frac{p}{2(q-p+1)}$, we have

$$\begin{aligned}
 & -(p-1) \left[2B_+ \frac{r}{R^2} \frac{b^*(r)}{B(r)} + \beta B_+ \left(\frac{r}{R} \right)^2 \right]^{p-2} \cdot \left[2B_+ \frac{1}{R^2} \left(\frac{b^*(r)}{B(r)} \right)^2 + 4\beta B_+ \frac{r}{R^2} \frac{b^*(r)}{B(r)} \right. \\
 & \quad \left. + \beta(\beta+1) B_+ \left(\frac{r}{R} \right)^2 - \beta B_+ \left(\frac{r}{R} \right)^2 \frac{b^*(r)b(r)}{B^2(r)} \right] \\
 & \quad - \frac{N-1}{r} \frac{b^*(r)}{B(r)} \left[2B_+ \frac{r}{R^2} \frac{b^*(r)}{B(r)} + \beta B_+ \left(\frac{r}{R} \right)^2 \right]^{p-1} \\
 & \geq \lambda \left(\frac{b^*(r)}{B(r)} \right)^p \left[A(b^*(r))^\beta + B_+ \left(\frac{r}{R} \right)^2 \right]^{p-1} \\
 & \quad - b(r)(b^*(r))^{-\beta p} \frac{(b^*(r))^{\beta+1}(p-1)+1}{(B(r))^p} (1-\epsilon)H \left[A(b^*(r))^\beta + B_+ \left(\frac{r}{R} \right)^2 \right]^q \\
 & = \lambda \left(\frac{b^*(r)}{B(r)} \right)^p \left[A(b^*(r))^\beta + B_+ \left(\frac{r}{R} \right)^2 \right]^{p-1} \\
 & \quad - b^{1-p/2}(r) \left(\frac{b^*(r)b(r)}{B^2(r)} \right)^{p/2} (1-\epsilon)H \left[A(b^*(r))^\beta + B_+ \left(\frac{r}{R} \right)^2 \right]^q.
 \end{aligned}$$

Since when $r \rightarrow R$, $b^*(r) \rightarrow 0$, $\frac{b^*(r)}{B(r)} \rightarrow 0$, $\frac{(B(r))^2}{b^*(r)b(r)} \rightarrow C_0 \geq 1$ by the assumption (A1), then as $r \rightarrow R$,

$$-(p-1)(\beta B_+)^{p-2} \left[\beta(\beta+1)B_+ - \beta B_+ \frac{1}{C_0} \right] \geq -b_0^{1-p/2} C_0^{-p/2} (1-\epsilon)H B_+^p,$$

which is

$$B_+ \geq [(p-1)\beta^{p-1}((\beta+1)C_0-1)(b_0C_0)^{p/2-1}]^{\frac{1}{q-p+1}} [(1-\epsilon)H]^{\frac{-1}{q-p+1}}.$$

Let $B_+ = (1+\epsilon)[(1-\epsilon)H]^{-2\beta/p}[(p-1)\beta^{p-1}((\beta+1)C_0-1)(b_0C_0)^{p/2-1}]^{2\beta/p} = (1+\epsilon)(1-\epsilon)^{-2\beta/p}H^{-2\beta/p}K$. Therefore, by making the choice B_+ , the inequality (3.8) is satisfied in a left neighborhood of $r = R$, say $(R-\delta, R]$, for some $\delta = \delta(\epsilon) > 0$. Finally, by choosing A sufficiently large (larger than A_0) it is clear that the inequality is satisfied in the whole interval $[0, R]$ since $q > p-1$ and $b^*(r)$ is bounded away from zero in $[0, R-\delta]$. Then $\bar{\psi}_\epsilon$ is our required super-solution of problem (3.1).

Next, we construct a sub-solution with the same blow-up rate as the above super-solution. Due to the assumption (A2) on h , for $u \geq A_0$ large,

$$(1 - \epsilon)Hu^q \leq h(u) \leq (1 + \epsilon)Hu^q.$$

For given $A_0 > 0$ and $0 < R_0 < R$, we consider the auxiliary problem

$$\begin{cases} -\left[(|\psi'|^{p-2}\psi')' + \frac{N-1}{r}|\psi'|^{p-2}\psi' \right] = \lambda\psi^{p-1} - b(r)h(\psi) & \text{in } (0, R_0), \\ \psi(R_0) = A_0, \\ \psi'(0) = 0. \end{cases} \quad (3.9)$$

By the assumptions on b and h , we have

$$\min_{r \in [0, R_0]} b(r) > 0, \quad h(0) = 0, \quad \text{and} \quad h(u)/u \rightarrow \infty \quad \text{as } u \rightarrow \infty.$$

Then it is easy to know that

$$\underline{\psi}_{A_0} := 0, \quad \overline{\psi}_{A_0} := A_0$$

provides us with an ordered sub-super-solution pair of (3.9). Thus (3.9) possesses a solution ψ_{A_0} such that $\psi_{A_0}(r) \in [0, A_0]$ for all $r \in [0, R_0]$.

For each $\epsilon > 0$ sufficiently small, we claim that there exists $0 < C < A_0$ for which the function

$$\underline{\psi}_\epsilon(r) = \begin{cases} \psi_{A_0}(r), & r \in [0, R_0], \\ \max\{A_0, C + B_-(\frac{r}{R})^2(b^*(r))^{-\beta}\}, & r \in (R_0, R], \end{cases}$$

provides a sub-solution, where R_0 and C are to be determined later and

$$B_- = (1 - \epsilon)(1 + \epsilon)^{-2\beta/p} H^{-2\beta/p} K.$$

In fact, denoting $f_C(r) = C + B_-(\frac{r}{R})^2(b^*(r))^{-\beta}$ we have

$$\begin{aligned} f'_C(r) &= 2B_- \frac{r}{R^2} (b^*(r))^{-\beta} - \beta B_- \left(\frac{r}{R}\right)^2 (b^*(r))^{-\beta-1} (b^*(r))' \\ &= 2B_- \frac{r}{R^2} (b^*(r))^{-\beta} + \beta B_- \left(\frac{r}{R}\right)^2 (b^*(r))^{-\beta-1} B(r) \end{aligned}$$

which is strictly bigger than zero in $(0, R)$. It follows that $f_C(r)$ is increasing and

$$\lim_{r \rightarrow R} f_C(r) = +\infty, \quad \lim_{r \rightarrow 0} f_C(r) = C < A_0.$$

By the continuity of $f_C(r)$ and the intermediate-value theorem, there exists a unique $Z = Z(C) \in (0, R)$ such that

$$\begin{aligned} C + B_- \left(\frac{r}{R}\right)^2 (b^*(r))^{-\beta} &< A_0 \quad \text{when } r \in [0, Z(C)), \\ C + B_- \left(\frac{r}{R}\right)^2 (b^*(r))^{-\beta} &\geq A_0 \quad \text{when } r \in [Z(C), R]. \end{aligned}$$

Moreover, $Z(C)$ is decreasing and

$$\lim_{C \rightarrow -\infty} Z(C) = R, \quad \lim_{C \rightarrow A_0} Z(C) = 0.$$

Let $R_0 = Z(C)$. From the definition of $\underline{\psi}_\epsilon(r)$ and R_0 , $\underline{\psi}_\epsilon(r) \equiv \psi_{A_0}(r)$ in $[0, Z(C)]$, and then the inequality $-(p-1)(\underline{\psi}'_\epsilon(r))^{p-2}\underline{\psi}''_\epsilon(r) - \frac{N-1}{r}(\underline{\psi}'_\epsilon(r))^{p-1} \leq \lambda \underline{\psi}_\epsilon^{p-1}(r) - b(r)h(\underline{\psi}_\epsilon(r))$ holds in $[0, Z(C)]$. So $\underline{\psi}_\epsilon(r)$ is a sub-solution of (3.1) if the following inequality is satisfied in $[Z(C), R]$

$$-(p-1)(\underline{\psi}'_\epsilon(r))^{p-2}\underline{\psi}''_\epsilon(r) - \frac{N-1}{r}(\underline{\psi}'_\epsilon(r))^{p-1} \leq \lambda \underline{\psi}_\epsilon^{p-1}(r) - b(r)h(\underline{\psi}_\epsilon(r)). \quad (3.10)$$

For each $r \in [Z(C), R]$,

$$-\frac{N-1}{r}(\underline{\psi}'_\epsilon(r))^{p-1} \leq 0 \leq \lambda \underline{\psi}_\epsilon^{p-1}(r).$$

By using the fact $h(\underline{\psi}_\epsilon(r)) \leq (1+\epsilon)H\underline{\psi}_\epsilon(r)^p$ in $[Z(C), R]$, (3.10) holds if

$$-(p-1)(\underline{\psi}'_\epsilon(r))^{p-2}\underline{\psi}''_\epsilon(r) \leq -b(r)(1+\epsilon)H\underline{\psi}_\epsilon^q(r), \quad \text{for each } r \in [Z(C), R],$$

i.e.

$$\begin{aligned} & -(p-1) \left[2B_- - \frac{r}{R^2} \frac{b^*(r)}{B(r)} + \beta B_- \left(\frac{r}{R} \right)^2 \right]^{p-2} \cdot \left[2B_- - \frac{1}{R^2} \left(\frac{b^*(r)}{B(r)} \right)^2 + 4\beta B_- \frac{r}{R^2} \frac{b^*(r)}{B(r)} \right. \\ & \quad \left. + \beta(\beta+1)B_- \left(\frac{r}{R} \right)^2 - \beta B_- \left(\frac{r}{R} \right)^2 \frac{b^*(r)b(r)}{B^2(r)} \right] \\ & \leq -b^{1-p/2}(r) \left(\frac{b^*(r)b(r)}{B^2(r)} \right)^{p/2} (1+\epsilon)H \left[C(b^*(r))^\beta + B_- \left(\frac{r}{R} \right)^2 \right]^q. \end{aligned}$$

Taking $r \rightarrow R$, it becomes

$$-(p-1)(\beta B_-)^{p-2} \left[\beta(\beta+1)B_- - \beta B_- \frac{1}{C_0} \right] \leq -b_0^{1-p/2} C_0^{-p/2} (1+\epsilon)H B_-^p,$$

which is

$$B_- \leq \left[(p-1)\beta^{p-1}((\beta+1)C_0 - 1)(b_0 C_0)^{p/2-1} \right]^{\frac{1}{q-p+1}} [(1+\epsilon)H]^{\frac{-1}{q-p+1}}.$$

Let $B_- = (1-\epsilon)[(1+\epsilon)H]^{-2\beta/p}[(p-1)\beta^{p-1}((\beta+1)C_0 - 1)(b_0 C_0)^{p/2-1}]^{2\beta/p} = (1-\epsilon)(1+\epsilon)^{-2\beta/p} H^{-2\beta/p} K$. It is easy to see that a constant $\delta = \delta(\epsilon) > 0$ exists for which the inequality (3.10) is satisfied in $[R-\delta, R]$, then we choose C such that $Z(C) = R - \delta(\epsilon)$ (therefore $R_0 = R - \delta(\epsilon)$). For this choice of C , it readily follows that $\underline{\psi}_\epsilon$ is a sub-solution to the problem.

So we have constructed a sub-solution and a super-solution with the same blow-up rate of problem (3.1). Because $\bar{\psi}_\epsilon(r) \geq \underline{\psi}_\epsilon(r)$ in $[0, R]$ and $\lim_{r \rightarrow R} \bar{\psi}_\epsilon(r) = \lim_{r \rightarrow R} \underline{\psi}_\epsilon(r) = \infty$, then by Lemma 2.2 there exists a solution $\Psi_\epsilon(r)$ of (3.1) such that

$$1 - \epsilon \leq \liminf_{r \rightarrow R} \frac{\Psi_\epsilon(r)}{KH^{-2\beta/p}(b^*(r))^{-\beta}} \leq \limsup_{r \rightarrow R} \frac{\Psi_\epsilon(r)}{KH^{-2\beta/p}(b^*(r))^{-\beta}} \leq 1 + \epsilon.$$

Proof of uniqueness. The proof of uniqueness basically follows the proofs in [10,12,25]. Let u be an arbitrary solution of (1.3a–c) with assumptions on nonlinear function $h(u)$ and weight function b as in Theorem 1.1. We first show that

$$\lim_{d(x) \rightarrow 0} \frac{u(x)}{KH^{-2\beta/p}(b^*(\|x - x_0\|))^{-\beta}} = 1.$$

Consequently, for any pair of solutions u, v of (1.3a–c)

$$\lim_{d(x) \rightarrow 0} \frac{u(x)}{v(x)} = 1.$$

In doing so, for any $\epsilon > 0$, there exists a radially symmetric positive large solution u_ϵ of (1.3a–c) satisfying (3.3). Choose $0 < \delta < \frac{R}{3}$ small, fix $0 < \tau < \frac{\delta}{4}$ and introduce the region

$$Q_\tau := \left\{ x \mid \tau < d(x, \partial B_R(x_0)) < \frac{\delta}{2} \right\}.$$

Let $M \geq \max_{\|x - x_0\| \leq (R - \frac{\delta}{4})} u(x)$ be large. Thus for every $\tau \in (0, \frac{\delta}{4})$,

$$\bar{V}_\epsilon(x) = u_\epsilon \left(x + \tau \frac{(x - x_0)}{\|x - x_0\|} \right) + M = u_\epsilon(\|x - x_0\| + \tau) + M$$

is a super-solution to

$$\begin{cases} -\Delta_p v = \lambda v^{p-1} - bh(v) & \text{in } Q_\tau, \\ v = u & \text{on } \partial Q_\tau \end{cases} \quad (3.11)$$

with u an arbitrary fixed solution to (1.3a–c) since $\bar{V}_\epsilon(x) \geq u$ for $x \in \partial Q_\tau, \tau \in (0, \frac{\delta}{4})$. Note that $\bar{V}_\epsilon(x) \rightarrow \infty$ as $x \rightarrow \partial B_{R-\tau}(x_0)$. $\bar{V}_\epsilon(x) \geq M \geq u$ as $x \rightarrow \partial B_{R-\frac{\delta}{2}}(x_0)$. In addition, the auxiliary problem (3.11) has $v = u$ as its unique solution. Since 0 is a sub-solution ($h(0) = 0$ by the assumption $h(u)/u$ is increasing), we conclude $u(x) \leq \bar{V}_\epsilon(x) = u_\epsilon(\|x - x_0\| + \tau) + M$ for every $x \in Q_\tau, 0 < \tau < \frac{\delta}{4}$. Letting $\tau \rightarrow 0^+$, we arrive at $u(x) \leq u_\epsilon(x) + M$ for every $x \in A_{\frac{\delta}{2}, R}(x_0)$ and we obtain

$$\limsup_{d(x) \rightarrow 0} \frac{u(x)}{KH^{-2\beta/p}(b^*(\|x - x_0\|))^{-\beta}} \leq \limsup_{d(x) \rightarrow 0} \frac{u_\epsilon(x)}{KH^{-2\beta/p}(b^*(\|x - x_0\|))^{-\beta}} \leq 1 + \epsilon.$$

We now prove

$$1 - \epsilon \leq \liminf_{d(x) \rightarrow 0} \frac{u_\epsilon(x)}{KH^{-2\beta/p}(b^*(\|x - x_0\|))^{-\beta}} \leq \liminf_{d(x) \rightarrow 0} \frac{u}{KH^{-2\beta/p}(b^*(\|x - x_0\|))^{-\beta}}.$$

For the same $\epsilon > 0$ and the radially symmetric positive large solution u_ϵ of (1.3a–c) satisfying (3.3), we choose $0 < \delta < \frac{R}{3}$ small, fix $0 < \tau < \frac{\delta}{4}$ and introduce the annuli region

$$A_{R-\delta, R-\tau} = \{x: R - \delta < \|x - x_0\| < R - \tau\}.$$

Let $M_1 \geq \max_{\|x-x_0\| \leq R-\frac{3\delta}{4}} u(x)$ be large. Thus for every $\tau \in (0, \frac{\delta}{4})$,

$$\tilde{V}_\epsilon(x) = \max \left\{ u \left(x + \tau \frac{(x-x_0)}{\|x-x_0\|} \right) - M_1, 0 \right\} = \max \{ u(\|x-x_0\| - \tau) - M_1, 0 \}$$

is a sub-solution to

$$\begin{cases} -\Delta_p v = \lambda v^{p-1} - bh(v) & \text{in } A_{R-\delta, R-\tau}, \\ v = u_\epsilon & \text{on } \partial A_{R-\delta, R-\tau} \end{cases} \quad (3.12)$$

for all $\tau \in (0, \frac{\delta}{4})$. By the same argument as above, we obtain

$$1 - \epsilon \leq \liminf_{d(x) \rightarrow 0} \frac{u_\epsilon}{KH^{-2\beta/p}(b^*(\|x-x_0\|))^{-\beta}} \leq \liminf_{d(x) \rightarrow 0} \frac{u}{KH^{-2\beta/p}(b^*(\|x-x_0\|))^{-\beta}}.$$

Then

$$1 - \epsilon \leq \liminf_{d(x) \rightarrow 0} \frac{u}{KH^{-2\beta/p}(b^*(\|x-x_0\|))^{-\beta}} \leq \limsup_{d(x) \rightarrow 0} \frac{u}{KH^{-2\beta/p}(b^*(\|x-x_0\|))^{-\beta}} \leq 1 + \epsilon.$$

Letting $\epsilon \rightarrow 0^+$, we have

$$\lim_{d(x) \rightarrow 0} \frac{u}{KH^{-2\beta/p}(b^*(\|x-x_0\|))^{-\beta}} = 1.$$

Now let u and v be large positive solutions to (1.3a–c). By virtue of above, u and v satisfy $\lim_{d(x) \rightarrow 0} \frac{u}{v} = 1$. Thus, for every $\epsilon > 0$, we can find $\delta > 0$ (as small as we are please) such that

$$(1 - \epsilon)v(x) \leq u(x) \leq (1 + \epsilon)v(x)$$

when $0 < d(x) \leq \delta$. On the other hand $\underline{w} = (1 - \epsilon)v(x)$ and $\overline{w} = (1 + \epsilon)v(x)$ are a sub-solution and a super-solution respectively to

$$\begin{cases} -\Delta_p w = \lambda w^{p-1} - bh(w) & \text{in } B_{R-\delta}(x_0), \\ w = u & \text{on } \partial B_{R-\delta}(x_0), \end{cases} \quad (3.13)$$

where we use the property $\frac{h((1-\epsilon)v)}{1-\epsilon} \leq h(v)$ and $\frac{h((1+\epsilon)v)}{1+\epsilon} \geq h(v)$. The unique solution to this problem is $w = u$. Then

$$(1 - \epsilon)v(x) \leq u(x) \leq (1 + \epsilon)v(x)$$

holds in $B_{R-\delta}(x_0)$, therefore it is true in $B_R(x_0)$. Letting $\epsilon \rightarrow 0$ we arrive at $u = v$.

References

- [1] C. Bandle, M. Essèn, On the solutions of quasilinear elliptic problems with boundary blow-up, *Symp. Math.* 35 (1994) 93–111.
- [2] R. Benguria, H. Brezis, E. Lieb, The Thomas–Fermi–Von Weizsacker theory of atoms and molecules, *Comm. Math. Phys.* 79 (1981) 167–180.
- [3] L. Bieberbach, $\Delta u = e^u$ und die automorphen Funktionen, *Math. Ann.* 77 (1916) 173–212.
- [4] M.-M. Boureau, Uniqueness of singular radial solutions for a class of quasilinear problems, *Bull. Belg. Math. Soc. Simon Stevin* 16 (4) (2009) 665–685.

- [5] S. Cano-Casanova, J. López-Gómez, Existence, uniqueness and blow-up rate of large solutions for a canonical class of one-dimensional problems on the half-line, *J. Differential Equations* 244 (2008) 3180–3203.
- [6] S. Cano-Casanova, J. López-Gómez, Blow-up rates of radially symmetric large solutions, *J. Math. Anal. Appl.* 352 (2009) 166–174.
- [7] F.C. Cirstea, Y. Du, General uniqueness results and variation speed for blow-up solutions of elliptic equations, *Proc. London Math. Soc.* (3) 91 (2005) 459–482.
- [8] J. Díaz, *Nonlinear Partial Differential Equations and Free Boundaries*, vol. I: Elliptic Equations, Res. Notes Math., vol. 106, Pitman Advanced Publishing Program, Boston, 1985.
- [9] Y. Du, Z.M. Guo, Boundary blow-up solutions and their applications in quasilinear elliptic equations, *J. Anal. Math.* 89 (2003) 277–302.
- [10] Y. Du, Q. Huang, Blow-up solutions for a class of semilinear elliptic and parabolic equations, *SIAM J. Math. Anal.* 31 (1) (1999) 1–18.
- [11] J. García-Melián, R. Gómez-Reñasco, J. López-Gómez, J.C. Sabina De Lis, Point-wise growth and uniqueness of positive solutions for a class of sublinear elliptic problems where bifurcation from infinity occurs, *Arch. Ration. Mech. Anal.* 145 (1998) 261–289.
- [12] J. García-Melián, R. Letelier-Albornoz, J. Sabina De Lis, Uniqueness and asymptotic behaviour for solutions of semilinear problems with boundary blow-up, *Proc. Amer. Math. Soc.* 129 (12) (2001) 3593–3602.
- [13] J. García-Melián, J. Sabina De Lis, Maximum and comparison principles for operators involving the p -Laplacian, *J. Math. Anal. Appl.* 218 (1998) 49–65.
- [14] J.B. Keller, On solutions of $\Delta u = f(u)$, *Comm. Pure Appl. Math.* 10 (1957) 503–510.
- [15] A.C. Lazer, P.J. McKenna, On a problem of Bieberbach and Rademacher, *Nonlinear Anal.* 21 (1993) 327–335.
- [16] J. López-Gómez, M. Molina-Meyer, The maximum principle for cooperative weakly coupled elliptic systems and some applications, *Differential Integral Equations* 7 (1994) 383–398.
- [17] J. López-Gómez, The boundary blow-up rate of large solutions, *J. Differential Equations* 195 (2003) 25–45.
- [18] J. López-Gómez, Optimal uniqueness theorems and exact blow-up rates of large solutions, *J. Differential Equations* 224 (2006) 385–439.
- [19] M. Marcus, L. Véron, Uniqueness and asymptotic behavior of solutions with boundary blow-up for a class of nonlinear elliptic equations, *Ann. Inst. H. Poincaré Anal. Non Linéaire* 14 (1997) 237–274.
- [20] R. Osserman, On the inequality $\Delta u \geq f(u)$, *Pacific J. Math.* 7 (1957) 1641–1647.
- [21] T. Ouyang, Z. Xie, The uniqueness of blow-up solution for radially symmetric semilinear elliptic equation, *Nonlinear Anal.* 64 (9) (2006) 2129–2142.
- [22] T. Ouyang, Z. Xie, The exact boundary blow-up rate of large solutions for semilinear elliptic problems, *Nonlinear Anal.* 68 (2008) 2791–2800.
- [23] F. Peng, Remarks on large solutions of a class of semilinear elliptic equations, *J. Math. Anal. Appl.* 356 (2009) 393–404.
- [24] H. Rademacher, Einige besondere Probleme der partiellen Differentialgleichungen, in: P. Frank, R. Von Mises (Eds.), *Die Differential und Integralgleichungen der Mechanik und Physik I*, 2nd. edition, Rosenberg, New York, 1943, pp. 838–845.
- [25] Z. Xie, Uniqueness and blow-up rate of large solutions for elliptic equation $-\Delta u = \lambda u - b(x)h(u)$, *J. Differential Equations* 247 (2009) 344–363.
- [26] Z. Xie, General uniqueness results and examples for blow-up solutions of elliptic equations, *Discrete Contin. Dyn. Syst. (Suppl.)* (2009) 828–837.