

Linear models

Samraat Pawar

Department of Life Sciences (Silwood Park)

Imperial College
London

LECTURE OUTLINE

Topics:

- What is a linear model?
 - Regression
 - ANOVA
 - Multiple explanatory variables (ANCOVA)
- Fitting linear models to your data
- Is the fitted linear model appropriate for the data?
- How well does a fitted linear model explain the data?

LECTURE OUTLINE

Topics:

- What is a linear model?
 - Regression
 - ANOVA
 - Multiple explanatory variables (ANCOVA)
- Fitting linear models to your data
- Is the fitted linear model appropriate for the data?
- How well does a fitted linear model explain the data?

Concepts:

- Types of variable: continuous versus categorical
- Terms and coefficients of a model
- Model fitting and model residuals
- Significance testing and p-values

WHAT PREDICTS THE WEIGHTS (w) OF LECTURERS?

Use *intuition* and *prior knowledge* to identify the *variables* to collect...

- Height (h) in metres

WHAT PREDICTS THE WEIGHTS (w) OF LECTURERS?

Use *intuition* and *prior knowledge* to identify the *variables* to collect...

- Height (h) in metres
- Exercise per week (e) in hours

WHAT PREDICTS THE WEIGHTS (w) OF LECTURERS?

Use *intuition* and *prior knowledge* to identify the *variables* to collect...

- Height (h) in metres
- Exercise per week (e) in hours
- Gender (g)

WHAT PREDICTS THE WEIGHTS (w) OF LECTURERS?

Use *intuition* and *prior knowledge* to identify the *variables* to collect...

- Height (h) in metres
- Exercise per week (e) in hours
- Gender (g)
- Distance from home to nearest Greggs bakery (d) in metres

WHAT PREDICTS THE WEIGHTS (w) OF LECTURERS?

Use *intuition* and *prior knowledge* to identify the *variables* to collect...

- Height (h) in metres
- Exercise per week (e) in hours
- Gender (g)
- Distance from home to nearest Greggs bakery (d) in metres
- Ownership of a games console (c)

WHAT PREDICTS THE WEIGHTS (w) OF LECTURERS?

Use *intuition* and *prior knowledge* to identify the *variables* to collect...

- Height (h) in metres
- Exercise per week (e) in hours
- Gender (g)
- Distance from home to nearest Greggs bakery (d) in metres
- Ownership of a games console (c)

WHAT PREDICTS THE WEIGHTS (w) OF LECTURERS?

Use *intuition* and *prior knowledge* to identify the *variables* to collect...

- Height (h) in metres
- Exercise per week (e) in hours
- Gender (g)
- Distance from home to nearest Greggs bakery (d) in metres
- Ownership of a games console (c)

... and build a mathematical model:

Lecturer weight (w) = *Combination of Independent Variables* (that determine w)

WHAT PREDICTS THE WEIGHTS (w) OF LECTURERS?

Use *intuition* and *prior knowledge* to identify the *variables* to collect...

- Height (h) in metres
- Exercise per week (e) in hours
- Gender (g)
- Distance from home to nearest Greggs bakery (d) in metres
- Ownership of a games console (c)


... and build a mathematical model:

Lecturer weight (w) = *Combination of Independent Variables* (that determine w)

$$w = \beta_1 + \beta_2 h + \beta_3 e + \beta_4 g_m + \beta_5 d + \beta_6 c_s + \beta_7 c_a + \varepsilon$$


THE LINEAR MODEL

A combination of four components:




- A **response variable** (w)
- A set of **explanatory variables** (h, e, g, d, c)
- A set of **coefficients** ($\beta_1 - \beta_7$)
- A set of **residuals** (ϵ)

THE VARIABLES




- The response variable is always continuous.

THE VARIABLES




- The response variable is always **continuous**.
- The explanatory variables can be a mix of:

THE VARIABLES




- The response variable is always continuous.
- The explanatory variables can be a mix of:
 - Continuous variables: height, exercise and distance.

THE VARIABLES




- The response variable is always **continuous**.
- The explanatory variables can be a mix of:
 - **Continuous** variables: height, exercise and distance.
 - **Categorical** variables: gender and console ownership.

THE VARIABLES




- The response variable is always **continuous**.
- The explanatory variables can be a mix of:
 - **Continuous** variables: height, exercise and distance.
 - **Categorical** variables: gender and console ownership.
- **Categorical** variables or *factors* have a number of *levels*:

THE VARIABLES



- The response variable is always **continuous**.
- The explanatory variables can be a mix of:
 - **Continuous** variables: height, exercise and distance.
 - **Categorical** variables: gender and console ownership.
- **Categorical** variables or *factors* have a number of *levels*:
 - Gender has two levels (Male / Female)

THE VARIABLES



- The response variable is always **continuous**.
- The explanatory variables can be a mix of:
 - **Continuous** variables: height, exercise and distance.
 - **Categorical** variables: gender and console ownership.
- **Categorical** variables or *factors* have a number of *levels*:
 - Gender has two levels (Male / Female)
 - Console has three levels (None / Sofa-based / Active)

THE TERMS AND COEFFICIENTS

The diagram illustrates the components of a linear regression model. The equation is $w = \beta_1 + \beta_2 h + \beta_3 e + \beta_4 g_m + \beta_5 d + \beta_6 c_s + \beta_7 c_a + \varepsilon$. Explanatory variables are shown in blue boxes with arrows pointing to their respective terms: Height points to h , Exercise points to e , and Distance points to d . These terms are in blue. Other terms are in red: g_m (Gender), c_s (Console), and c_a (Console). The coefficients $\beta_1, \beta_2, \beta_3, \beta_5, \beta_6, \beta_7$ and the error term ε are in grey.

$$w = \beta_1 + \beta_2 h + \beta_3 e + \beta_4 g_m + \beta_5 d + \beta_6 c_s + \beta_7 c_a + \varepsilon$$

- Each explanatory variable is a *term* in the model

THE TERMS AND COEFFICIENTS

The diagram shows a linear regression equation: $w = \beta_1 + \beta_2 h + \beta_3 e + \beta_4 g_m + \beta_5 d + \beta_6 c_s + \beta_7 c_a + \varepsilon$. The terms are color-coded: h (blue), e (blue), g_m (red), d (blue), c_s (red), and c_a (red). Above the equation, three red boxes labeled "Gender" and "Console" have red arrows pointing to the red terms g_m , c_s , and c_a respectively. Below the equation, three blue boxes labeled "Height", "Exercise", and "Distance" have blue arrows pointing to the blue terms h , e , and d respectively.

$$w = \beta_1 + \beta_2 h + \beta_3 e + \beta_4 g_m + \beta_5 d + \beta_6 c_s + \beta_7 c_a + \varepsilon$$

- Each explanatory variable is a *term* in the model
- Each term has at least one coefficient

THE TERMS AND COEFFICIENTS

The diagram illustrates a linear regression equation: $w = \beta_1 + \beta_2 h + \beta_3 e + \beta_4 g_m + \beta_5 d + \beta_6 c_s + \beta_7 c_a + \varepsilon$. The terms are categorized as follows:

- Continuous terms (blue boxes):** Height (points to h), Exercise (points to e), and Distance (points to d).
- Categorical terms (red boxes):** Gender (points to g_m) and Console (points to both c_s and c_a).

Each term is associated with a coefficient β_i . The error term ε is also shown.

- Each explanatory variable is a *term* in the model
- Each term has at least one coefficient
- **Continuous** terms always have one coefficient

THE TERMS AND COEFFICIENTS

The diagram shows the equation $w = \beta_1 + \beta_2 h + \beta_3 e + \beta_4 g_m + \beta_5 d + \beta_6 c_s + \beta_7 c_a + \varepsilon$. Above the equation, blue boxes labeled 'Height', 'Exercise', and 'Distance' have arrows pointing to the variables h , e , and d respectively. Above the equation, red boxes labeled 'Gender' and 'Console' have arrows pointing to the terms g_m and c_s and c_a respectively. The variables h , e , and d are blue, while g_m , c_s , and c_a are red.

$$w = \beta_1 + \beta_2 h + \beta_3 e + \beta_4 g_m + \beta_5 d + \beta_6 c_s + \beta_7 c_a + \varepsilon$$

- Each explanatory variable is a *term* in the model
- Each term has at least one coefficient
- **Continuous** terms always have one coefficient
- Categorical **Factors** have $N - 1$ coefficients, where N is the number of levels (*where are the missing coefficients??*)

WAIT! WHY $N - 1$ COEFFICIENTS? WHAT IS β_1 ?

$$w = \beta_1 + \beta_2 h + \beta_3 e + \beta_4 g_m + \beta_5 d + \beta_6 c_s + \beta_7 c_a + \varepsilon$$

- Two ways of thinking about β_1 :

WAIT! WHY $N - 1$ COEFFICIENTS? WHAT IS β_1 ?

$$w = \beta_1 + \beta_2 h + \beta_3 e + \beta_4 g_m + \beta_5 d + \beta_6 c_s + \beta_7 c_a + \varepsilon$$

- Two ways of thinking about β_1 :
 - Continuous variables: the *y intercept*

WAIT! WHY $N - 1$ COEFFICIENTS? WHAT IS β_1 ?

$$w = \beta_1 + \beta_2 h + \beta_3 e + \beta_4 g_m + \beta_5 d + \beta_6 c_s + \beta_7 c_a + \varepsilon$$

- Two ways of thinking about β_1 :
 - Continuous variables: the *y intercept*
 - Factors: the baseline or *reference* value

WAIT! WHY $N - 1$ COEFFICIENTS? WHAT IS β_1 ?

$$w = \beta_1 + \beta_2 h + \beta_3 e + \beta_4 g_m + \beta_5 d + \beta_6 c_s + \beta_7 c_a + \varepsilon$$

- Two ways of thinking about β_1 :
 - Continuous variables: the *y intercept*
 - Factors: the baseline or *reference* value
- This baseline is the value for the *first levels* of each factor

WAIT! WHY $N - 1$ COEFFICIENTS? WHAT IS β_1 ?

$$w = \beta_1 + \beta_2 h + \beta_3 e + \beta_4 g_m + \beta_5 d + \beta_6 c_s + \beta_7 c_a + \varepsilon$$

- Two ways of thinking about β_1 :
 - Continuous variables: the *y intercept*
 - Factors: the baseline or *reference* value
- This baseline is the value for the *first levels* of each factor
- All response values start at this baseline

WAIT! WHY $N - 1$ COEFFICIENTS? WHAT IS β_1 ?

$$w = \beta_1 + \beta_2 h + \beta_3 e + \beta_4 g_m + \beta_5 d + \beta_6 c_s + \beta_7 c_a + \varepsilon$$

- Two ways of thinking about β_1 :
 - Continuous variables: the *y intercept*
 - Factors: the baseline or *reference* value
- This baseline is the value for the *first levels* of each factor
- All response values start at this baseline
- All the other coefficients measure *differences* from β_1 :

WAIT! WHY $N - 1$ COEFFICIENTS? WHAT IS β_1 ?

$$w = \beta_1 + \beta_2 h + \beta_3 e + \beta_4 g_m + \beta_5 d + \beta_6 c_s + \beta_7 c_a + \varepsilon$$

- Two ways of thinking about β_1 :
 - Continuous variables: the *y intercept*
 - Factors: the baseline or *reference* value
- This baseline is the value for the *first levels* of each factor
- All response values start at this baseline
- All the other coefficients measure *differences* from β_1 :
 - along a continuous slope

WAIT! WHY $N - 1$ COEFFICIENTS? WHAT IS β_1 ?

$$w = \beta_1 + \beta_2 h + \beta_3 e + \beta_4 g_m + \beta_5 d + \beta_6 c_s + \beta_7 c_a + \varepsilon$$

- Two ways of thinking about β_1 :
 - Continuous variables: the *y intercept*
 - Factors: the baseline or *reference* value
- This baseline is the value for the *first levels* of each factor
- All response values start at this baseline
- All the other coefficients measure *differences* from β_1 :
 - along a continuous slope
 - as an offset to a different level

So, TO PUT IT SIMPLY,

Linear models are just a sum of *terms* that are *linear* in the *coefficients*:

$$w = \beta_1 + \beta_2 h + \beta_3 e + \beta_4 g_m + \beta_5 d + \beta_6 c_s + \beta_7 c_a + \varepsilon$$

So, TO PUT IT SIMPLY,

Linear models are just a sum of *terms* that are *linear* in the *coefficients*:

$$w = \beta_1 + \beta_2 h + \beta_3 e + \beta_4 g_m + \beta_5 d + \beta_6 c_s + \beta_7 c_a + \varepsilon$$

What our example linear model means (literally):

- The coefficient β_1 is the baseline value of *weight* for *women* with *no games console*

So, TO PUT IT SIMPLY,

Linear models are just a sum of *terms* that are *linear* in the *coefficients*:

$$w = \beta_1 + \beta_2 h + \beta_3 e + \beta_4 g_m + \beta_5 d + \beta_6 c_s + \beta_7 c_a + \varepsilon$$

What our example linear model means (literally):

- The coefficient β_1 is the baseline value of *weight* for *women* with *no games console*
- The model tells us how much (determined by the other coefficients) to add to this baseline weight ...

So, TO PUT IT SIMPLY,

Linear models are just a sum of *terms* that are *linear* in the *coefficients*:

$$w = \beta_1 + \beta_2 h + \beta_3 e + \beta_4 g_m + \beta_5 d + \beta_6 c_s + \beta_7 c_a + \varepsilon$$

What our example linear model means (literally):

- The coefficient β_1 is the baseline value of *weight* for *women* with *no games console*
- The model tells us how much (determined by the other coefficients) to add to this baseline weight ...
 - for a height of 1.82 metres?

So, TO PUT IT SIMPLY,

Linear models are just a sum of *terms* that are *linear* in the *coefficients*:

$$w = \beta_1 + \beta_2 h + \beta_3 e + \beta_4 g_m + \beta_5 d + \beta_6 c_s + \beta_7 c_a + \varepsilon$$

What our example linear model means (literally):

- The coefficient β_1 is the baseline value of *weight* for *women* with *no games console*
- The model tells us how much (determined by the other coefficients) to add to this baseline weight ...
 - for a height of 1.82 metres?
 - for doing 150 minutes of exercise a week?

So, TO PUT IT SIMPLY,

Linear models are just a sum of *terms* that are *linear* in the *coefficients*:

$$w = \beta_1 + \beta_2 h + \beta_3 e + \beta_4 g_m + \beta_5 d + \beta_6 c_s + \beta_7 c_a + \varepsilon$$

What our example linear model means (literally):

- The coefficient β_1 is the baseline value of *weight* for *women* with *no games console*
- The model tells us how much (determined by the other coefficients) to add to this baseline weight ...
 - for a height of 1.82 metres?
 - for doing 150 minutes of exercise a week?
 - for being male?

SO, TO PUT IT SIMPLY,

Linear models are just a sum of *terms* that are *linear* in the *coefficients*:

$$w = \beta_1 + \beta_2 h + \beta_3 e + \beta_4 g_m + \beta_5 d + \beta_6 c_s + \beta_7 c_a + \varepsilon$$

What our example linear model means (literally):

- The coefficient β_1 is the baseline value of *weight* for *women* with *no games console*
- The model tells us how much (determined by the other coefficients) to add to this baseline weight ...
 - for a height of 1.82 metres?
 - for doing 150 minutes of exercise a week?
 - for being male?
 - for living 2416 metres from a Greggs?

SO, TO PUT IT SIMPLY,




Linear models are just a sum of *terms* that are *linear* in the *coefficients*:

$$w = \beta_1 + \beta_2 h + \beta_3 e + \beta_4 g_m + \beta_5 d + \beta_6 c_s + \beta_7 c_a + \varepsilon$$

What our example linear model means (literally):


- The coefficient β_1 is the baseline value of *weight* for *women* with *no games console*
- The model tells us how much (determined by the other coefficients) to add to this baseline weight ...
 - for a height of 1.82 metres?
 - for doing 150 minutes of exercise a week?
 - for being male?
 - for living 2416 metres from a Greggs?
 - for owning an Xbox?

EXAMPLES OF LINEAR MODELS



- These are *all* linear models (fitted to data)
- Each model a *sum of terms that are linear in coefficients*
- *Linear models can include curved relationships (e.g. polynomials) — this is a common point of confusion!*

LINEAR MODEL WITH ONE CONTINUOUS VARIABLE



$$y = \beta_1 x$$

$$4 = 4 \times 1$$


$$8 = 4 \times 2$$

$$12 = 4 \times 3$$

$$16 = 4 \times 4$$

$$\beta_1 = 4$$

LINEAR MODEL WITH ONE CONTINUOUS VARIABLE



$$y = \beta_1 x$$

$$4 = 4 \times 1$$

$$8 = 4 \times 2$$


$$12 = 4 \times 3$$

$$16 = 4 \times 4$$

$$\beta_1 = 4$$

Regression with *known baseline value (intercept)*

LINEAR MODEL WITH ONE CONTINUOUS VARIABLE



$$y = \beta_1 + \beta_2 x$$

$$9 = 5 + 4 \times 1$$


$$13 = 5 + 4 \times 2$$

$$17 = 5 + 4 \times 3$$

$$21 = 5 + 4 \times 4$$

$$\beta_1 = 5; \beta_2 = 4$$

LINEAR MODEL WITH ONE CONTINUOUS VARIABLE



$$y = \beta_1 + \beta_2 x$$

$$9 = 5 + 4 \times 1$$

$$13 = 5 + 4 \times 2$$


$$17 = 5 + 4 \times 3$$

$$21 = 5 + 4 \times 4$$

$$\beta_1 = 5; \beta_2 = 4$$

Regression with *unknown baseline value (intercept)*

LINEAR MODEL WITH ONE FACTOR (CATEGORICAL VARIABLE)



$$y = \beta_1 + \beta_2 g_m$$

$$2 = 2 + 3 \times 0$$

$$2 = 2 + 3 \times 0$$

$$2 = 2 + 3 \times 0$$


$$5 = 2 + 3 \times 1$$

$$5 = 2 + 3 \times 1$$

$$5 = 2 + 3 \times 1$$

$$\beta_1 = 2; \beta_2 = 3$$

LINEAR MODEL WITH ONE FACTOR (CATEGORICAL VARIABLE)



$$y = \beta_1 + \beta_2 g_m$$

$$2 = 2 + 3 \times 0$$

$$2 = 2 + 3 \times 0$$

$$2 = 2 + 3 \times 0$$

$$5 = 2 + 3 \times 1$$


$$5 = 2 + 3 \times 1$$

$$5 = 2 + 3 \times 1$$

$$\beta_1 = 2; \beta_2 = 3$$

Analysis of Variance (ANOVA)

LINEAR MODEL WITH ONE CONTINUOUS VARIABLE AND ONE FACTOR



$$y = \beta_1 + \beta_2 x + \beta_3 g_m$$

$$3 = 1 + 2 \times 1 + 3 \times 0$$

$$5 = 1 + 2 \times 2 + 3 \times 0$$

$$7 = 1 + 2 \times 3 + 3 \times 0$$


$$6 = 1 + 2 \times 1 + 3 \times 1$$

$$8 = 1 + 2 \times 2 + 3 \times 1$$

$$10 = 1 + 2 \times 3 + 3 \times 1$$

$$\beta_1 = 1; \beta_2 = 2; \beta_3 = 3$$

LINEAR MODEL WITH ONE CONTINUOUS VARIABLE AND ONE FACTOR



$$y = \beta_1 + \beta_2 x + \beta_3 g_m$$

$$3 = 1 + 2 \times 1 + 3 \times 0$$

$$5 = 1 + 2 \times 2 + 3 \times 0$$

$$7 = 1 + 2 \times 3 + 3 \times 0$$

$$6 = 1 + 2 \times 1 + 3 \times 1$$


$$8 = 1 + 2 \times 2 + 3 \times 1$$

$$10 = 1 + 2 \times 3 + 3 \times 1$$

$$\beta_1 = 1; \beta_2 = 2; \beta_3 = 3$$

Multiple Explanatory variables, Analysis of Covariance (ANCOVA)

CLOSER LOOK AT THE ANCOVA EXAMPLE



$$y = \beta_1 + \beta_2 x + \beta_3 g_m$$

$$3 = 1 + 2 \times 1 + 3 \times 0$$

$$5 = 1 + 2 \times 2 + 3 \times 0$$

$$7 = 1 + 2 \times 3 + 3 \times 0$$


$$6 = 1 + 2 \times 1 + 3 \times 1$$

$$8 = 1 + 2 \times 2 + 3 \times 1$$

$$10 = 1 + 2 \times 3 + 3 \times 1$$

$$\beta_1 = 1; \beta_2 = 2; \beta_3 = 3$$


“FITTING” A LINEAR MODEL TO DATA



Rizzuto et al. 2017, Nat Ecol Evol

- Data always shows variation from a perfect model (deviations)
 - Missing variables (age, lab vs. field biology, time of day)


“FITTING” A LINEAR MODEL TO DATA



Rizzuto et al. 2017, Nat Ecol Evol

- Data always shows variation from a perfect model (deviations)
 - Missing variables (age, lab vs. field biology, time of day)
 - Measurement error


“FITTING” A LINEAR MODEL TO DATA



Rizzuto et al. 2017, Nat Ecol Evol


- Data always shows variation from a perfect model (deviations)
 - Missing variables (age, lab vs. field biology, time of day)
 - Measurement error
 - Stochastic variation

FITTING A LINEAR MODEL TO DATA



*What line best passes through
(describes) these data?*


FITTING A LINEAR MODEL TO DATA



*What line best passes through
(describes) these data?*

$$y = \beta_1 + \beta_2 x$$

FITTING A LINEAR MODEL TO DATA



*What line best passes through
(describes) these data?*

$$y = \beta_1 + \beta_2 x$$


$$9.50 = ? + ? \times 1$$

$$11.00 = ? + ? \times 2$$

$$19.58 = ? + ? \times 3$$

$$20.00 = ? + ? \times 4$$

FITTING A LINEAR MODEL TO DATA: GUESS



$$y = \beta_1 + \beta_2 x + \varepsilon$$

$$9.50 = 12.52 + 1 \times 1 - 4.02$$


$$11.00 = 12.52 + 1 \times 2 - 3.52$$

$$19.58 = 12.52 + 1 \times 3 + 4.06$$

$$20.00 = 12.52 + 1 \times 4 + 3.48$$

$$\beta_1 = 12.52; \beta_2 = 1$$

FITTING A LINEAR MODEL TO DATA: GUESS AGAIN!



$$y = \beta_1 + \beta_2 x + \epsilon$$

$$9.50 = -2.48 + 7 \times 1 + 4.98$$


$$11.00 = -2.48 + 7 \times 2 - 0.52$$

$$19.58 = -2.48 + 7 \times 3 + 1.06$$

$$20.00 = -2.48 + 7 \times 4 - 5.52$$

$$\beta_1 = -2.48; \beta_2 = 7$$

FITTING A LINEAR MODEL TO DATA: GUESS AGAIN!



$$y = \beta_1 + \beta_2 x + \epsilon$$

$$9.50 = -2.48 + 7 \times 1 + 4.98$$

$$11.00 = -2.48 + 7 \times 2 - 0.52$$

$$19.58 = -2.48 + 7 \times 3 + 1.06$$

$$20.00 = -2.48 + 7 \times 4 - 5.52$$


$$\beta_1 = -2.48; \beta_2 = 7$$

There must be a better way to do this!

FITTING A LINEAR MODEL: LEAST SQUARES SOLUTION

Minimize the *sum* of the *squared residuals*:

THE (ORDINARY) LEAST SQUARES FITTING SOLUTION



$$y = \beta_1 + \beta_2 x + \epsilon$$

$$9.50 = 5 + 4 \times 1 + 0.50$$

$$11.00 = 5 + 4 \times 2 - 2.00$$

$$19.58 = 5 + 4 \times 3 + 2.58$$

$$20.00 = 5 + 4 \times 4 - 1.00$$

$$\beta_1 = 5; \beta_2 = 4$$

THE MATHS MAGIC UNDER THE HOOD

$$\mathbf{Y} = \mathbf{X}\beta + \varepsilon$$

Observed values

$$\begin{bmatrix} y_1 \\ y_2 \\ y_3 \\ y_4 \end{bmatrix}$$

Coefficients

$$\begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix}$$

Model matrix

$$\begin{bmatrix} 1 & x_1 \\ 1 & x_2 \\ 1 & x_3 \\ 1 & x_4 \end{bmatrix}$$

Residuals

$$\begin{bmatrix} \varepsilon_1 \\ \varepsilon_2 \\ \varepsilon_3 \\ \varepsilon_4 \end{bmatrix}$$

THE MATHS MAGIC UNDER THE HOOD

$$\mathbf{Y} = \mathbf{X}\beta + \varepsilon$$

Observed values

$$\begin{bmatrix} 9.50 \\ 11.00 \\ 19.58 \\ 20.00 \end{bmatrix}$$

Coefficients

$$\begin{bmatrix} 5 \\ 4 \end{bmatrix}$$

Model matrix

$$\begin{bmatrix} 1 & 1 \\ 1 & 2 \\ 1 & 3 \\ 1 & 4 \end{bmatrix}$$

Residuals

$$\begin{bmatrix} 0.50 \\ -2.00 \\ 2.58 \\ -1.00 \end{bmatrix}$$

=

+

THE MATHS MAGIC UNDER THE HOOD

$$\mathbf{Y} = \mathbf{X}\beta + \varepsilon$$

Observed values

$$\begin{bmatrix} 9.50 \\ 11.00 \\ 19.58 \\ 20.00 \end{bmatrix}$$

Coefficients

$$\begin{bmatrix} 5 \\ 4 \end{bmatrix}$$

Model matrix

$$\begin{bmatrix} 1 & 1 \\ 1 & 2 \\ 1 & 3 \\ 1 & 4 \end{bmatrix}$$

Residuals

$$\begin{bmatrix} 0.50 \\ -2.00 \\ 2.58 \\ -1.00 \end{bmatrix}$$

THE MATHS MAGIC UNDER THE HOOD

$$\mathbf{Y} = \mathbf{X}\beta + \varepsilon$$

Observed values

$$\begin{bmatrix} 9.50 \\ 11.00 \\ 19.58 \\ 20.00 \end{bmatrix}$$

Coefficients

$$\begin{bmatrix} 5 \\ 4 \end{bmatrix}$$

Model matrix

$$\begin{bmatrix} 1 & 1 \\ 1 & 2 \\ 1 & 3 \\ 1 & 4 \end{bmatrix}$$

Residuals

$$\begin{bmatrix} 0.50 \\ -2.00 \\ 2.58 \\ -1.00 \end{bmatrix}$$

=

+

THE MATHS MAGIC UNDER THE HOOD

$$\mathbf{Y} = \mathbf{X}\beta + \varepsilon$$

Observed values

$$\begin{bmatrix} 9.50 \\ 11.00 \\ 19.58 \\ 20.00 \end{bmatrix}$$

Coefficients

$$\begin{bmatrix} 5 \\ 4 \end{bmatrix}$$

Model matrix

$$\begin{bmatrix} 1 & 1 \\ 1 & 2 \\ 1 & 3 \\ 1 & 4 \end{bmatrix}$$

Residuals

$$\begin{bmatrix} 0.50 \\ -2.00 \\ 2.58 \\ -1.00 \end{bmatrix}$$

=

+

THE MATHS MAGIC UNDER THE HOOD

$$\mathbf{Y} = \mathbf{X}\beta + \varepsilon$$

Observed values

$$\begin{bmatrix} 9.50 \\ 11.00 \\ 19.58 \\ 20.00 \end{bmatrix}$$

Coefficients

$$\begin{bmatrix} 5 \\ 4 \end{bmatrix}$$

Model matrix

$$\begin{bmatrix} 1 & 1 \\ 1 & 2 \\ 1 & 3 \\ 1 & 4 \end{bmatrix}$$

Residuals

$$\begin{bmatrix} 0.50 \\ -2.00 \\ 2.58 \\ -1.00 \end{bmatrix}$$

THE MATHS MAGIC UNDER THE HOOD

$$\mathbf{Y} = \mathbf{X}\beta + \varepsilon$$

Given these ...

$$\begin{bmatrix} 9.50 \\ 11.00 \\ 19.58 \\ 20.00 \end{bmatrix}$$

=

$$\begin{bmatrix} 1 & 1 \\ 1 & 2 \\ 1 & 3 \\ 1 & 4 \end{bmatrix}$$

... find the set of these...

$$\begin{bmatrix} 5 \\ 4 \end{bmatrix}$$

+

$$\begin{bmatrix} 0.50 \\ -2.00 \\ 2.58 \\ -1.00 \end{bmatrix}$$

... that minimize the sum of the squares of these.

THE MATHS MAGIC UNDER THE HOOD

$$\hat{\mathbf{Y}} = \mathbf{X}\beta$$

Predicted or fitted values


Coefficients

The diagram illustrates the matrix equation $\hat{\mathbf{Y}} = \mathbf{X}\beta$. On the left, a red box labeled "Predicted or fitted values" has a red arrow pointing down to the vector $\begin{bmatrix} 9 \\ 13 \\ 17 \\ 21 \end{bmatrix}$. In the middle is an equals sign. To the right of the equals sign is the matrix $\begin{bmatrix} 1 & 1 \\ 1 & 2 \\ 1 & 3 \\ 1 & 4 \end{bmatrix}$, which has a red box labeled "Model matrix" below it with a red arrow pointing up to it. To the right of the matrix is a vector $\begin{bmatrix} 5 \\ 4 \end{bmatrix}$, which has a red box labeled "Coefficients" above it with a red arrow pointing down to it.

$$\begin{bmatrix} 9 \\ 13 \\ 17 \\ 21 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & 2 \\ 1 & 3 \\ 1 & 4 \end{bmatrix} \begin{bmatrix} 5 \\ 4 \end{bmatrix}$$

Model matrix

PREDICTED VALUES AND RESIDUALS



$$\hat{y} = \beta_1 + \beta_2 x$$

$$9 = 5 + 4 \times 1$$

$$13 = 5 + 4 \times 2$$

$$17 = 5 + 4 \times 3$$

$$21 = 5 + 4 \times 4$$



FITTING A LINEAR MODEL: ASSUMPTIONS

- Linear models are fitted with the following assumptions:
 - No measurement error in explanatory variables
 - The explanatory variables are not very highly (inter-) correlated
 - The model has constant normal variance
- **If these assumptions are not met, the model can be very wrong**
- The first two you will should consider *before* even fitting a linear model

FITTING A LINEAR MODEL: ASSUMPTIONS




- Linear models are fitted with the following assumptions:
 - No measurement error in explanatory variables
 - The explanatory variables are not very highly (inter-) correlated
 - **The model has constant normal variance**
- **If these assumptions are not met, the model can be very wrong**
- The first two you will should consider *before* even fitting a linear model
- The last one needs can be tested *after* fitting a linear model

'THE MODEL HAS CONSTANT NORMAL VARIANCE'






- The data have a similar spread around any predicted point in the model
- Overall, the residuals are *normally distributed*: mostly small but a few larger values
- Points *should* be spaced so as to best capture the normal (gaussian) curve

CHECKING IF THE LINEAR MODEL IS APPROPRIATE






- All these three linear model fits appropriate for the data? Are assumptions of the linear model fit satisfied?

CHECKING IF THE LINEAR MODEL IS APPROPRIATE






- All these three linear model fits appropriate for the data? Are assumptions of the linear model fit satisfied?

CHECKING IF THE LINEAR MODEL IS APPROPRIATE



- All these three linear model fits appropriate for the data? Are assumptions of the linear model fit satisfied?
 - The spread of the real data around the fitted line (fitted values) is about the same across the x-axis – good

CHECKING IF THE LINEAR MODEL IS APPROPRIATE



- All these three linear model fits appropriate for the data? Are assumptions of the linear model fit satisfied?
 - The spread of the real data around the fitted line (fitted values) is about the same across the x-axis – good
 - But are the residuals normally distributed?

DIAGNOSTICS FOR A FITTED LINEAR MODEL




- *The spread of the real data around the fitted line (fitted values) is about the same across the x-axis*

DIAGNOSTICS FOR A FITTED LINEAR MODEL

- *The spread of the real data around the fitted line (fitted values) is about the same across the x-axis*


DIAGNOSTICS FOR A FITTED LINEAR MODEL

- *The spread of the real data around the fitted line (fitted values) is about the same across the x-axis*



DIAGNOSTICS FOR A FITTED LINEAR MODEL


- *The spread of the real data around the fitted line (fitted values) is about the same across the x-axis*



- That is, the residuals have about the same spread irrespective of the fitted values

DIAGNOSTICS FOR A FITTED LINEAR MODEL


- *The spread of the real data around the fitted line (fitted values) is about the same across the x-axis*



- That is, the residuals have about the same spread irrespective of the fitted values
- The three numbered points in each plot are the three most 'badly behaved' data points.

DIAGNOSTICS FOR A FITTED LINEAR MODEL

- *The spread of the real data around the fitted line (fitted values) is about the same across the x-axis*



- That is, the residuals have about the same spread irrespective of the fitted values
- The three numbered points in each plot are the three most 'badly behaved' data points.
 - Each number is the datum's row number in the R data frame

DIAGNOSTICS FOR A FITTED LINEAR MODEL




- *Are the residuals normally distributed?*

DIAGNOSTICS FOR A FITTED LINEAR MODEL

- *Are the residuals normally distributed?*




DIAGNOSTICS FOR A FITTED LINEAR MODEL

- *Are the residuals normally distributed?*






DIAGNOSTICS FOR A FITTED LINEAR MODEL

- Are the residuals normally distributed?






- Residuals from the first (simple regression) and third (polynomial) model's fits show some deviations from normality at the ends (high and low ends of their distributions), but it's acceptable

THREE BAD LINEAR MODEL FITS






- These are three bad linear model fits

THREE BAD LINEAR MODEL FITS






- These are three bad linear model fits
 - The data spread is not the same for all fitted values

THREE BAD LINEAR MODEL FITS






- These are three bad linear model fits
 - The data spread is not the same for all fitted values
 - The first model clearly spread is not the same for all fitted values

THREE BAD LINEAR MODEL FITS









- These are three bad linear model fits
 - The data spread is not the same for all fitted values
 - The first model clearly spread is not the same for all fitted values
 - Are the residuals normally distributed?

DIAGNOSTICS FOR A (BADLY) FITTED LINEAR MODEL



DIAGNOSTICS FOR A (BADLY) FITTED LINEAR MODEL



IS A LINEAR MODEL APPROPRIATE?

Plot the data!
Plot the residuals!

HOW EXPLANATORY IS THE FITTED LINEAR MODEL?

- The role of F and t tests in Linear Model fitting

HOW EXPLANATORY IS THE FITTED LINEAR MODEL?

- The role of F and t tests in Linear Model fitting
- Significance of *Terms*: F test

HOW EXPLANATORY IS THE FITTED LINEAR MODEL?

- The role of F and t tests in Linear Model fitting
- Significance of *Terms*: F test
 - Does the model explain enough variation?

HOW EXPLANATORY IS THE FITTED LINEAR MODEL?

- The role of F and t tests in Linear Model fitting
- Significance of *Terms*: F test
 - Does the model explain enough variation?
 - Does each term explain enough variation?

HOW EXPLANATORY IS THE FITTED LINEAR MODEL?

- The role of F and t tests in Linear Model fitting
- Significance of *Terms*: F test
 - Does the model explain enough variation?
 - Does each term explain enough variation?
- Significance of *Coefficients*: t tests

HOW EXPLANATORY IS THE FITTED LINEAR MODEL?

- The role of F and t tests in Linear Model fitting
- Significance of *Terms*: F test
 - Does the model explain enough variation?
 - Does each term explain enough variation?
- Significance of *Coefficients*: t tests
 - Are the coefficients different from zero?

IS THE FITTED LINEAR MODEL SIGNIFICANT?: F TEST

- **Total sum of squares (TSS):** Sum of the squared difference between the observed dependent variable (y) and the mean of y (\bar{y}), or, $TSS = \sum_{i=1}^n (y_i - \bar{y})^2$
TSS tells us how much variation there is in the dependent variable

IS THE FITTED LINEAR MODEL SIGNIFICANT?: F TEST

- **Total sum of squares (TSS):** Sum of the squared difference between the observed dependent variable (y) and the mean of y (\bar{y}), or, $TSS = \sum_{i=1}^n (y_i - \bar{y})^2$
TSS tells us how much variation there is in the dependent variable
- **Explained sum of squares (ESS):** Sum of the squared differences between the predicted y (\hat{y}) and \bar{y} , or, $ESS = \sum_{i=1}^n (\hat{y}_i - \bar{y})^2$
ESS tells us how much of the variation in the dependent variable our model was able to explain


IS THE FITTED LINEAR MODEL SIGNIFICANT?: F TEST

- **Total sum of squares (TSS):** Sum of the squared difference between the observed dependent variable (y) and the mean of y (\bar{y}), or, $TSS = \sum_{i=1}^n (y_i - \bar{y})^2$
TSS tells us how much variation there is in the dependent variable
- **Explained sum of squares (ESS):** Sum of the squared differences between the predicted y (\hat{y}) and \bar{y} , or, $ESS = \sum_{i=1}^n (\hat{y}_i - \bar{y})^2$
ESS tells us how much of the variation in the dependent variable our model was able to explain
- **Residual sum of squares (RSS):** Sum of the squared differences between the observed y and the predicted \hat{y} (residuals), or, $RSS = \sum_{i=1}^n (\hat{y}_i - y_i)^2$
RSS tells us how much of the variation in the dependent variable our model could not explain


IS THE FITTED LINEAR MODEL SIGNIFICANT?: F TEST

- **Total sum of squares (TSS):** Sum of the squared difference between the observed dependent variable (y) and the mean of y (\bar{y}), or, $TSS = \sum_{i=1}^n (y_i - \bar{y})^2$
TSS tells us how much variation there is in the dependent variable
- **Explained sum of squares (ESS):** Sum of the squared differences between the predicted y (\hat{y}) and \bar{y} , or, $ESS = \sum_{i=1}^n (\hat{y}_i - \bar{y})^2$
ESS tells us how much of the variation in the dependent variable our model was able to explain
- **Residual sum of squares (RSS):** Sum of the squared differences between the observed y and the predicted \hat{y} (residuals), or, $RSS = \sum_{i=1}^n (\hat{y}_i - y_i)^2$
RSS tells us how much of the variation in the dependent variable our model could not explain
- Of course, $TSS = ESS + RSS$

NULL VS. OVER-SPECIFIED MODELS: TWO ENDPOINTS




- The null model (H_0)
- Nothing is going on
- Biggest possible residuals
- Residual sum of squares (RSS) is as big as it can be



- The saturated model
- One coefficient per data point
- RSS is zero - all the sums of squares are now explained (ESS)

THE 'RIGHT' (INTERESTING) MODEL



- Added a term with three levels
- Some but not all of the residual sums of squares are explained
- Is this enough to be interesting?

F STATISTIC OF THE FITTED LINEAR MODEL

$$F = \frac{\text{ESS} / N_c}{\text{RSS} / N_r} = \frac{9.92 / 2}{6.59 / 6} = 4.52$$

F STATISTIC OF THE FITTED LINEAR MODEL

Large ESS
is good

$$F = \frac{\text{ESS} / N_c}{\text{RSS} / N_r} = \frac{9.92 / 2}{6.59 / 6} = 4.52$$

F STATISTIC OF THE FITTED LINEAR MODEL

Large ESS
is good



$$F = \frac{\text{ESS} / N_c}{\text{RSS} / N_r} = \frac{9.92 / 2}{6.59 / 6} = 4.52$$

Small RSS
is good



F STATISTIC OF THE FITTED LINEAR MODEL

Large ESS
is good

Fewer coefficients
is better

$$F = \frac{\text{ESS} / N_c}{\text{RSS} / N_r} = \frac{9.92 / 2}{6.59 / 6} = 4.52$$

Small RSS
is good

F STATISTIC OF THE FITTED LINEAR MODEL

The diagram illustrates the components of the F-statistic formula. Four blue-bordered boxes with arrows point to the terms in the formula:

- Large ESS is good** points to ESS.
- Fewer coefficients is better** points to N_c .
- Small RSS is good** points to RSS.
- Residual degrees of freedom: larger sample size is better** points to N_r .

$$F = \frac{\text{ESS} / N_c}{\text{RSS} / N_r} = \frac{9.92 / 2}{6.59 / 6} = 4.52$$


WHAT IT REALLY MEANS: F VALUE BY CHANCE?

What would be the distribution of F if nothing is going on?


WHAT IT REALLY MEANS: F VALUE BY CHANCE?

What would be the distribution of F if nothing is going on?


$$y = \beta_1 + \beta_2 f_b + \beta_3 f_c$$



$$y = \beta_1 + \beta_2 f_b + \beta_3 f_c$$






$$y = \beta_1 + \beta_2 f_b + \beta_3 f_c$$



WHAT IT REALLY MEANS: F VALUE BY CHANCE?


What would be the distribution of F if nothing is going on?



- Simulate 10,000 datasets where nothing is going on (H_0 is true)

WHAT IT REALLY MEANS: F VALUE BY CHANCE?


What would be the distribution of F if nothing is going on?



- Simulate 10,000 datasets where nothing is going on (H_0 is true)
- Calculate F for each random dataset under H_1

WHAT IT REALLY MEANS: F VALUE BY CHANCE?

What would be the distribution of F if nothing is going on?



- Simulate 10,000 datasets where nothing is going on (H_0 is true)
- Calculate F for each random dataset under H_1
- H_1 typically has a low F – but sometimes it is high *by chance*

WHAT IT REALLY MEANS: F VALUE BY CHANCE?

- In our possibly interesting model, $F = 4.52$

WHAT IT REALLY MEANS: F VALUE BY CHANCE?

- In our possibly interesting model, $F = 4.52$
- 95% of the random data sets have $F \leq 5.5$

WHAT IT REALLY MEANS: F VALUE BY CHANCE?

- In our possibly interesting model, $F = 4.52$
- 95% of the random data sets have $F \leq 5.5$
- A model this good would be found by chance 1 in 16 times ($p = 0.063$)

WHAT IT REALLY MEANS: F VALUE BY CHANCE?

- In our possibly interesting model, $F = 4.52$
- 95% of the random data sets have $F \leq 5.5$
- A model this good would be found by chance 1 in 16 times ($p = 0.063$)
- Close, but not quite interesting (significant) enough!

ARE THE COEFFICIENTS DIFFERENT FROM ZERO?

$$t = \frac{\text{Effect size}}{\text{Precision}} = \frac{\text{Coefficient value}}{\text{Standard error}}$$

ARE THE COEFFICIENTS DIFFERENT FROM ZERO?

Large is good: bigger changes

$$t = \frac{\text{Effect size}}{\text{Precision}} = \frac{\text{Coefficient value}}{\text{Standard error}}$$

ARE THE COEFFICIENTS DIFFERENT FROM ZERO?

Large is good: bigger changes

↓ ↓

$$t = \frac{\text{Effect size}}{\text{Precision}} = \frac{\text{Coefficient value}}{\text{Standard error}}$$

↑ ↑


Small is good: known more precisely

ARE THE COEFFICIENTS DIFFERENT FROM ZERO?

Large is good: bigger changes

$$t = \frac{\text{Effect size}}{\text{Precision}} = \frac{\text{Coefficient value}}{\text{Standard error}}$$

Small is good: known more precisely



- The value of a coefficient in a model is an *effect size*

ARE THE COEFFICIENTS DIFFERENT FROM ZERO?

Large is good: bigger changes

↓ ↓

$$t = \frac{\text{Effect size}}{\text{Precision}} = \frac{\text{Coefficient value}}{\text{Standard error}}$$

↑ ↑

Small is good: known more precisely


- The value of a coefficient in a model is an *effect size*
- How much does changing that predictor variable change the response variable?

ARE THE COEFFICIENTS DIFFERENT FROM ZERO?

Large is good: bigger changes


$$t = \frac{\text{Effect size}}{\text{Precision}} = \frac{\text{Coefficient value}}{\text{Standard error}}$$

Small is good: known more precisely



- The value of a coefficient in a model is an *effect size*
- How much does changing that predictor variable change the response variable?
- The *standard error* estimates how precisely we know the value

VARIATION IN EFFECT SIZE AND PRECISION






WHAT IT REALLY MEANS: t VALUES BY CHANCE

What is the distribution of t if nothing is going on?


WHAT IT REALLY MEANS: t VALUES BY CHANCE

What is the distribution of t if nothing is going on?



WHAT IT REALLY MEANS: t VALUES BY CHANCE


What is the distribution of t if nothing is going on?



- Simulate 10,000 datasets where nothing is going on (H_0 is true)

WHAT IT REALLY MEANS: t VALUES BY CHANCE


What is the distribution of t if nothing is going on?



- Simulate 10,000 datasets where nothing is going on (H_0 is true)
- Calculate t for each random dataset under H_1

WHAT IT REALLY MEANS: t VALUES BY CHANCE

What is the distribution of t if nothing is going on?



- Simulate 10,000 datasets where nothing is going on (H_0 is true)
- Calculate t for each random dataset under H_1
- H_1 typically has a t near zero but can be strongly positive or negative *by chance*

DISTRIBUTION OF t




DISTRIBUTION OF t

- 95% of the random data sets have $t \leq \pm 2.09$

DISTRIBUTION OF t




- 95% of the random data sets have $t \leq \pm 2.09$
- Only the two higher precision models are expected to occur less than 1 time in 20 by chance.

SOME MORE EXAMPLES OF LINEAR MODEL FITTING






- The null hypothesis (H_0): Nothing is going on (model is just β_1 !)

SOME MORE EXAMPLES OF LINEAR MODEL FITTING



- The null hypothesis (H_0): Nothing is going on (model is just β_1 !)
- The residuals (and therefore, RSS) will get *smaller* as we include more terms to the model




SOME MORE EXAMPLES OF LINEAR MODEL FITTING



- The null hypothesis (H_0): Nothing is going on (model is just β_1 !)
- The residuals (and therefore, RSS) will get *smaller* as we include more terms to the model
- *How much smaller is enough?*

SOME MORE EXAMPLES OF LINEAR MODEL FITTING




First try: Add one continuous term



- Fitted an *alternative* model (H_1) using a predictor variable x

SOME MORE EXAMPLES OF LINEAR MODEL FITTING




First try: Add one continuous term



- Fitted an *alternative* model (H_1) using a predictor variable x
- i.e., Added one term (x) to the model to give (H_1)

SOME MORE EXAMPLES OF LINEAR MODEL FITTING




First try: Add one continuous term



- Fitted an *alternative* model (H_1) using a predictor variable x
- i.e., Added one term (x) to the model to give (H_1)
- Do we reject H_0 and accept this new model?

SOME MORE EXAMPLES OF LINEAR MODEL FITTING




Second try: Add one continuous term



- Fitted another model (H_2) with continuous predictor x and factor f

SOME MORE EXAMPLES OF LINEAR MODEL FITTING




Second try: Add one continuous term



- Fitted another model (H_2) with continuous predictor x and factor f
- The RSS gets still smaller

SOME MORE EXAMPLES OF LINEAR MODEL FITTING

Second try: Add one continuous term



- Fitted another model (H_2) with continuous predictor x and factor f
- The RSS gets still smaller
- Is this *even* better than H_1 ?

COMPARE THE THREE MODELS

		Model A	Model B	Model C
H_0	Unexplained SS	241.97	185.02	259.80
	Explained SS	0	0	0
H_1	Unexplained SS	241.97	173.21	62.95
	Explained SS	0.00	11.81	196.85
H_2	Unexplained SS	238.07	123.75	25.05
	Explained SS	3.9	61.27	234.75

- Which model would you choose between H_1 and H_2 ?

COMPARE THE THREE MODELS

		Model A	Model B	Model C
H_0	Unexplained SS	241.97	185.02	259.80
	Explained SS	0	0	0
H_1	Unexplained SS	241.97	173.21	62.95
	Explained SS	0.00	11.81	196.85
H_2	Unexplained SS	238.07	123.75	25.05
	Explained SS	3.9	61.27	234.75

- Which model would you choose between H_1 and H_2 ?
- Every alternative model is an *alternative hypothesis*

LINEAR MODELS: SUMMARY

- Linear models predict a continuous response variable
- A LM is a sum of terms that are linear in the coefficients capturing the effect sizes of explanatory variables
- LMs are fitted using (ordinary) least squares — minimizes sum of squared residuals
- Need to check if the fitted LM is appropriate
- Then check if the LM is explanatory
- Fitting alternative LMs = Testing alternative hypotheses