# Real Analysis Theorems

Zhiwei Zhang

April 13, 2019

### 1 Limsup and Liminf

Corollary 1.0.1. If  $\lim_{n \to \infty} \left| \frac{s_{n+1}}{s_n} \right|$  exists [ and equals L], then  $\lim_{n \to \infty} \left| s_n \right|^{1/n}$  exists [ and equals L].

## 2 Uniform Continuity

Theorem 2.1. Pass. ...

### 3 Power Series

Given power series  $\sum_{n=0}^{\infty} a_n x^n$ 

**Theorem 3.1.** Given any  $(a_n)$ , one of the following holds true:

- 1. The power series converges for all  $x \in \mathbb{R}$
- 2. The power series converges only for x = 0
- 3. The power series converges for all x in some bounded interval centered at 0; the intervalmay be open, half-open or closed.

#### Theorem 3.2. Let

$$\beta = \limsup |a_n|^{1/n} \quad and \quad R = \frac{1}{\beta}$$

Then

- 1. The power series converges for |x| < R
- 2. The power series diverges for |x| > R

Also notice that  $\lim_{n \to \infty} \left| \frac{a_{n+1}}{a_n} \right| = \beta$ , therefore most of the time we will use  $\lim_{n \to \infty} \left| \frac{a_{n+1}}{a_n} \right|$  as it's easier to compute that  $\beta$ .

### 4 More on Uniform Convergence

**Theorem 4.1.** Let  $(f_n)$  be a sequence of continuous functions on [a,b], and suppose  $f_n \to f$  uniformly on [a,b]. Then

$$\lim_{n \to \infty} \int_a^b f_n(x) dx = \int_a^b f(x) dx$$

**Definition 4.2.** A sequence  $(f_n)$  of functions defined on a set  $S \subseteq \mathbb{R}$  is uniformly Cauchy on S if

for each 
$$\epsilon > 0$$
 there exists a number N such that  $|f_n(x) - f_m(x)| < \epsilon$  for all  $x \in S$  and all  $m, n > N$ 

**Theorem 4.3.** Let  $(f_n)$  be a sequence of functions defined and uniformly Cauchy on a set  $S \subseteq \mathbb{R}$ . Then there exists a function f on S such that  $f_n \to f$  uniformly on S.

**Theorem 4.4.** Consider a series  $\sum_{k=0}^{\infty} g_k$  of functions on a set  $S \subseteq \mathbb{R}$ . Suppose each  $g_k$  is continuous on S and the series converges uniformly on S. Then the series  $\sum_{k=0}^{\infty} g_k$  represents a continuous function on S.

**Theorem 4.5.** If a series  $\sum_{k=0}^{\infty} g_k$  of functions satisfies the Cauchy criterion uniformly on a set S, then the series converges uniformly on S.

**Theorem 4.6.** Let  $(M_k)$  be a sequence of nonnegative real numbers where  $\sum M_k < \infty$ . If  $|g_k(x)| \leq M_k$  for all x in a set S, then  $\sum g_k$  converges uniformly on S.

**Theorem 4.7.** Show that if the series  $\sum g_n$  converges uniformly on a set S, then  $\lim_{n\to\infty} \sup \{|g_n(x)| : x \in S\}$ 

## 5 Differentiation and Integration of Power Series

**Theorem 5.1.** Let  $\sum_{n=0}^{\infty} a_n x^n$  be a power series with radius of convergence R > 0 [possibly  $R = +\infty$ ]. If  $0 < R_1 < R$ , then the power series converges uniformly on  $[-R_1, R_1]$  to a continuous function.

**Lemma 5.2.** If the power series  $\sum_{n=0}^{\infty} a_n x^n$  has radius of convergence R, then the power series

$$\sum_{n=1}^{\infty} n a_n x^{n-1}$$

and

$$\sum_{n=0}^{\infty} \frac{a_n}{n+1} x^{n+1}$$

also have radius of convergence R.

**Theorem 5.3** (Abel's Theorem). Let  $f(x) = \sum_{n=0}^{\infty} a_n x^n$  be a power series with finite positive radius of convergence R. If the series converges at x = R, then f is continuous at x = R. If the series converges at x = -R, then f is continuous at x = -R.

### 6 Basic Properties of the Derivative

**Theorem 6.1.** Differentiability implies continuity.

### 7 Mean Value Theorem

**Theorem 7.1.** Let f be a continuous function on [a,b] that is differentiable on (a,b). Then there exists  $[at least one] \ x \ in (a,b) \ such that$ 

$$f'(x) = \frac{f(b) - f(a)}{b - a}$$

**Theorem 7.2** (Intermediate Value Theorem for Derivatives). Let f be a differentiable function on (a,b). If  $a < x_1 < x_2 < b$ , and if c lies between  $f'(x_1)$  and  $f'(x_2)$ , there exists [at least one] x in  $(x_1, x_2)$  such that f'(x) = c

**Theorem 7.3.** Let f be a one-to-one continuous function on an open intervalI, and let J = f(I). If f is differentiable at  $x_0 \in I$  and if  $f'(x_0) \neq 0$ , then  $f^{-1}$  is differentiable at  $y_0 = f(x_0)$  and

$$(f^{-1})'(y_0) = \frac{1}{f'(x_0)}$$