

PH 712 Probability and Statistical Inference

Part VI: Evaluating Estimators I

Zhiyang Zhou (zhou67@uwm.edu, zhiyanggeezhou.github.io)

2024/Oct/08 17:52:39

Bias

- Bias of $\hat{\theta}$: $\text{Bias}(\hat{\theta}) = E(\hat{\theta}) - \theta$
- Unbiased: if $E(\hat{\theta}) = \theta$

Mean squared error (MSE)

- $\text{MSE}(\hat{\theta}) = E(\hat{\theta} - \theta)^2 = \text{Bias}^2(\hat{\theta}) + \text{var}(\hat{\theta})$
 - The lower the better
- For unbiased estimators, minimizing the MSE \Leftrightarrow minimizing the variance

Cramér-Rao lower bound (CRLB, CB Thm 7.3.9 & Lemma 7.3.11)

- Recall the score $S(\theta) = \ell'(\theta)$
- $\text{CRLB} = I^{-1}(\theta) \left\{ \frac{d}{d\theta} E(\hat{\theta}) \right\}^2$
 - Reducing to $I^{-1}(\theta)$ if $E(\hat{\theta}) = \theta$ (i.e., unbiased $\hat{\theta}$)
 - Where $I(\theta) = \text{var}\{S(\theta)\} = E[\{S(\theta)\}^2] = -E\{H(\theta)\}$ is called the Fisher information
 - * Where $H(\theta) = S'(\theta) = \ell''(\theta)$ is called the Hessian
 - * The most convenient way to calculate $I(\theta)$: $I(\theta) = -E\{H(\theta)\}$
- Under regularity conditions, $\text{var}(\hat{\theta}) \geq \text{CRLB}$.

Example Lec6.1

- Find the CRLB for all the UNBIASED estimators in the following cases.
 - a. $X_1, \dots, X_n \stackrel{\text{iid}}{\sim} \mathcal{N}(\mu, \sigma^2)$ with UNKNOWN μ and GIVEN σ^2 .
 - b. $X_1, \dots, X_n \stackrel{\text{iid}}{\sim} \mathcal{N}(\mu, \sigma^2)$ with GIVEN μ and UNKNOWN σ^2 .