

# STAT 3690 Lecture 29

zhiyanggeezhou.github.io

Zhiyang Zhou (zhiyang.zhou@umanitoba.ca)

Apr 08, 2022

## CCA (Con'd)

- Sample version
  - $(\mathbf{Y}_1, \mathbf{X}_1), \dots, (\mathbf{Y}_n, \mathbf{X}_n) \stackrel{\text{iid}}{\sim} (\mathbf{Y}, \mathbf{X})$ 
    - \*  $\mathbf{Y}_i$  and  $\mathbf{X}_i$  jointly sampled
    - \*  $p \leq q < n$
  - $n \times p$  matrix  $\mathbb{Y} = [\mathbf{Y}_1, \dots, \mathbf{Y}_n]^\top$  and  $n \times q$  matrix  $\mathbb{X} = [\mathbf{X}_1, \dots, \mathbf{X}_n]^\top$
  - Sample covariance matrices
    - \*  $\mathbf{S}_\mathbf{Y} = (n-1)^{-1} \sum_i (\mathbf{Y}_i - \bar{\mathbf{Y}})(\mathbf{Y}_i - \bar{\mathbf{Y}})^\top$
    - \*  $\mathbf{S}_\mathbf{X} = (n-1)^{-1} \sum_i (\mathbf{X}_i - \bar{\mathbf{X}})(\mathbf{X}_i - \bar{\mathbf{X}})^\top$
    - \*  $\mathbf{S}_{\mathbf{YX}} = \mathbf{S}_{\mathbf{XY}}^\top = (n-1)^{-1} \sum_i (\mathbf{Y}_i - \bar{\mathbf{Y}})(\mathbf{X}_i - \bar{\mathbf{X}})^\top$
  - Vocabulary
    - \* (The  $k$ th pair of) sample canonical directions:  $(\hat{\mathbf{a}}_k \in \mathbb{R}^p, \hat{\mathbf{b}}_k \in \mathbb{R}^q)$
    - \* (The  $k$ th pair of) sample canonical variates:  $(\mathbb{Y}_C \hat{\mathbf{a}}_k, \mathbb{X}_C \hat{\mathbf{b}}_k)$
    - \* (The  $k$ th) canonical correlation:  $\hat{\rho}_k$
  - Goal: find  $\hat{\mathbf{a}}_k$  and  $\hat{\mathbf{b}}_k$ ,  $k = 1, \dots, r \leq p$ , to maximize

$$\hat{\rho}_k = \frac{\hat{\mathbf{a}}_k^\top \mathbf{S}_{\mathbf{YX}} \hat{\mathbf{b}}_k}{\sqrt{\hat{\mathbf{a}}_k^\top \mathbf{S}_\mathbf{Y} \hat{\mathbf{a}}_k} \sqrt{\hat{\mathbf{b}}_k^\top \mathbf{S}_\mathbf{X} \hat{\mathbf{b}}_k}}$$

subject to

- \*  $\hat{\mathbf{a}}_k^\top \mathbf{S}_\mathbf{Y} \hat{\mathbf{a}}_k = 1$
- \*  $\hat{\mathbf{b}}_k^\top \mathbf{S}_\mathbf{X} \hat{\mathbf{b}}_k = 1$
- \*  $\hat{\mathbf{a}}_k^\top \mathbf{S}_\mathbf{Y} \hat{\mathbf{a}}_\ell = 0, \ell = 1, \dots, k-1$
- \*  $\hat{\mathbf{a}}_k^\top \mathbf{S}_{\mathbf{YX}} \hat{\mathbf{b}}_\ell = 0, \ell = 1, \dots, k-1$
- \*  $\hat{\mathbf{b}}_k^\top \mathbf{S}_\mathbf{X} \hat{\mathbf{b}}_\ell = 0, \ell = 1, \dots, k-1$
- \*  $\hat{\mathbf{b}}_k^\top \mathbf{S}_{\mathbf{XY}} \hat{\mathbf{a}}_\ell = 0, \ell = 1, \dots, k-1$
- Solution: Let  $\widehat{\mathbf{M}} = \mathbf{S}_\mathbf{Y}^{-1/2} \mathbf{S}_{\mathbf{YX}} \mathbf{S}_\mathbf{X}^{-1/2}$ 
  - \*  $\hat{\rho}_k = \sqrt{\hat{\lambda}_k}$  is the  $k$ th largest singular value of  $\widehat{\mathbf{M}}$ 
    - $\hat{\lambda}_k$ : the  $k$ th largest eigenvalue of  $\widehat{\mathbf{M}} \widehat{\mathbf{M}}^\top$  (or  $\widehat{\mathbf{M}}^\top \widehat{\mathbf{M}}$ )
  - \*  $\hat{\mathbf{a}}_k = \mathbf{S}_\mathbf{Y}^{-1/2} \hat{\mathbf{e}}_k$ 
    - $\hat{\mathbf{e}}_k$ : the left-singular vector corresponding to the  $k$ th largest singular value of  $\widehat{\mathbf{M}}$ , i.e., the eigenvector corresponding to the  $k$ th largest eigenvalue of  $\widehat{\mathbf{M}} \widehat{\mathbf{M}}^\top$
  - \*  $\hat{\mathbf{b}}_k = \mathbf{S}_\mathbf{X}^{-1/2} \hat{\mathbf{f}}_k$ 
    - $\hat{\mathbf{f}}_k$ : the right-singular vector corresponding to the  $k$ th largest singular value of  $\widehat{\mathbf{M}}$ , i.e., the eigenvector corresponding to the  $k$ th largest eigenvalue of  $\widehat{\mathbf{M}}^\top \widehat{\mathbf{M}}$

- Example: olive oil data
  - 572 olive oils
  - 10 features
    - \* **region** indicates the general region (in Italy) of origin.
    - \* **area** details the area of Italy.
    - \* Remaining variables are continuous valued and measure the percentage composition of 8 different fatty acids

```
options(digits=4)
Y = as.matrix(dslabs::olive[,3:4])
X = as.matrix(dslabs::olive[,5:10])
p = ncol(Y)
q = ncol(X)

# by definition
S_Y = cov(Y)
S_X = cov(X)
S_YX = cov(Y, X)
S_Y_sqrt = expm::sqrtm(S_Y)
S_X_sqrt = expm::sqrtm(S_X)
M = solve(S_Y_sqrt) %*% S_YX %*% solve(S_X_sqrt)
decomp1 = svd(M)
decomp1$d
A1 = solve(S_Y_sqrt) %*% decomp1$u
B1 = solve(S_X_sqrt) %*% decomp1$v
YA1 = scale(Y, scale=F) %*% A1 # canonical variates
XB1 = scale(X, scale=F) %*% B1 # canonical variates

# by cancel (not recommended if you try to know the specific values of canonical directions/variates)
decomp2 = cancel(x=X, y=Y)
decomp2$cor
A2 = decomp2$ycoef # identical to A1 up to a constant
B2 = decomp2$xcoef[, 1:min(p,q)] # identical to B1 up to a constant
YA2 = scale(Y, scale=F) %*% A2 # canonical variates
XB2 = scale(X, scale=F) %*% B2 # canonical variates

# comparison
A1/A2
B1/B2
t(A1) %*% S_Y %*% A1
t(B1) %*% S_X %*% B1
t(A2) %*% S_Y %*% A2
t(B2) %*% S_X %*% B2
```

- 
- Proportion of explained correlation
    - Determining  $r$ , the number of pairs of canonical variates to retain
    - $p \times r$  matrix  $\text{corr}(\mathbf{Y}, \mathbf{A}_r^\top \mathbf{Y})$  and  $q \times r$  matrix  $\text{corr}(\mathbf{X}, \mathbf{B}_r^\top \mathbf{X})$ 
      - \* The correlation matrix between  $\mathbf{Y}$  (or  $\mathbf{X}$ ) and canonical variates
      - \*  $\mathbf{A}_r = [\mathbf{a}_1, \dots, \mathbf{a}_r]$  and  $\mathbf{B}_r = [\mathbf{b}_1, \dots, \mathbf{b}_r]$
    - $\|\text{corr}(\mathbf{Y}, \mathbf{A}_r^\top \mathbf{Y})\|_F^2/p$  and  $\|\text{corr}(\mathbf{X}, \mathbf{B}_r^\top \mathbf{X})\|_F^2/q$ 
      - \* Proportion of explained correlation of  $\mathbf{Y}$  (or  $\mathbf{X}$ )
      - \*  $\|\cdot\|_F^2$ : squared Frobenius norm, i.e., sum of squared entries

```

Y = as.matrix(dslabs::olive[,3:6])
X = as.matrix(dslabs::olive[,7:10])
p = ncol(Y)
q = ncol(X)

S_Y = cov(Y)
S_X = cov(X)
S_YX = cov(Y, X)
S_Y_sqrt = expm::sqrtm(S_Y)
S_X_sqrt = expm::sqrtm(S_X)
M = solve(S_Y_sqrt) %*% S_YX %*% solve(S_X_sqrt)
decomp1 = svd(M)
A1 = solve(S_Y_sqrt) %*% decomp1$u
B1 = solve(S_X_sqrt) %*% decomp1$v
YA1 = scale(Y, scale=F) %*% A1
XB1 = scale(X, scale=F) %*% B1

cor(Y, YA1)
cor(X, XB1)

cumsum(colMeans(cor(Y, YA1)^2))
cumsum(colMeans(cor(X, XB1)^2))

```

- 
- Interpreting canonical variates

- $\text{corr}(\mathbf{Y}, \mathbf{A}_r^\top \mathbf{Y})$
- $\text{corr}(\mathbf{X}, \mathbf{B}_r^\top \mathbf{X})$
- $\text{corr}(\mathbf{Y}, \mathbf{B}_r^\top \mathbf{X})$
- $\text{corr}(\mathbf{X}, \mathbf{A}_r^\top \mathbf{Y})$

```

colnames(U1) = paste0("Y'a", seq_len(q))
colnames(V1) = paste0("X'b", seq_len(q))
cor(Y, YA1)
cor(Y, XB1)
cor(X, YA1)
cor(X, XB1)

```