PH 716 Applied Survival Analysis

Part 6: Diagnostics of Cox PH Models

Zhiyang Zhou (zhou67@uwm.edu, zhiyanggeezhou.github.io)

2026/04/09 01:17:36

Types of residuals

- Cox-Snell residuals: assessing the overall fit of the final model
- Martingale residuals: determining the functional form of a covariate included in the model
- Deviance residuals: detecting outliers
- Schoenfeld residuals: checking the appropriateness of the PH assumption

Cox-Snell residuals

- Cox-Snell residuals: $r_{i,\text{CS}} = \widehat{\Lambda}_{T_i}(\widetilde{T}_i)$
  - $\widehat{\Lambda}_{T_i}(\cdot)$: estimated $\Lambda_{T_i}(\cdot)$ given by the Cox PH model
  - (For uncensored subjects) $r_{i,\text{CS}} \overset{\text{iid}}{\approx} \exp(1)$ if the Cox PH model is correct
  - (For all subjects) $\widehat{\Lambda}_{H_i,\text{NA}}(\tilde{t}_i) \approx \tilde{t}_i$ if the Cox PH model is correct, where
    - $\widehat{\Lambda}_{H_i,\text{NA}}(\cdot)$: NA estimator of $\Lambda_{H_i}(\cdot)$ based on right-censored $\{(r_{i,\text{CS}}, \Delta_i) : i = 1, \ldots, n\}$
    - $H_i = \widehat{\Lambda}_{T_i}(T_i)$: a transformation of the true event time T_i
      - Note that $r_{i,\text{CS}} \neq H_i$ and instead $r_{i,\text{CS}} = \min\{H_i, \widehat{\Lambda}_{T_i}(C_i)\}$
- Cox-Snell residual plot
  - (For uncensored subjects) compare $r_{i,\text{CS}}$ to $\exp(1)$ samples via Q-Q plot
  - (For all subjects) plot $\widehat{\Lambda}_{H_i,\text{NA}}(\tilde{t}_i)$ against $\tilde{t}_i$
  - Used to diagnose poor model fit
  - No insight into how model assumptions are violated
- Theoretical notes
  - Inverse cdf theorem: arbitrary r.v. X with cdf $F_X(x) = \Pr(X \leq x) \Rightarrow F_X(X) \sim U(0,1)$
  - It follows that $T_i \overset{\text{independent}}{\sim} S_{T_i}(\cdot) \Rightarrow S_{T_i}(T_i) \overset{\text{iid}}{\sim} U(0,1) \Rightarrow \Lambda_{T_i}(T_i) = -\ln S_{T_i}(T_i) \overset{\text{iid}}{\sim} \exp(1)$
  - $\{(r_{i,\text{CS}}, \Delta_i) : i = 1, \ldots, n\}$ is a right-censored dataset
    - $r_{i,\text{CS}} = \min(H_i, \widehat{\Lambda}_{T_i}(C_i)) \Leftarrow \widetilde{T}_i = \min(T_i, C_i)$ and monotonically ascending $\widehat{\Lambda}_{T_i}(\cdot)$
      - $H_i = \widehat{\Lambda}_{T_i}(T_i) \approx \Lambda_{T_i}(T_i) \Rightarrow H_i \approx \exp(1) \Rightarrow \Lambda_{H_i}(\tilde{t}_i) \approx \tilde{t}_i$

Ex. 6.1 Multi-center acute leukemia study

- This multi-center acute leukemia study consists of 137 patients with acute myelocytic leukemia (AML) or acute lymphoblastic leukemia (ALL) aged 7 to 52 from March 1, 1984 to June 30, 1989 at four institutions.
- The disease-free survival time (`t2`) on study is defined as time (in days) to relapse or death
- `d3` is the disease free survival indicator: 1 - Dead or Relapsed, 0 - Alive Disease Free.
- Focus on effects of the following 9 covariates on disease-free survival:

- z1: Patient age in years.
- z2: Donor age in years.
- z3: Patient sex: 1 - Male, 0 - Female.
- z4: Doner sex: 1 - Male, 0 - Female.
- z5: Patient Cytomegalovirus (CMV) status: 1 - CMV positive, 0 - CMV negative.
- z6: Donor CMV status: 1 - CMV positive, 0 - CMV negative.
- z7: Waiting time to transplant in days.
- z8: French–American–British classification (FAB): 1 - FAB Grade 4 or 5 and AML, 0 - otherwise.
- Z9: Hospital: 1 - The Ohio State University, 2 - Alferd , 3 - St. Vincent, 4 - Hahnemann.
- z10: Methotrexate (MTX): used as a Graft-Versus-Host-Prophylactic 1 - Yes, 0 - No.

```r
options(digits=4)
library(survival)
# model fitting
data.ex61 = read.csv("bmt.csv")
fit.ex61 <- coxph(Surv(t2,d3) ~ z1+z2+z3+z4+z5+z6+z7+z8+z10, data=data.ex61)

# Cox-Snell residual
r.cs = data.ex61$d3-residuals(fit.ex61, type='martingale') # Cox-Snell

# Cox-Snell residual plot
set.seed(716)
exp.rnd = rexp(10000)
qqplot(
  x = exp.rnd, y = r.cs[as.logical(data.ex61$d3)],
  xlab = "Theoretical Quantiles", ylab = "Sample Quantiles"
)
qqline(r.cs[as.logical(data.ex61$d3)], distribution = qexp)
# Or
cum.haz.r.cs <- basehaz(coxph(Surv(r.cs, d3)~1, data=data.ex61), centered = FALSE)
plot(
  x=cum.haz.r.cs[,2], y=cum.haz.r.cs[,1],
  xlab='t', ylab='Cumulative hazard of r.cs'
)
abline(a=0,b=1,col='red')
```

Martingale residuals

- Martingale residuals: $r_{i,\mathrm{M}} = \Delta_i - r_{i,\mathrm{CS}}$
- To explore the proper functional form of the j_0th covariate, say $f_{j_0}(\cdot)$
  1. Fit a Cox PH model without the j_0th covariate $\lambda_{T_i}(t) = \lambda_0(t)\exp\{\sum_{j\neq j_0} f_j(x_{ij})\beta_j\}$
     - $f_j(\cdot)$: known proper functional form of the jth covariate
  2. Compute martingale residuals $r_{i,\mathrm{M}}$ for the above model
  3. Scatterplot of $r_{i,\mathrm{M}}$ against x_{ij_0} with a fitted locally estimated scatterplot smoothing (loess) curve
     - $f_{j_0}(\cdot)$ (approximately) proportional to the loess curve
  4. Update the model by adding $f_{j_0}(x_{ij_0})$ and check the scatterplot of updated $r_{i,\mathrm{M}}$ against x_{ij_0} with a fitted loess line
     - No further transformation needed If the new loess curve lies at the x-axis
- Theoretical notes:
  - loess: to predict $f(x_0)$ by an weighted average $\hat{f}(x_0) = \sum_{i=1}^n w_i y_i$, where w_i relies on the distance between x_i and x_0; a closer x_i enjoys a higher w_i.
  - Why is the residual bearing such a name?

    * Martingale: a stochastic process $M(t)$ such that $\mathrm{E}\{M(t)\} = 0$ and $\mathrm{E}\{M(t) \mid M(s)\} = M(s)$ for all $s < t$
    * $r_{i,\mathrm{M}}$ obtained by evaluating a martingale at $\tilde{t}_i$
  - Why would the martingale residuals reveal the correct functional forms of covariates?
    * With $r_{i,\mathrm{M}}$ obtained from the Cox PH model excluding the j_0th covariate, $E(r_{i,\mathrm{M}}) \approx (n_D/n)\{f_{j_0}(x_{ij_0}) - \text{constant}\}$ [KM, pp. 362]
      · n_D/n: the ratio of total number of events to total number of subjects
  - Zero-sum of martingale residuals: $\sum_i r_{i,\mathrm{M}} = 0$
    * Specific for Cox PH model with the Breslow estimator for the baseline cumulative hazard
    * Proof: $\sum_i r_{i,\mathrm{CS}} = \sum_i \sum_{k:t_k \le \tilde{t}_i} \frac{d_k \exp(\sum_j x_{ij}\hat{\beta}_j)}{\sum_{\ell \in \mathcal{R}(t_k)} \exp(\sum_j x_{\ell j}\hat{\beta}_j)} = \sum_k \sum_{i \in \mathcal{R}(t_k)} \frac{d_k \exp(\sum_j x_{ij}\hat{\beta}_j)}{\sum_{\ell \in \mathcal{R}(t_k)} \exp(\sum_j x_{\ell j}\hat{\beta}_j)} = \sum_k d_k = \sum_i \delta_i$

Revisit Ex. 6.1

```r
options(digits=4)
library(survival)
# [DM, pp. 208] a function to add the smooth curve and confidence limits to scatter plots
smoothSEcurve <- function(yy, xx) {
  # use after a call to "plot"
  # fit a lowess curve and 95% confidence interval curve
  # make list of x values
  xx.list <- min(xx) + ((0:100)/100)*(max(xx) - min(xx))
  # Then fit loess function through the points (xx, yy)
  # at the listed values
  yy.xx <- predict(loess(yy ~ xx, span = 1), se=T, newdata=data.frame(xx=xx.list))
  lines(yy.xx$fit ~ xx.list, lwd=2)
  lines(yy.xx$fit -
  qt(0.975, yy.xx$df)*yy.xx$se.fit ~ xx.list, lty=2)
  lines(yy.xx$fit +
  qt(0.975, yy.xx$df)*yy.xx$se.fit ~ xx.list, lty=2)
}

# model fitting without z1
data.ex61 = read.csv("bmt.csv")
fit.ex61 <- coxph(Surv(t2,d3) ~ z2+z3+z4+z5+z6+z7+z8+z10, data=data.ex61, ties = 'exact')

# Martingale residual plot (for the model without z1) vs. z1
r.m = residuals(fit.ex61, type='martingale')
sum(r.m)
plot(
  x=data.ex61$z1, y=r.m,
  main = 'Martingale residuals \n (for the model without z1) \n versus z1')
smoothSEcurve(r.m, data.ex61$z1) # indicating a cubic function?

# model fitting with a cubic function of z1
fit.ex61.1 <- coxph(Surv(t2,d3) ~ z1+I(z1^2)+I(z1^3)+z2+z3+z4+z5+z6+z7+z8+z10, data=data.ex61)
summary(fit.ex61.1)

# Martingale residual plot (for the model with a cubic function of z1) vs. z1
r.m.1 = residuals(fit.ex61.1, type='martingale')
plot(
  x=data.ex61$z1, y=r.m.1,
```

```
  main = 'Martingale residual \n (for the model with a cubic function of z1) \n versus z1')
smoothSEcurve(r.m.1, data.ex61$z1)
```

Deviance residuals

- Outlier: an observation for which the outcome is not sufficiently well predicted by the fitted model
- Deviance residuals: $r_{i,\mathrm{D}} = \mathrm{sign}(r_{i,M})\sqrt{-2\{r_{i,M} + \delta_i \ln(\delta_i - r_{i,M})\}}$
  - Symmetrically distributed with expected value 0 (if the fitted model is correct); de-skewed/transformed martingale residuals
    - $r_{i,\mathrm{D}} = 0 \Leftrightarrow r_{i,M} = 0$
    - Inflating $r_{i,\mathrm{D}}$ when $r_{i,M}$ is close to 1
    - Shrinking large negative $r_{i,M}$
  - Analogous to the deviance in GLMs
- Detecting outliers: plotting $r_{i,\mathrm{D}}$ against $\sum_{j=1}^{p} x_{ij}\hat{\beta}_j$ (called linear predictors or risk scores)
  - With moderate (or less) censoring, this plot should look like randomly-distributed noise without discernible pattern
  - Large absolute values of deviance residuals indicating observations that are poorly explained by the model, potentially pointing to outliers or influential points
    - 95% of absolute deviance residuals ≤ 2
    - 99.7% of absolute deviance residuals ≤ 3

Revisit Ex. 6.1

```
options(digits=4)
library(survival)
# [DM, pp. 208] a function to add the smooth curve and confidence limits
smoothSEcurve <- function(yy, xx) {
  # use after a call to "plot"
  # fit a lowess curve and 95% confidence interval curve
  # make list of x values
  xx.list <- min(xx) + ((0:100)/100)*(max(xx) - min(xx))
  # Then fit loess function through the points (xx, yy)
  # at the listed values
  yy.xx <- predict(loess(yy ~ xx, span = 1), se=T, newdata=data.frame(xx=xx.list))
  lines(yy.xx$fit ~ xx.list, lwd=2)
  lines(yy.xx$fit -
  qt(0.975, yy.xx$df)*yy.xx$se.fit ~ xx.list, lty=2)
  lines(yy.xx$fit +
  qt(0.975, yy.xx$df)*yy.xx$se.fit ~ xx.list, lty=2)
}

# model fitting
fit.ex61.1 <- coxph(
  Surv(t2,d3) ~ z1+I(z1^2)+I(z1^3)+z2+z3+z4+z5+z6+z7+z8+z10,
  data=data.ex61,
  x = T
)

# Two ways to calculate linear predictors
risk.score.1 = fit.ex61.1$x %*% coef(fit.ex61.1) # reliable
risk.score.2 = fit.ex61.1$linear.predictors # Unreliable due to the default centering
```

```r
sum((risk.score.1-risk.score.2)^2) # typically different

# Deviance residual plot vs. risk scores
r.d = residuals(fit.ex61.1, type='deviance')
plot(
  x=risk.score.1, y=r.d,
  main = 'Deviance residuals \n versus risk scores')
smoothSEcurve(yy=r.d, xx=risk.score.1)
abline(a=2,b=0,col='red')
abline(a=-2,b=0,col='red')
abline(a=3,b=0,col='red')
abline(a=-3,b=0,col='red')

(1:nrow(data.ex61))[abs(r.d) > 2] # Potential outliers
sum(abs(r.d) > 2)/nrow(data.ex61) # percent of r.d over 2
sum(abs(r.d) > 3)/nrow(data.ex61) # percent of r.d over 3
```

Schoenfeld residuals

- Definition (Schoenfeld residuals): for UNCENSORED subject i and the jth covariate,

$$r_{ij,\mathrm{S}} = x_{ij} - \bar{x}_{\cdot j}$$

  where

$$\bar{x}_{\cdot j} = \sum_{k \in \text{uncensored subjects}} \frac{z_{kj} \cdot \exp(\sum_{j=1}^{p} x_{kj}\beta_j)}{\sum_{\ell \in \mathcal{R}(\tilde{t}_k)} \exp(\sum_{j=1}^{p} x_{\ell j}\beta_j)}$$

- Key property [DM, Sec. 7.2.2]: Schoenfeld residuals are components of the score function, and thus $\sum_{i \in \text{uncensored subjects}} r_{ij,\mathrm{S}} = 0$ for each j
- Definition (scaled Schoenfeld residuals):

$$r^*_{ij,\mathrm{S}} = d \cdot r_{ij,\mathrm{S}} \cdot \mathrm{var}(\hat{\beta}_j)$$

  - d: total number of events
  - If the hazard ratio is constant over time (i.e., PH assumption holds), then $\mathrm{E}(r^*_{ij,\mathrm{S}}) + \beta_j$ is time-invariant
- Investigating the PH assumption
  - Direct approach (inconvenient to be implemented in `R`)
    - Plotting $r_{ij,\mathrm{S}}$ versus x_{ij} for each j
    - Interpretation: PH assumption holds $\Rightarrow$ points centered around 0
  - Standard approach
    - Plot $r^*_{ij,\mathrm{S}} + \hat{\beta}_j$ vs time t
    - Interpretation: PH assumption holds $\Rightarrow$ no systematic trend over time
  - Formal test (score test) via `survival::cox.zph`

Revisit Ex. 6.1

```r
options(digits=4)
library(survival)

# model fitting
fit.ex61.1 <- coxph(
```

```r
  Surv(t2,d3) ~ z1+I(z1^2)+I(z1^3)+z2+z3+z4+z5+z6+z7+z8+z10,
  data=data.ex61,
  x=T
)

# (unscaled) Schoenfeld residuals
r.s.unscaled = residuals(fit.ex61.1, type='schoenfeld')

# (scaled) Schoenfeld residual plot
plot(cox.zph(fit.ex61.1, transform="identity", terms=F, global=F))

# Score test
cox.zph(fit.ex61.1, transform="identity", terms=F, global=T)
## Identify which covariate(s) violate the PH assumption
## How to address the violation? See the next lecture on stratified Cox PH models.
```