

Lie Groups, Lie Algebras, and Representations

Reading Notes and Exercises

Zhiyuan Xu
Department of Electrical and Computer Engineering
McGill University

Contents

1	Matrix Lie Groups	3
1.1	Definitions	3
1.2	Examples	3
1.2.1	General and Special Linear Groups	3
1.2.2	Unitary and Orthogonal Groups	3
1.2.3	Generalized Orthogonal and Lorentz Groups	4
1.2.4	Symplectic Groups	4
1.2.5	The Euclidean and Poincaré Groups	5
1.2.6	The Heisenberg Group	5
1.2.7	The Groups $\mathbb{R}^*$, $\mathbb{C}^*$, S^1 , $\mathbb{R}$, and $\mathbb{R}^n$	5
1.2.8	The Compact Symplectic Group	5
1.3	Topological Properties	6
1.3.1	Compactness	6
1.3.2	Connectedness	6
1.3.3	The Topology of $SO(3)$	7
1.4	Homomorphisms	7
1.5	Lie Groups	8
1.6	Exercises	8
2	The Matrix Exponential	10
2.1	The Exponential of a Matrix	10
2.2	Computing the Exponential	10
2.3	The Matrix Logarithm	11
2.4	Further Properties of the Exponential	12
2.5	The Polar Decomposition	12
2.6	Exercises	16
3	Lie Algebras	18
3.1	Definitions and First Examples	18
3.2	Simple, Solvable, and Nilpotent Lie Algebras	20
3.3	The Lie Algebra of a Matrix Lie Group	21
3.4	Examples	21
3.5	Lie Group and Lie Algebra Homomorphisms	21
3.6	The Complexification of a Real Lie Algebra	21
3.7	The Exponential Map	21
3.8	Consequences of Theorem 3.42	21
3.9	Exercises	21
4	Basic Representation Theory	23
4.1	Representations	23
4.2	Examples of Representations	23
4.3	New Representations from Old	23
4.4	Complete Reducibility	23
4.5	Schur's Lemma	23
4.6	Representations of $\mathfrak{sl}(2; \mathbb{C})$	23
4.7	Group Versus Lie Algebra Representations	23
4.8	A Nonmatrix Lie Group	23
4.9	Exercises	23

1 Matrix Lie Groups

1.1 Definitions

Definition 1.1. The **general linear group** over the real numbers, denoted $GL(n, \mathbb{R})$, is the group of all $n \times n$ invertible matrices with real entries. The general linear group over the complex numbers, denoted $GL(n, \mathbb{C})$, is the group of all $n \times n$ invertible matrices with complex entries.

Definition 1.2. Let $M_n(\mathbb{C})$ denote the space of all $n \times n$ matrices with complex entries.

Definition 1.3. Let A_m be a sequence of complex matrices in $M_n(\mathbb{C})$. We say A_m **converges** to a matrix A if each entry of A_m converges to the corresponding entry of A as $m \rightarrow \infty$, i.e. if $(A_m)_{jk}$ converges to A_{jk} for all $1 \leq j, k \leq n$.

Note that Definition 1.3 means we can identify $M_n(\mathbb{C})$ with $\mathbb{C}^{n^2}$, and use the standard notion of convergence in $\mathbb{C}^{n^2}$. All topological properties of a matrix Lie group G will be considered with respect to the topology G inherits as a subset of $M_n(\mathbb{C}) \cong \mathbb{C}^{n^2}$.

We now consider subgroups of $GL(n, \mathbb{C})$, which are subsets $G \subset GL(n, \mathbb{C})$ such that $Id \in G$, and for all $A, B \in G$ we have $AB \in G$ and $A^{-1} \in G$.

Definition 1.4. A **matrix Lie group** is a subgroup G of $GL(n, \mathbb{C})$ with the property: if A_m is any sequence of matrices in G , and A_m converges to some matrix A , then either A is in G or A is not invertible.

This property means that G is a closed subset of $GL(n, \mathbb{C})$. So this definition is equivalent to saying a matrix Lie group is a closed subgroup of $GL(n, \mathbb{C})$. However, G is not necessarily closed in $M_n(\mathbb{C})$. For this to be true, we require a stronger property: if A_m is any sequence of matrices in G and A_m converges to some matrix A , then $A \in G$.

1.2 Examples

We introduce some important examples of (matrix) Lie groups, including the classical groups.

1.2.1 General and Special Linear Groups

The general linear groups $GL(n, \mathbb{R})$ and $GL(n, \mathbb{C})$, defined in 1.1, are matrix Lie groups.

The special linear groups $SL(n, \mathbb{R})$ and $SL(n, \mathbb{C})$ is the group of $n \times n$ invertible matrices (with real or complex entries) having determinant 1. They are matrix Lie groups.

1.2.2 Unitary and Orthogonal Groups

A $n \times n$ complex matrix A is **unitary** if its column vectors are orthonormal, that is,

$$\sum_{l=1}^n \bar{A}_{lj} A_{lk} = \delta_{jk}$$

where δ_{jk} is the Kronecker delta. This can be rewritten as

$$\sum_{l=1}^n (A^*)_{jl} A_{lk} = \delta_{jk}$$

where A^* is the adjoint of A , defined by $A^* = \bar{A}^t$. A is unitary if and only if $A^* = A^{-1}$.

The **unitary group** $U(n)$ is the collection of all unitary matrices. The **special unitary group** $SU(n)$ is the subgroup of $U(n)$ consisting of unitary matrices with determinant 1. $U(n)$ and $SU(n)$ are matrix Lie groups.

A $n \times n$ real matrix A is **orthogonal** if its column vectors are orthonormal. The equivalent conditions are the same as before, except that when A is real, $A^* = A^t$. A is orthogonal if and only if $A^t = A^{-1}$, if and only if A preserves the inner product on $\mathbb{R}^n$.

The **orthogonal group** $O(n)$ is the collection of all orthogonal matrices. The **special orthogonal group** $SO(n)$ is the set of $n \times n$ orthogonal matrices with determinant 1. Elements of $SO(n)$ are rotations, while the elements of $O(n)$ are either rotations or combinations of rotations and reflections.

Consider the bilinear form $(\cdot, \cdot)$ on $\mathbb{C}^n$ defined by

$$(x, y) = \sum_j x_j y_j$$

Note that this form is not an inner product, since it is symmetric, not conjugate-symmetric. The **complex orthogonal group** $O(n, \mathbb{C})$ is the set of all $n \times n$ complex matrices A that preserve this form. The group $SO(n, \mathbb{C})$ is the set of all matrices in $O(n, \mathbb{C})$ with determinant 1. $O(n, \mathbb{C})$ and $SO(n, \mathbb{C})$ are matrix Lie groups.

1.2.3 Generalized Orthogonal and Lorentz Groups

Consider $\mathbb{R}^{n+k}$, with n and k positive integers. Define a symmetric bilinear form $[\cdot, \cdot]_{n,k}$ on $\mathbb{R}^{n+k}$ by

$$[x, y]_{n,k} = x_1 y_1 + \cdots + x_n y_n - x_{n+1} y_{n+1} - \cdots - x_{n+k} y_{n+k}$$

The **generalized orthogonal group** $O(n, k)$ is the set of $(n+k) \times (n+k)$ real matrices A that preserve this form, i.e. $[Ax, Ay]_{n,k} = [x, y]_{n,k}$. It is a matrix Lie group. In particular, the **Lorentz group** is $O(3, 1)$. $SO(n, k)$ is the subgroup of $O(n, k)$ consisting of elements with determinant 1.

For any $A \in O(n, k)$, we have $\det A = \pm 1$.

1.2.4 Symplectic Groups

Define the skew-symmetric bilinear form ω on $\mathbb{R}^{2n}$ by

$$\omega(x, y) = \sum_{j=1}^n (x_j y_{n+j} - x_{n+j} y_j)$$

The **real symplectic group** $Sp(n, \mathbb{R})$ is the set of all $2n \times 2n$ real matrices A which preserve ω . It is a closed subgroup of $GL(2n, \mathbb{R})$, hence a matrix Lie group. $\det A = \pm 1$ for all $A \in Sp(n, \mathbb{R})$; in fact, $\det A = 1$.

Using the same formula, we can define a bilinear form ω on $\mathbb{C}^{2n}$. Over $\mathbb{C}$, we have the relation $\omega(z, w) = (z, \Omega w)$, where $(\cdot, \cdot)$ is the complex bilinear form from 1.2.2. The **complex symplectic group** $Sp(n, \mathbb{C})$ is the set of all $2n \times 2n$ complex matrices which preserve this form. We can show that for all $A \in Sp(n, \mathbb{C})$, $\det A = \pm 1$, and in fact, $\det A = 1$.

The **compact symplectic group** $Sp(n)$ is defined as

$$Sp(n) = Sp(n, \mathbb{C}) \cap U(2n)$$

That is to say, $Sp(n)$ is the group of $2n \times 2n$ matrices that preserve both the inner product and the bilinear form ω .

1.2.5 The Euclidean and Poincaré Groups

The **Euclidean group** $E(n)$ is the group of all transformations of $\mathbb{R}^n$ that can be expressed as a composition of a translation and an orthogonal linear transformation. Elements of $E(n)$ are written as pairs $\{x, R\}$ with $x \in \mathbb{R}^n$ and $R \in O(n)$, and $\{x, R\}$ acts on $\mathbb{R}^n$ by

$$\{x, R\}y = Ry + x$$

The product operation for $E(n)$ is

$$\{x_1, R_1\}\{x_2, R_2\} = \{x_1 + R_1x_2, R_1R_2\}$$

The inverse of an element of $E(n)$ is

$$\{x, R\}^{-1} = \{-R^{-1}x, R^{-1}\}$$

$E(n)$ is not a subgroup of $GL(n, \mathbb{R})$, but it is isomorphic to a closed subgroup of $GL(n+1, \mathbb{R})$.

The **Poincaré group** $P(n, 1)$ is the group of all transformations of $\mathbb{R}^{n+1}$ of the form

$$T = T_x A$$

with $x \in \mathbb{R}^{n+1}$ and $A \in O(n, 1)$.

1.2.6 The Heisenberg Group

The **Heisenberg group** H is the set of all 3×3 real matrices

$$A = \begin{pmatrix} 1 & a & b \\ 0 & 1 & c \\ 0 & 0 & 1 \end{pmatrix}$$

where $a, b, c \in \mathbb{R}$. H is a subgroup of $GL(3, \mathbb{R})$.

1.2.7 The Groups $\mathbb{R}^*$, $\mathbb{C}^*$, S^1 , $\mathbb{R}$, and $\mathbb{R}^n$

These groups are not defined as groups of matrices, but can be thought of as such.

- The group $\mathbb{R}^*$ of non-zero real numbers under multiplication is isomorphic to $GL(1, \mathbb{R})$
- The group $\mathbb{C}^*$ of non-zero complex numbers under multiplication is isomorphic to $GL(1, \mathbb{C})$
- The group S^1 of complex numbers with absolute value 1 is isomorphic to $U(1)$
- The group $\mathbb{R}$ under addition is isomorphic to $GL(1, \mathbb{R})^+$, the 1×1 matrices with positive determinant
- The group $\mathbb{R}^n$ under vector addition is isomorphic to the group of diagonal real matrices with positive diagonal entries

1.2.8 The Compact Symplectic Group

We try to understand the structure of $Sp(n)$, and show that it can be understood as the "unitary group over the quaternions".

Define a conjugate-linear map $J : \mathbb{C}^{2n} \rightarrow \mathbb{C}^{2n}$ by

$$J(\alpha, \beta) = (-\bar{\beta}, \bar{\alpha})$$

where $\alpha, \beta \in \mathbb{C}^n$ and $(\alpha, \beta) \in \mathbb{C}^{2n}$. For all $z, w \in \mathbb{C}^{2n}$, we have

$$\omega(z, w) = \langle Jz, w \rangle$$

Furthermore, for all $z, w \in \mathbb{C}^{2n}$, we have

$$\langle Jz, w \rangle = -\langle Jw, z \rangle$$

and

$$J^2 = -I$$

Proposition 1.5. *If $U \in U(2n)$, then $U \in Sp(n)$ if and only if U commutes with J .*

Theorem 1.6. *$U \in Sp(n)$ if and only if there exists an orthonormal basis $u_1, \dots, u_n, v_1, \dots, v_n$ for $\mathbb{C}^{2n}$ such that the following properties hold:*

1. $Ju_j = v_j$
2. For some real numbers $\theta_1, \dots, \theta_n$, we have

$$\begin{aligned} Uu_j &= e^{i\theta_j}u_j \\ Uv_j &= e^{-i\theta_j}v_j \end{aligned}$$

3. We have

$$\begin{aligned} \omega(u_j, u_k) &= \omega(v_j, v_k) = 0 \\ \omega(u_j, v_k) &= \delta_{jk} \end{aligned}$$

Lemma 1.7. *Suppose V is a complex subspace of $\mathbb{C}^{2n}$ that is invariant under the conjugate-linear map J . Then the orthogonal complement $V^\perp$ of V (with respect to the inner product $\langle \cdot, \cdot \rangle$) is also invariant under J . Furthermore, V and $V^\perp$ are orthogonal with respect to ω , i.e.*

$$\omega(z, w) = 0$$

for all $z \in V$ and $w \in V^\perp$.

1.3 Topological Properties

1.3.1 Compactness

Definition 1.8. *A matrix Lie group $G \subset GL(n, \mathbb{C})$ is **compact** if it is compact in the usual topological sense as a subset of $M_n(\mathbb{C}) \cong \mathbb{R}^{2n^2}$.*

By the Heine-Borel theorem, a matrix Lie group G is compact if and only if it is closed and bounded.

$O(n)$, $SO(n)$, $U(n)$, $SU(n)$, and $Sp(n)$ are compact.

1.3.2 Connectedness

Definition 1.9. *A matrix Lie group G is **connected** if for all $A, B \in G$, there exists a continuous path $A(t)$, $a \leq t \leq b$, lying in G with $A(a) = A$ and $A(b) = B$. For any matrix Lie group G , the **identity component** of G , denoted G_0 , is the set of $A \in G$ for which there exists a continuous path $A(t)$, $a \leq t \leq b$, lying in G with $A(a) = I$ and $A(b) = A$.*

This definition is actually called **path connected** in topology. A matrix Lie group is connected if and only if it is path-connected.

To show that a matrix Lie group G is connected, it suffices to show that each $A \in G$ can be connected to the identity by a continuous path lying in G .

Proposition 1.10. *If G is a matrix Lie group, the identity component G_0 of G is a normal subgroup of G .*

G_0 is closed, and hence a matrix Lie group.

Proposition 1.11. *$GL(n, \mathbb{C})$ is connected for all $n \geq 1$.*

Proposition 1.12. *$SL(n, \mathbb{C})$ is connected for all $n \geq 1$.*

Proposition 1.13. *$U(n)$, $SU(n)$, and $SO(n)$ are connected for all $n \geq 1$.*

1.3.3 The Topology of $SO(3)$

Definition 1.14. *The **real projective space** of dimension n , denoted $\mathbb{R}P^n$, is the set of lines through the origin in $\mathbb{R}^{n+1}$. Since each line through the origin intersects the unit sphere exactly twice, we may think of $\mathbb{R}P^n$ as the unit sphere S^n with "antipodal" points u and $-u$ identified.*

Points in $\mathbb{R}P^n$ can be thought of as pairs $\{u, -u\}$ with $u \in S^n$. There is a natural map $\pi : S^n \rightarrow \mathbb{R}P^n$

$$\pi(u) = \{u, -u\}$$

A distance function can be defined on $\mathbb{R}P^n$ by

$$d(\{u, -u\}, \{v, -v\}) = \min(d(u, v), d(u, -v))$$

The space $\mathbb{R}P^n$ is the upper hemisphere in S^n with antipodal points on the equator identified. $\mathbb{R}P^n$ is the closed unit ball $B^n \subset \mathbb{R}^n$, with antipodal points on the boundary of B^n identified.

Proposition 1.15. *There is a continuous bijection between $SO(3)$ and $\mathbb{R}P^3$.*

$\mathbb{R}P^n$ is not simply connected, and hence $SO(3)$ is not simply connected.

1.4 Homomorphisms

Definition 1.16. *Let G and H be matrix Lie groups. A map Φ from G to H is called a **Lie group homomorphism** if*

1. Φ is a group homomorphism, and
2. Φ is continuous.

*If, in addition, Φ is one-to-one and onto and the inverse map Φ^{-1} is continuous, then Φ is called a **Lie group isomorphism**.*

If G and H are matrix Lie groups, and there exists a Lie group isomorphism from G to H , then G and H are isomorphic, $G \cong H$.

The relationship between $SU(2)$ and $SO(3)$: they are not isomorphic, but there exists a Lie group homomorphism $\Phi : SU(2) \rightarrow SO(3)$ that is two-to-one and onto.

Proposition 1.17. *The map $U \mapsto \Phi_U$ is a 2-1 and onto map of $SU(2)$ to $SO(3)$, with kernel equal to $\{I, -I\}$.*

$SU(2)$ is homeomorphic to S^3 . The proposition shows that $SO(3)$ is homeomorphic to $\mathbb{R}P^3$, that is, S^3 with antipodal points identified.

1.5 Lie Groups

The text first gives a very brief discussion of manifolds and smooth manifolds.

Definition 1.18. A **Lie group** is a smooth manifold G which is also a group, and such that the group product $G \times G \rightarrow G$ and the inverse map $G \rightarrow G$ are smooth.

Every matrix Lie group is a Lie group, but the reverse is not true. However, we focus on matrix Lie groups.

Usually, we call a map Φ between two Lie groups a Lie group homomorphism if Φ is a group homomorphism and Φ is smooth.

Every matrix Lie group G is a manifold, so G must be locally path connected, so G is connected if and only if it is path connected.

1.6 Exercises

Exercise 1.7 Using Theorem 1.6, show that $Sp(n)$ is connected and that every element of $Sp(n)$ has determinant 1.

Proof. Let $U \in Sp(n)$. Note that the matrix U represents a linear transformation $L_u : \mathbb{C}^{2n} \rightarrow \mathbb{C}^{2n}$. By Theorem 1.6, there exists an orthonormal basis $u_1, \dots, u_n, v_1, \dots, v_n$ for $\mathbb{C}^{2n}$ such that

1. $Ju_j = v_j$

2. for some $\theta_1, \dots, \theta_n \in \mathbb{R}$,

$$Uu_j = e^{i\theta_j}u_j$$

$$Uv_j = e^{-i\theta_j}v_j$$

3. and we have

$$\omega(u_j, u_k) = \omega(v_j, v_k) = 0$$

$$\omega(u_j, v_k) = \delta_{jk}$$

The second point is the most important. Notice that it implies $e^{i\theta_j}$ are eigenvalues of U with eigenvectors u_j , and $e^{-i\theta_j}$ are eigenvalues of U with eigenvectors v_j .

Denote $B = \{u_1, \dots, u_n, v_1, \dots, v_n\}$. B is a basis for $\mathbb{C}^{2n}$ consisting entirely of eigenvectors of U , so U is diagonalizable. Thus there exists an invertible matrix $Q \in M_{2n}(\mathbb{C})$ such that

$$QUQ^{-1} = \text{diag}(e^{i\theta_1}, \dots, e^{i\theta_n}, e^{-i\theta_1}, \dots, e^{-i\theta_n})$$

and so

$$U = Q^{-1} \text{diag}(e^{i\theta_1}, \dots, e^{i\theta_n}, e^{-i\theta_1}, \dots, e^{-i\theta_n})Q$$

Now, we make a slight modification and define

$$U(t) = Q^{-1} \text{diag}(e^{i(1-t)\theta_1}, \dots, e^{i(1-t)\theta_n}, e^{-i(1-t)\theta_1}, \dots, e^{-i(1-t)\theta_n})Q$$

for $0 \leq t \leq 1$. $U(t)$ is continuous, $U(0) = U$, and $U(1) = Q^{-1}IQ = I$. By applying the converse of Theorem 1.6 to $U(t)$, it is clear that $U(t) \in Sp(n)$ for all $0 \leq t \leq 1$.

Hence $Sp(n)$ is connected.

For any $U \in Sp(n)$,

$$\begin{aligned} \det(U) &= \det(Q^{-1} \text{diag}(e^{i\theta_1}, \dots, e^{i\theta_n}, e^{-i\theta_1}, \dots, e^{-i\theta_n})Q) \\ &= \det(Q^{-1}) \det(\text{diag}(e^{i\theta_1}, \dots, e^{i\theta_n}, e^{-i\theta_1}, \dots, e^{-i\theta_n})) \det(Q) \\ &= \frac{1}{\det(Q)} e^{i\theta_1} \dots e^{i\theta_n} e^{-i\theta_1} \dots e^{-i\theta_n} \det(Q) \\ &= 1 \end{aligned}$$

□

Exercise 1.9 Suppose a is an irrational real number. Show that the set E_a of numbers of the form $e^{2\pi ina}$, $n \in \mathbb{Z}$, is dense in the unit circle S^1 .

Proof. We follow the given hint, and divide S^1 into N equally sized bins of length $\frac{2\pi}{N}$. E_a has an infinite number of elements, and N may be large but finite. There must be at least one bin that has an infinite number of elements of E_a (otherwise, if every bin has finite number of elements and N is finite, then E_a would be finite).

Now, choose a bin with infinitely many elements of E_a . In this bin, it is possible to choose two distinct elements of E_a of the form $e^{2\pi ina}$ and $e^{2\pi ima}$, with $n, m \in \mathbb{R}$, $n \neq m$.

We use the fact that E_a is a subgroup of S^1 . $e^{2\pi ina}, e^{2\pi ima} \in E_a$, so the inverse $e^{-2\pi ima} \in E_a$, so $e^{2\pi i(n-m)a} \in E_a$, and so $(e^{2\pi i(n-m)a})^k = e^{2k\pi i(n-m)a} \in E_a$ for any $k \in \mathbb{Z}$.

By taking N to be large, we can force $e^{2\pi ina}$ and $e^{2\pi ima}$ to be arbitrarily close to each other on S^1 . Then $e^{2k\pi i(n-m)a}$ is a rotation by an arbitrarily small angle. Putting it to the power of k allows us to approximate any point on S^1 by an element of E_a .

More formally, let $e^{i\theta} \in S^1$ for any $\theta \in \mathbb{R}$.

$$\begin{aligned} |e^{i\theta} - e^{2k\pi i(n-m)a}| &= \left| e^{2k\pi i(n-m)a} \left(\frac{e^{i\theta}}{e^{2k\pi i(n-m)a}} - 1 \right) \right| \\ &= \left| \frac{e^{i\theta}}{e^{2k\pi i(n-m)a}} - 1 \right| \\ &= |e^{i(\theta - 2k\pi(n-m)a)} - 1| \\ &= |e^{i(\theta - 2k\pi(n-m)a)} - e^0| \end{aligned}$$

By continuity, $\forall \varepsilon > 0$, $\exists \delta > 0$ such that $|\theta - 2k\pi(n-m)a| < \delta \Rightarrow |e^{i(\theta - 2k\pi(n-m)a)} - e^0| < \varepsilon$. So $\forall \varepsilon > 0$, we can choose N large, and a suitable k such that

$$|e^{i\theta} - e^{2k\pi i(n-m)a}| < \varepsilon$$

Hence E_a is dense in S^1 . □

2 The Matrix Exponential

2.1 The Exponential of a Matrix

If X is a $n \times n$ matrix, we define the **exponential** of X , denoted e^X , by the power series

$$e^X = \sum_{m=0}^{\infty} \frac{X^m}{m!} \quad (2.1)$$

where X^0 is the identity matrix I and X^m is the repeated matrix product of X with itself.

Proposition 2.1. *The series 2.1 converges for all $X \in M_n(\mathbb{C})$ and e^X is a continuous function of X .*

Definition 2.2. *For any $X \in M_n(\mathbb{C})$, we define the **Hilbert-Schmidt** norm of X ,*

$$\|X\| = \left(\sum_{j,k=1}^n |X_{jk}|^2 \right)^{1/2}$$

The Hilbert-Schmidt norm may be computed in a basis-independent way as

$$\|X\| = (\text{tr}(X^* X))^{1/2}$$

This norm satisfies

$$\|X + Y\| \leq \|X\| + \|Y\|$$

$$\|XY\| \leq \|X\| \|Y\|$$

for all $X, Y \in M_n(\mathbb{C})$.

Proposition 2.3. *Let X and Y be arbitrary $n \times n$ matrices. Then we have the following:*

1. $e^0 = I$
2. $(e^X)^* = e^{X^*}$
3. e^X is invertible and $(e^X)^{-1} = e^{-X}$
4. $e^{(\alpha+\beta)X} = e^{\alpha X} e^{\beta X}$ for all α and β in $\mathbb{C}$
5. If $XY = YX$, then $e^{X+Y} = e^X e^Y = e^Y e^X$
6. If $C \in GL(N, \mathbb{C})$, then $e^{CXC^{-1}} = Ce^XC^{-1}$

Proposition 2.4. *Let X be a $n \times n$ complex matrix. Then e^{tX} is a smooth curve in $M_n(\mathbb{C})$ and*

$$\frac{d}{dt} e^{tX} = X e^{tX} = e^{tX} X$$

In particular,

$$\left. \frac{d}{dt} e^{tX} \right|_{t=0} = X$$

2.2 Computing the Exponential

The book explains some methods for computing the matrix exponential. If X is diagonalizable, then $X = CDC^{-1}$, and $e^X = Ce^DC^{-1}$. If X is nilpotent, ($X^k = 0$ for some k), then the power series of e^X terminates. Finally, every matrix X can be written uniquely as $X = S + N$ where S is diagonalizable, N is nilpotent, and $SN = NS$. Then $e^X = e^{S+N} = e^S e^N$.

2.3 The Matrix Logarithm

We want to define a matrix logarithm, which should be an inverse function to the matrix exponential. The simplest way is by a power series. Recall that in the case of complex numbers,

Lemma 2.5. *The function*

$$\log z = \sum_{m=1}^{\infty} (-1)^{m+1} \frac{(z-1)^m}{m}$$

is defined and holomorphic in a circle of radius 1 about $z = 1$. For all z with $|z - 1| < 1$,

$$e^{\log z} = z$$

For all u with $|u| < \log 2$, we have $|e^u - 1| < 1$ and

$$\log e^u = u$$

Definition 2.6. *For an $n \times n$ matrix A , define $\log A$ by*

$$\log A = \sum_{m=1}^{\infty} (-1)^{m+1} \frac{(A - I)^m}{m}$$

whenever the series converges.

Theorem 2.7. *The function*

$$\log A = \sum_{m=1}^{\infty} (-1)^{m+1} \frac{(A - I)^m}{m}$$

is defined and continuous on the set of all $n \times n$ complex matrices A with $\|A - I\| < 1$. For all $A \in M_n(\mathbb{C})$ with $\|A - I\| < 1$,

$$e^{\log A} = A$$

For all $X \in M_n(\mathbb{C})$ with $\|X\| < \log 2$, $\|e^X - I\| < 1$ and

$$\log e^X = X$$

Notice from the last sentence that it is not true that $\log e^X = X$ whenever the series for the logarithm converges.

Proposition 2.8. *There exists a constant c such that for all $n \times n$ matrices B with $\|B\| < 1/2$, we have*

$$\|\log(I + B) - B\| \leq c\|B\|^2$$

More concisely, the proposition can be stated as

$$\log(I + B) = B + O(\|B\|^2)$$

Theorem 2.9. *Every invertible $n \times n$ matrix can be expressed as e^X for some $X \in M_n(\mathbb{C})$.*

2.4 Further Properties of the Exponential

Theorem 2.10 (Lie Product Formula). *For all $X, Y \in M_n(\mathbb{C})$, we have*

$$e^{X+Y} = \lim_{m \rightarrow \infty} \left(e^{\frac{X}{m}} e^{\frac{Y}{m}} \right)^m$$

Theorem 2.11. *For any $X \in M_n(\mathbb{C})$, we have*

$$\det(e^X) = e^{\text{tr}(X)}$$

Definition 2.12. *A function $A : \mathbb{R} \rightarrow GL(n, \mathbb{C})$ is called a **one-parameter subgroup** of $GL(n, \mathbb{C})$ if*

1. *A is continuous,*
2. *$A(0) = I$,*
3. *$A(t+s) = A(t)A(s)$ for all $t, s \in \mathbb{R}$.*

Theorem 2.13 (One-Parameter Subgroups). *If $A(\cdot)$ is a one-parameter subgroup of $GL(n, \mathbb{C})$, there exists a unique $n \times n$ complex matrix X such that*

$$A(t) = e^{tX}$$

Lemma 2.14. *Fix some ε with $\varepsilon < \log 2$. Let $B_{\varepsilon/2}$ be the ball of radius $\varepsilon/2$ around the origin in $M_n(\mathbb{C})$, and let $U = \exp(B_{\varepsilon/2})$. Then every $B \in U$ has a unique square root C in U , given by $C = \exp(\frac{1}{2} \log B)$.*

Proposition 2.15. *The exponential map is an infinitely differentiable map of $M_n(\mathbb{C})$ into $M_n(\mathbb{C})$.*

2.5 The Polar Decomposition

The polar decomposition for a nonzero complex number z states that z can be written uniquely as $z = up$, where $|u| = 1$ and p is real and positive. Since p is real and positive, it can be written as $p = e^x$ for a unique real number x . This gives the polar decomposition for z ,

$$z = ue^x$$

with $x \in \mathbb{R}$ and $|u| = 1$.

We want to establish a similar polar decomposition for $GL(n, \mathbb{C})$, then for various subgroups. If P is a self-adjoint $n \times n$ matrix ($P^* = P$), we say that P is **positive** if $\langle v, Pv \rangle > 0$ for all nonzero $v \in \mathbb{C}^n$. A self-adjoint matrix P is positive if and only if all the eigenvalues of P are positive.

Theorem 2.16. *Polar decomposition for $GL(n, \mathbb{C})$.*

1. *Every $A \in GL(n, \mathbb{C})$ can be written uniquely in the form*

$$A = UP$$

where U is unitary and P is self-adjoint and positive.

2. *Every self-adjoint positive matrix P can be written uniquely in the form*

$$P = e^X$$

with X self-adjoint. Conversely, if X is self-adjoint, then e^X is self-adjoint and positive.

3. If we decompose each $A \in GL(n, \mathbb{C})$ (uniquely) as

$$A = Ue^X$$

with U unitary and X self-adjoint, then U and X depend continuously on A .

Lemma 2.17. *If Q is a self-adjoint, positive matrix, then Q has a unique self-adjoint, positive square root.*

Proof. Recall that a matrix Q is self-adjoint (Hermitian) if $Q^* = Q$, and unitary if $Q^* = Q^{-1}$. From Appendix A, we know that a matrix Q self-adjoint $\Rightarrow$ normal $\Rightarrow$ diagonalizable, and it is possible to find an orthonormal basis of eigenvectors for Q . Here, we can write $Q = UDU^{-1}$ where U can be taken to be unitary.

Let Q be a self-adjoint, positive matrix, then Q can be written as

$$Q = U \begin{pmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{pmatrix} U^{-1}$$

with U unitary. The λ_j are eigenvalues corresponding to the orthonormal basis of eigenvectors of Q . Since Q is self-adjoint and positive, $\lambda_j > 0$ for all j .

Thus, we can construct a square root of Q as

$$Q^{1/2} = U \begin{pmatrix} \lambda_1^{1/2} & & \\ & \ddots & \\ & & \lambda_n^{1/2} \end{pmatrix} U^{-1}$$

$Q^{1/2}$ is still self-adjoint and positive. This proves the existence of the square root.

Consider P a self-adjoint, positive matrix with eigenvalues λ_j and eigenvectors v_j , and so

$$Pv_j = \lambda_j v_j$$

Notice that the eigenspaces of P^2 are the same as the eigenspaces of P , with the eigenvalues of P^2 equal to the squares of the eigenvalues of P . This is because

$$P^2 v_j = P(Pv_j) = P(\lambda_j v_j) = \lambda_j (\lambda_j v_j) = \lambda_j^2 v_j$$

The mapping $x \mapsto x^2$ is injective for $x \in \mathbb{R}^{\geq 0}$. So eigenspaces with distinct eigenvalues remain with distinct eigenvalues after squaring.

If a self-adjoint positive Q has a self-adjoint positive square root P , then the eigenspaces of P must be the same as the eigenspaces of Q , and the eigenvalues of P must be the positive square roots of the eigenvalues of Q . Thus, P is uniquely determined by Q , proving uniqueness. $\square$

Proof. of Theorem 2.16.

1. Note that if $A = UP$, then $A^*A = P^*U^*UP = PU^{-1}UP = P^2$. For any matrix A , the matrix A^*A is self-adjoint, since $(A^*A)^* = A^*A^{**} = A^*A$. If, in addition, A is invertible, then for all nonzero $v \in \mathbb{C}^n$, we have

$$\langle v, A^*Av \rangle = \langle Av, Av \rangle > 0$$

where the last inequality is due to the definition of the inner product and v being nonzero. So A^*A is positive. For all invertible A , let us define P by

$$P = (A^*A)^{1/2}$$

where the square root is the unique self-adjoint positive square root from Lemma 2.17. Then, we define

$$U = AP^{-1} = A[(A^*A)^{1/2}]^{-1}$$

P is self-adjoint and positive by construction, and $A = UP$ by the definition of U , so it remains only to check that U is unitary. To do this, we check

$$U^*U = [(A^*A)^{1/2}]^{-1}A^*A[(A^*A)^{1/2}]^{-1}$$

noting that the inverse of a self-adjoint positive matrix is self-adjoint. Since A^*A is the square of $(A^*A)^{1/2}$, we have

$$U^*U = [(A^*A)^{1/2}]^{-1}(A^*A)^{1/2}(A^*A)^{1/2}[(A^*A)^{1/2}]^{-1} = I$$

showing that U is unitary. This proves existence.

We already showed that if $A = UP$, then $A^*A = P^2$ where A^*A is self-adjoint positive. The uniqueness of P follows from Lemma 2.17. The uniqueness of U then follows from the definition $U = AP^{-1}$.

2. The existence and uniqueness of the decomposition is proven in the same way as in Lemma 2.17. Instead of the square root function, we use the logarithm function, which is a bijection between $(0, \infty)$ and $\mathbb{R}$.

Let P be a self-adjoint matrix, so it is normal, so it is diagonalizable, and so P can be written as

$$P = U \begin{pmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{pmatrix} U^{-1}$$

where U can be taken to be unitary, and λ_j is an orthonormal basis of P . Since P is positive, $\lambda_j > 0$ for all j . Next, we define

$$\log P = U \begin{pmatrix} \log \lambda_1 & & \\ & \ddots & \\ & & \log \lambda_n \end{pmatrix} U^{-1} =: X$$

Then,

$$P = \exp \left(U \begin{pmatrix} \log \lambda_1 & & \\ & \ddots & \\ & & \log \lambda_n \end{pmatrix} U^{-1} \right) = e^X$$

We can check that X is self-adjoint

$$\begin{aligned} X^* &= (U^{-1})^* \text{diag}(\log \lambda_1, \dots, \log \lambda_n)^* U^* \\ &= (U^*)^{-1} \text{diag}(\log \lambda_1, \dots, \log \lambda_n)^* U^* \\ &= U \text{diag}(\log \lambda_1, \dots, \log \lambda_n) U^{-1} \\ &= X \end{aligned}$$

So P can be written as e^X where X is self-adjoint, which proves existence.

Moreover, the mapping $P \mapsto \log P$ is a bijection, so every P corresponds to a unique X such that $P = e^X$, which proves uniqueness.

For a self-adjoint matrix X , we know from before that we can write

$$X = U \begin{pmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{pmatrix} U^{-1}$$

with U unitary, and λ_j an orthonormal basis of X . Define

$$e^X = U \begin{pmatrix} e_1^\lambda & & \\ & \ddots & \\ & & e_n^\lambda \end{pmatrix} U^{-1}$$

e^X is self-adjoint because

$$\begin{aligned} (e^X)^* &= (U^{-1})^* \text{diag}(e_1^\lambda, \dots, e_n^\lambda)^* U^* \\ &= (U^*)^{-1} \text{diag}(e_1^\lambda, \dots, e_n^\lambda)^* U^* \\ &= U \text{diag}(e_1^\lambda, \dots, e_n^\lambda) U^{-1} \\ &= e^X \end{aligned}$$

Furthermore, e^X is positive because

$$\begin{aligned} \langle v, e^X v \rangle &= \langle v, U \text{diag}(e_1^\lambda, \dots, e_n^\lambda) U^{-1} v \rangle \\ &= \langle U w, U \text{diag}(e_1^\lambda, \dots, e_n^\lambda) w \rangle \\ &= \langle w, U^* U \text{diag}(e_1^\lambda, \dots, e_n^\lambda) w \rangle \\ &= \langle w, \text{diag}(e_1^\lambda, \dots, e_n^\lambda) w \rangle \\ &= w^t \text{diag}(e_1^\lambda, \dots, e_n^\lambda) w \\ &= (w_1 \ \dots \ w_n) \begin{pmatrix} w_1 \\ \vdots \\ w_n \end{pmatrix} \\ &= \sum_{j=1}^n e^{\lambda_j} w_j^2 > 0 \end{aligned}$$

Hence for any self-adjoint X self-adjoint, e^X is self-adjoint and positive.

X is referred to as the "logarithm" of P .

3. We first show that the logarithm X of a self-adjoint positive matrix P depends continuously on P . First, we consider the case where the eigenvalues of P are between 0 and 2. The proof of Theorem 2.7 establishes that if A is diagonalizable and all eigenvalues λ of A satisfy $|\lambda - 1| < 1$, then $\log A$ is defined and $e^{\log A} = A$. Similarly if X is diagonalizable and all eigenvalues λ of X satisfy $|\lambda| < \log 2$, then $\log e^X$ is defined and $\log e^X = X$. So the power series for $\log P$ converges to X .

$$\log P = \sum_{m=1}^{\infty} (-1)^{m+1} \frac{(P - I)^m}{m}$$

is defined for $\|P - I\| < 1$. $(P - I)^m$ is a continuous function of P , so the partial sum is continuous. Note that

$$\left| (-1)^{m+1} \frac{(P - I)^m}{m} \right| = \frac{\|(P - I)^m\|}{m} \leq \frac{\|P - I\|^m}{m} \leq \frac{R^m}{m}, \quad R < 1$$

and that

$$\sum_{m=0}^{\infty} \frac{R^m}{m}$$

converges since

$$\lim_{m \rightarrow \infty} \left| \frac{R^{m+1}}{m+1} \frac{m}{R^m} \right| = \lim_{m \rightarrow \infty} R \frac{m}{m+1} = R < 1$$

By the Weierstrass M-test, the series converges uniformly on all sets of the form $\{\|P - I\| \leq R\}$, for $R < 1$. So $\log P$ is continuous for all $P \in M_n(\mathbb{C})$ such that $\|P - I\| < 1$.

Next, we consider the general case. Fix some self-adjoint positive matrix P_0 . Fix a small neighbourhood $\mathcal{U}$ of P_0 , choose $a > 0$ large. For $P \in \mathcal{U}$, write

$$P = e^a(e^{-a}P) = e^a I(e^{-a}P) = e^{aI}(e^{-a}P)$$

It is easy to see that e^{aI} and $e^{-a}P$ are commutative, and so

$$\log P = \log(e^{aI}(e^{-a}P)) = aI + \log(e^{-a}P)$$

If a is large enough, then for any $P \in \mathcal{U}$, the positive eigenvalues of $e^{-a}P$ all become less than 2, so the series for $\log(e^{-a}P)$ converge and depends continuously on P , thus X depends continuously on P .

By writing

$$P^{1/2} = \exp(\log(P)/2)$$

we see that the square root operation on P is also continuous. The mapping $A \mapsto A^*A$ is continuous, and the mapping $A^*A \mapsto (A^*A)^{1/2}$ is continuous. Since $P = (A^*A)^{1/2}$, P is continuous on A , and so X is continuous on A . Since $U = AP^{-1}$, U is continuous on A .

□

Proposition 2.18. *Polar decomposition for various subgroups of $GL(n, \mathbb{C})$.*

1. Every $A \in GL(n, \mathbb{R})$ can be written uniquely as

$$A = Re^X$$

where $R \in O(n)$ and X is real and symmetric.

2. Every $A \in SL(n, \mathbb{C})$ can be written uniquely as

$$A = Ue^X$$

where $U \in SU(n)$ and X is self-adjoint with trace zero.

3. Every $A \in SL(n, \mathbb{R})$ can be written uniquely as

$$A = Re^X$$

where $R \in SO(n)$ and X is real and symmetric with trace zero.

2.6 Exercises

Exercise 2.6 Show that every 2×2 matrix X with $\text{tr}(X) = 0$ satisfies

$$X^2 = -\det(X)I$$

If X is 2×2 with trace zero, show by direct calculation using the power series for the exponential that

$$e^X = \cos(\sqrt{\det X})I + \frac{\sin \sqrt{\det X}}{\sqrt{\det X}}X$$

where $\sqrt{\det X}$ is either of the two (possibly complex) square roots of $\det X$. Use this to give an alternative computation of the exponential e^{X_1} in Example 2.5.

Proof. Let X be a 2×2 matrix with $\text{tr}(X) = 0$, taking the form

$$X = \begin{pmatrix} a & b \\ c & -a \end{pmatrix}$$

The determinant is

$$\det(X) = -a^2 - bc$$

Hence, we have

$$X^2 = \begin{pmatrix} a & b \\ c & -a \end{pmatrix} \begin{pmatrix} a & b \\ c & -a \end{pmatrix} = \begin{pmatrix} a^2 + bc & 0 \\ 0 & a^2 + bc \end{pmatrix} = (a^2 + bc)I = -\det(X)I$$

which proves the required statement.

The power series of the matrix exponential is

$$e^X = \sum_{n=0}^{\infty} \frac{X^n}{n!} = I + X + \frac{X^2}{2!} + \frac{X^3}{3!} + \frac{X^4}{4!} + \cdots$$

Using the result we just proved, $X^2 = -\det(X)I$, we can generalize and say that

$$X^{2n} = (-1)^n \det(X)^n I$$

$$X^{2n+1} = X^{2n} X = (-1)^n \det(X)^n X$$

The power series of the matrix exponential is absolutely convergent, so we can rearrange terms without worrying about the series converging to some other limit (source).

$$\begin{aligned} e^X &= I + X + \frac{X^2}{2!} + \frac{X^3}{3!} + \frac{X^4}{4!} + \cdots \\ &= \left(I + \frac{X^2}{2!} + \frac{X^4}{4!} + \cdots \right) + \left(X + \frac{X^3}{3!} + \frac{X^5}{5!} + \cdots \right) \\ &= \left(\sum_{n=0}^{\infty} \frac{(-1)^n \det(X)^n}{(2n)!} \right) I + \left(\sum_{n=0}^{\infty} \frac{(-1)^n \det(X)^n}{(2n+1)!} \right) X \\ &= \left(\sum_{n=0}^{\infty} \frac{(-1)^n \sqrt{\det X}^{2n}}{(2n)!} \right) I + \left(\sum_{n=0}^{\infty} \frac{(-1)^n \sqrt{\det X}^{2n+1}}{\sqrt{\det X} (2n+1)!} \right) X \\ &= \cos(\sqrt{\det X}) I + \frac{\sin \sqrt{\det X}}{\sqrt{\det X}} X \end{aligned}$$

where the last equality is from the power series of $\cos$ and $\sin$. This proves the desired identity.

In Example 2.5, the matrix X_1 is

$$X_1 = \begin{pmatrix} 0 & -a \\ a & 0 \end{pmatrix}$$

which is a 2×2 matrix with trace zero. Its determinant is $\det X_1 = a^2$. We apply the above formula to compute its matrix exponential.

$$\begin{aligned} e^{X_1} &= \cos(\sqrt{a^2}) I + \frac{\sin \sqrt{a^2}}{\sqrt{a^2}} X_1 \\ &= \begin{pmatrix} \cos \sqrt{a^2} & 0 \\ 0 & \cos \sqrt{a^2} \end{pmatrix} + \begin{pmatrix} 0 & -a \frac{\sin \sqrt{a^2}}{\sqrt{a^2}} \\ a \frac{\sin \sqrt{a^2}}{\sqrt{a^2}} & 0 \end{pmatrix} \\ &= \begin{pmatrix} \cos a & -\sin a \\ \sin a & \cos a \end{pmatrix} \end{aligned}$$

This is the same result as in Example 2.5. Note that $\sqrt{\det X_1} = \sqrt{a^2}$ can be taken to be either $+a$ or $-a$. The final result is the same. $\square$

3 Lie Algebras

3.1 Definitions and First Examples

Definition 3.1. A *finite-dimensional real or complex Lie algebra* is a finite-dimensional real or complex vector space $\mathfrak{g}$, together with a map $[\cdot, \cdot]$ from $\mathfrak{g} \times \mathfrak{g}$ into $\mathfrak{g}$, with the following properties:

1. $[\cdot, \cdot]$ is bilinear.
2. $[\cdot, \cdot]$ is skew symmetric: $[X, Y] = -[Y, X]$ for all $X, Y \in \mathfrak{g}$.
3. The Jacobi identity holds:

$$[X, [Y, Z]] + [Y, [Z, X]] + [Z, [X, Y]] = 0$$

for all $X, Y, Z \in \mathfrak{g}$.

Two elements X and Y of $\mathfrak{g}$ commute if $[X, Y] = 0$. $\mathfrak{g}$ is commutative if $[X, Y] = 0$ for all $X, Y \in \mathfrak{g}$.

Example 3.2. Let $\mathfrak{g} = \mathbb{R}^3$, let $[\cdot, \cdot] : \mathbb{R}^3 \times \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be given by

$$[x, y] = x \times y$$

where $x \times y$ is the cross product. Then $\mathfrak{g}$ is a Lie algebra.

Example 3.3. Let $\mathcal{A}$ be an associative algebra and let $\mathfrak{g}$ be a subspace of $\mathcal{A}$ such that $XY - YX \in \mathfrak{g}$ for all $X, Y \in \mathfrak{g}$. Then $\mathfrak{g}$ is a Lie algebra with bracket operation given by $[X, Y] = XY - YX$.

Example 3.4. Let $\mathfrak{sl}(n, \mathbb{C})$ denote the space of all $X \in M_n(\mathbb{C})$ for which $\text{tr}(X) = 0$. Then $\mathfrak{sl}(n, \mathbb{C})$ is a Lie algebra with bracket $[X, Y] = XY - YX$.

Definition 3.5. A *subalgebra* of a real or complex Lie algebra $\mathfrak{g}$ is a subspace $\mathfrak{h}$ of $\mathfrak{g}$ such that $[H_1, H_2] \in \mathfrak{h}$ for all $H_1, H_2 \in \mathfrak{h}$. If $\mathfrak{g}$ is a complex Lie algebra and $\mathfrak{h}$ is a real subspace of $\mathfrak{g}$ which is closed under brackets, then $\mathfrak{h}$ is said to be a *real subalgebra* of $\mathfrak{g}$.

A subalgebra $\mathfrak{h}$ of a Lie algebra $\mathfrak{g}$ is said to be an **ideal** in $\mathfrak{g}$ if $[X, H] \in \mathfrak{h}$ for all $X \in \mathfrak{g}$ and $H \in \mathfrak{h}$.

The **center** of a Lie algebra $\mathfrak{g}$ is

$$\mathfrak{z}(\mathfrak{g}) = \{X \in \mathfrak{g} : [X, Y] = 0 \ \forall Y \in \mathfrak{g}\}$$

Example. The vector space $\mathfrak{gl}(n, \mathbb{C})$ of all $n \times n$ complex matrices is a Lie algebra with Lie bracket $[X, Y] = XY - YX$. The subspace $\mathfrak{sl}(n, \mathbb{C})$ of all $n \times n$ complex matrices with trace zero is a Lie subalgebra of $\mathfrak{gl}(n, \mathbb{C})$.

Example. Every Lie algebra $\mathfrak{g}$ has at least two ideals: $\{0\}$ and $\mathfrak{g}$.

$\mathfrak{sl}(n, \mathbb{C})$ is a non-trivial ideal of $\mathfrak{gl}(n, \mathbb{C})$ for $n > 1$.

For a Lie algebra $\mathfrak{g}$, its center $\mathfrak{z}(\mathfrak{g})$ and its derived algebra $\mathcal{D}(\mathfrak{g}) = [\mathfrak{g}, \mathfrak{g}]$ are ideals.

Example. The center of the Heisenberg Lie algebra is

$$\mathfrak{z}(\mathfrak{h}) = \left\{ \begin{pmatrix} 0 & 0 & c \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} : c \in \mathbb{R} \right\} = \{cZ : c \in \mathbb{R}\}$$

Definition 3.6. If $\mathfrak{g}$ and $\mathfrak{h}$ are Lie algebras, then a linear map $\phi : \mathfrak{g} \rightarrow \mathfrak{h}$ is called a **Lie algebra homomorphism** if $\phi([X, Y]) = [\phi(X), \phi(Y)]$ for all $X, Y \in \mathfrak{g}$. If, in addition, ϕ is one-to-one and onto (i.e. bijective), then ϕ is called a **Lie algebra isomorphism**. A Lie algebra isomorphism of a Lie algebra with itself is called a **Lie algebra automorphism**.

Definition 3.7. If $\mathfrak{g}$ is a Lie algebra and X is an element of $\mathfrak{g}$, define a linear map $\text{ad}_X : \mathfrak{g} \rightarrow \mathfrak{g}$ by

$$\text{ad}_X(Y) = [X, Y].$$

The map $X \mapsto \text{ad}_X$ is the **adjoint map** or **adjoint representation**.

We can view ad (the map $X \mapsto \text{ad}_X$) as a linear map of $\mathfrak{g}$ into $\text{End}(\mathfrak{g})$, the space of linear operators on $\mathfrak{g}$.

Proposition 3.8. If $\mathfrak{g}$ is a Lie algebra, then

$$\text{ad}_{[X, Y]} = \text{ad}_X \text{ad}_Y - \text{ad}_Y \text{ad}_X = [\text{ad}_X, \text{ad}_Y];$$

that is, $\text{ad} : \mathfrak{g} \mapsto \text{End}(\mathfrak{g})$ is a Lie algebra homomorphism.

Example. If $\mathfrak{g}$ is a Lie algebra, then the identity map $x \mapsto x$ is a Lie algebra isomorphism, and also a Lie algebra automorphism.

Let K be a field and $\mathfrak{gl}(n, K)$ be the Lie algebra given by all $n \times n$ matrices over K , with Lie bracket given by $[X, Y] = XY - YX$. For $X \in \mathfrak{gl}(K)$ and $g \in GL(n, K)$, the map $X \mapsto gXg^{-1}$ is an isomorphism from $\mathfrak{gl}(n, K)$ to itself, i.e. an automorphism of $\mathfrak{gl}(n, K)$.

Definition 3.9. If $\mathfrak{g}_1$ and $\mathfrak{g}_2$ are Lie algebras, the **direct sum** of $\mathfrak{g}_1$ and $\mathfrak{g}_2$ is the vector space direct sum of $\mathfrak{g}_1$ and $\mathfrak{g}_2$, with bracket given by

$$[(X_1, X_2), (Y_1, Y_2)] = ([X_1, Y_1], [X_2, Y_2]).$$

If $\mathfrak{g}$ is a Lie algebra and $\mathfrak{g}_1$ and $\mathfrak{g}_2$ are subalgebras, we say that $\mathfrak{g}$ **decomposes as the Lie algebra direct sum** of $\mathfrak{g}_1$ and $\mathfrak{g}_2$ if $\mathfrak{g}$ is the direct sum of $\mathfrak{g}_1$ and $\mathfrak{g}_2$ as vector spaces and $[X_1, X_2] = 0$ for all $X_1 \in \mathfrak{g}_1$ and $X_2 \in \mathfrak{g}_2$.

Example. Examples of Lie algebra direct sums

- $\mathfrak{g} = \mathbb{R} \oplus \mathbb{R}$
- $\mathfrak{gl}(2, \mathbb{C}) = \mathfrak{sl}(2, \mathbb{C}) \oplus \mathbb{C}I$

Definition 3.10. Let $\mathfrak{g}$ be a finite-dimensional real or complex Lie algebra, and let $X_1, \dots, X_N$ be a basis for $\mathfrak{g}$ (as a vector space). Then the unique constants c_{jkl} such that

$$[X_j, X_k] = \sum_{l=1}^N c_{jkl} X_l$$

are called the **structure constants** of $\mathfrak{g}$ (with respect to the chosen basis).

The structure constants satisfy the following two conditions:

$$c_{jkl} + c_{kjl} = 0$$

$$\sum_n (c_{jkn} c_{nlm} + c_{kln} c_{njm} + c_{ljn} c_{nkm}) = 0$$

for all j, k, l, m .

Example. A basis for $\mathfrak{sl}(2, \mathbb{C})$ is

$$X_1 = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}; \quad X_2 = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}; \quad X_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

and we have the relations

$$[X_3, X_1] = 2X_1; \quad [X_3, X_2] = -2X_2; \quad [X_1, X_2] = X_3$$

The structure constants are

$$c_{311} = 2, \quad c_{322} = -2, \quad c_{123} = 1$$

with all others following from skew symmetry or equal to 0.

3.2 Simple, Solvable, and Nilpotent Lie Algebras

Definition 3.11. A Lie algebra $\mathfrak{g}$ is called **irreducible** if the only ideals in $\mathfrak{g}$ are $\mathfrak{g}$ and $\{0\}$. A Lie algebra $\mathfrak{g}$ is called **simple** if it is irreducible and $\dim \mathfrak{g} \geq 2$

Example. Irreducible but not simple Lie algebras include the one-dimensional Lie algebras. For instance, take $\mathfrak{g} = \mathbb{R}$ with bracket $[x, y] = 0$.

Proposition 3.12. The Lie algebra $\mathfrak{sl}(2, \mathbb{C})$ is simple.

Definition 3.13. If $\mathfrak{g}$ is a Lie algebra, then the **commutator ideal** in $\mathfrak{g}$, denoted $[\mathfrak{g}, \mathfrak{g}]$, is the space of linear combinations of commutators, that is, the space of elements Z in $\mathfrak{g}$ that can be expressed as

$$Z = c_1[X_1, Y_1] + \cdots + c_m[X_m, Y_m]$$

for some constants c_j and vectors $X_j, Y_j \in \mathfrak{g}$.

Example. For $\mathfrak{g} = \mathfrak{gl}(n, \mathbb{C})$, the commutator ideal is $[\mathfrak{g}, \mathfrak{g}] = \mathfrak{sl}(n, \mathbb{C})$.

Definition 3.14. For any Lie algebra $\mathfrak{g}$, we define a sequence of subalgebras $\mathfrak{g}_0, \mathfrak{g}_1, \mathfrak{g}_2, \dots$ of $\mathfrak{g}$ inductively as follows: $\mathfrak{g}_0 = \mathfrak{g}$, $\mathfrak{g}_1 = [\mathfrak{g}_0, \mathfrak{g}_0]$, $\mathfrak{g}_2 = [\mathfrak{g}_1, \mathfrak{g}_1]$, etc. These subalgebras are called the **derived series** of $\mathfrak{g}$. A Lie algebra $\mathfrak{g}$ is called **solvable** if $\mathfrak{g}_j = \{0\}$ for some j .

It can be shown, using the Jacobi identity and induction on j , that each $\mathfrak{g}_j$ is an ideal in $\mathfrak{g}$.

Definition 3.15. For any Lie algebra $\mathfrak{g}$, we define a sequence of ideals $\mathfrak{g}^j$ in $\mathfrak{g}$ inductively as follows. We set $\mathfrak{g}^0 = \mathfrak{g}$ and then define $\mathfrak{g}^{j+1}$ to be the space of linear combinations of commutators of the form $[X, Y]$ with $X \in \mathfrak{g}$ and $Y \in \mathfrak{g}^j$. These algebras are called the **upper central series** of $\mathfrak{g}$. A Lie algebra $\mathfrak{g}$ is said to be **nilpotent** if $\mathfrak{g}^j = \{0\}$ for some j .

Proposition 3.16. If $\mathfrak{g} \subset M_3(\mathbb{R})$ denotes the space of 3×3 upper triangular matrices with zeros on the diagonal, then $\mathfrak{g}$ satisfies the assumptions of Example 3.3. The Lie algebra $\mathfrak{g}$ is a nilpotent Lie algebra.

Proposition 3.17. If $\mathfrak{g} \subset M_2(\mathbb{C})$ denotes the space of 2×2 matrices of the form

$$\begin{pmatrix} a & b \\ 0 & c \end{pmatrix}$$

with $a, b, c \in \mathbb{C}$, then $\mathfrak{g}$ satisfies the assumptions of Example 3.3. The Lie algebra $\mathfrak{g}$ is solvable but not nilpotent.

I will use the next two examples to develop Proposition 3.16 and 3.17.

Example. Take

$$\mathfrak{g} = \left\{ \begin{pmatrix} 0 & a & b \\ 0 & 0 & c \\ 0 & 0 & 0 \end{pmatrix} : a, b, c \in \mathbb{R} \right\}$$

The upper central series is as follows. Set $\mathfrak{g}^0 = \mathfrak{g}$. Then

$$\mathfrak{g}^1 = [\mathfrak{g}^0, \mathfrak{g}^0] = \left\{ \begin{pmatrix} 0 & 0 & b \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} : b \in \mathbb{R} \right\}$$

$$\mathfrak{g}^2 = \{0\}$$

Hence $\mathfrak{g}$ is nilpotent.

Example. Take

$$\mathfrak{g} = \left\{ \begin{pmatrix} a & b \\ 0 & c \end{pmatrix} : a, b, c \in \mathbb{C} \right\}$$

which is actually a subalgebra of $\mathfrak{gl}(2, \mathbb{C})$. The derived series is as follows. Define $\mathfrak{g}_0 = \mathfrak{g}$, then

$$\mathfrak{g}_1 = [\mathfrak{g}_0, \mathfrak{g}_0] = \left\{ \begin{pmatrix} 0 & b \\ 0 & 0 \end{pmatrix} : b \in \mathbb{C} \right\}$$

$$\mathfrak{g}_2 = [\mathfrak{g}_1, \mathfrak{g}_1] = \{0\}$$

Hence $\mathfrak{g}$ is solvable.

3.3 The Lie Algebra of a Matrix Lie Group

3.4 Examples

3.5 Lie Group and Lie Algebra Homomorphisms

3.6 The Complexification of a Real Lie Algebra

3.7 The Exponential Map

3.8 Consequences of Theorem 3.42

3.9 Exercises

Exercise 3.8 Show that two isomorphic matrix Lie groups have isomorphic Lie algebras.

Proof. Let G and H be two isomorphic matrix Lie groups ($G \cong H$) with corresponding Lie algebras $\mathfrak{g}$ and $\mathfrak{h}$, respectively. Since $G \cong H$, let $\Phi : G \rightarrow H$ be a Lie group isomorphism. By Theorem 3.28, there exists a unique real-linear map $\phi : \mathfrak{g} \rightarrow \mathfrak{h}$ such that

$$\Phi(e^X) = e^{\phi(X)}$$

for all $X \in \mathfrak{g}$. The map ϕ has the properties

1. $\phi(AXA^{-1}) = \Phi(A)\phi(X)\Phi(A)^{-1}$ for all $X \in \mathfrak{g}, A \in G$
2. $\phi([X, Y]) = [\phi(X), \phi(Y)]$, for all $X, Y \in \mathfrak{g}$
3. $\phi(X) = \frac{d}{dt}\Phi(e^{tX})|_{t=0}$ for all $X \in \mathfrak{g}$.

From property 2, it is clear that ϕ is a Lie algebra homomorphism. It remains to show that ϕ is bijective.

Since Φ is a Lie group isomorphism, it must be bijective, hence injective. So $\ker(\Phi) = \{e_G\}$. By Proposition 3.31, we have

$$\ker(\phi) = \text{Lie}(\ker(\Phi)) = \text{Lie}(\{e_G\}) = \{0\}$$

where 0 is the zero element of $\mathfrak{g}$. This implies ϕ is injective. I assume that we are only working with finite-dimensional Lie groups and finite-dimensional Lie algebras. In this case, the dimension of a Lie group equals the dimension of its Lie algebra. Also, since $G \cong H$, $\dim G = \dim H$. So we have

$$\dim \mathfrak{g} = \dim G = \dim H = \dim \mathfrak{h}$$

Since ϕ is injective and $\dim \mathfrak{g} = \dim \mathfrak{h}$, ϕ is bijective.

Hence ϕ is an isomorphism, and the Lie algebras $\mathfrak{g}$ and $\mathfrak{h}$ are isomorphic. $\square$

Remark. Based on the discussion during the groupe de travail, to complete the proof, we still need to show that ϕ^{-1} is a Lie group homomorphism, i.e. it satisfies

$$\phi^{-1}([X, Y]) = [\phi^{-1}(X), \phi^{-1}(Y)]$$

To do this, we can apply ϕ to both sides, and see that both yield $[X, Y]$. Since ϕ is injective, the condition has been proven. However, I am not sure why this extra step is needed.

Exercise 3.13 Let G be a matrix Lie group and let $\mathfrak{g}$ be its Lie algebra. For each $A \in G$, show that Ad_A is a Lie algebra automorphism of $\mathfrak{g}$.

Proof. For each $A \in G$,

$$\begin{aligned} [\text{Ad}_A(X), \text{Ad}_A(Y)] &= \text{Ad}_A(X)\text{Ad}_A(Y) - \text{Ad}_A(Y)\text{Ad}_A(X) \\ &= (AXA^{-1})(AYA^{-1}) - (AYA^{-1})(AXA^{-1}) \\ &= A(XY)A^{-1} - A(YX)A^{-1} \\ &= A(XY - YX)A^{-1} \\ &= A[X, Y]A^{-1} \\ &= \text{Ad}_A([X, Y]) \end{aligned}$$

for all $X, Y \in \mathfrak{g}$. Hence $\text{Ad}_A : \mathfrak{g} \rightarrow \mathfrak{g}$ is a Lie algebra homomorphism. It remains to prove that Ad_A is bijective.

Let $X, Y \in \mathfrak{g}$, with $X \neq Y$. We have $\text{Ad}_A(X) = AXA^{-1}$ and $\text{Ad}_A(Y) = AYA^{-1}$. Suppose $\text{Ad}_A(X) = \text{Ad}_A(Y)$, then $AXA^{-1} = AYA^{-1}$, then by left and right multiplication by A , $X = Y$, which is a contradiction. So Ad_A is an injective mapping.

Let $X \in \mathfrak{g}$. Define $P = A^{-1}XA$. By Theorem 3.20, $AXA^{-1} \in \mathfrak{g}$ for all $A \in G$. Since G is a group, $A^{-1} \in G$, so we see that $P = A^{-1}XA \in \mathfrak{g}$ for each $A \in G$. It is clear that

$$\text{Ad}_A(P) = APA^{-1} = X$$

This means $\forall X \in \mathfrak{g}, \exists P \in \mathfrak{g}$ such that $\text{Ad}_A(P) = X$. So Ad_A is a surjective mapping.

Hence $\text{Ad}_A : \mathfrak{g} \rightarrow \mathfrak{g}$ is a Lie algebra isomorphism, and furthermore a Lie algebra automorphism. $\square$

4 Basic Representation Theory

4.1 Representations

4.2 Examples of Representations

4.3 New Representations from Old

4.4 Complete Reducibility

4.5 Schur's Lemma

4.6 Representations of $\mathfrak{sl}(2; \mathbb{C})$

4.7 Group Versus Lie Algebra Representations

4.8 A Nonmatrix Lie Group

4.9 Exercises