

Gibbs Sampling - (continued)

$$P(x, y) = y \exp(-[xy + y])$$

$$P(x|y) = \frac{P(x,y)}{\int dx' P(x',y)}$$

$$P(x|y) \propto \exp(-xy)$$

$$\exp(-x)$$

(Correction from Wednesday's Lecture.)

A. G. M.

Constitutive
relation

Uniqueness

$$b(x(2)) = g_2(x(2))$$

$$b(x(2)) = \begin{cases} g_2(x(2)) \\ g_1(x(2)) \end{cases}$$

$$b(x(2)) = g_2(x(2))$$

(unique)

(simple) gives , it satisfies DB with $\sigma = P$.

(sweep) gives . does satisfy DB .
does have P as stationary dist.

$$\int P(x) \rho(x', x) dx = \rho(x')$$

if true, P is stationary w.r.t. Markov chain $\rho(x', x)$.

consider case $d=2$.

transition $x \rightarrow y$ happens in two stages $(x^o, x') \xrightarrow{\quad} (y^o, x') \xrightarrow{\quad} (y^o, y')$

using 1-dim
conditional dists.

(x^o, x') (y^o, y')

$$P([y^o, y^i], [x^o, x^i]) = P(y^o | x^i) P(y^i | y^o)$$

$$= \frac{P(x^o, y^i)}{\int da P(a, x^i)} \frac{P(y^o, b)}{\int db P(y^o, b)}$$

sub into start condition.

$$\int dx^o \int dx^i P(x^o, x^i) P([y^o, y^i], [x^o, x^i]) = ? P(y^o, y^i)$$

$$= \int dx^o \int dx^i P(x^o, x^i) \frac{P(y^o, x^i)}{\int da P(a, x^i)} \frac{P(y^o, b)}{\int db P(y^o, b)}$$

$$= \cancel{\int dx^o P(x^o, x^i) \frac{P(y^o, y^i)}{\int db P(y^o, b)}} \int dx^i \frac{P(y^o, x^i)}{\int da P(a, x^i)}$$

$$\begin{aligned} & \int_{\mathbb{R}^d} b(x) \int_{\mathbb{R}^d} b(x, y) \int_{\mathbb{R}^d} b(y, z) \int_{\mathbb{R}^d} b(z, w) \\ & \quad b(w, v) b(v, u) b(u, x) b(x, y) b(y, z) b(z, w) \\ & = b(x, y) b(y, z) b(z, w) b(w, v) b(v, u) b(u, x) \end{aligned}$$

$$\begin{aligned} & \int_{\mathbb{R}^d} b(x) \int_{\mathbb{R}^d} b(x, y) \int_{\mathbb{R}^d} b(y, z) \int_{\mathbb{R}^d} b(z, w) \\ & \quad b(w, v) b(v, u) b(u, x) b(x, y) b(y, z) b(z, w) \\ & = b(x, y) b(y, z) b(z, w) b(w, v) b(v, u) b(u, x) \end{aligned}$$

so we have
the condition.

$$b(2, 1) = b(2, 2)$$

$$\begin{aligned} & \int_{\mathbb{R}^d} b(x) \int_{\mathbb{R}^d} b(x, y) \int_{\mathbb{R}^d} b(y, z) \\ & \quad b(z, w) b(w, v) b(v, u) b(u, x) b(x, y) b(y, z) \end{aligned}$$

$$b([x_2, x_1], [x_3, x_4]) = b([x_2, x_4]) b([x_1, x_3])$$

$$= \frac{P(y^e, y')}{\int dy P(y, s)}$$

$$\left[\int dx' \frac{P(y^e, x')}{\int dx P(x, x')} \right]$$

$$\left(\int dx^e P(x^e, x') \right)$$

□

$$= P(y^e, y')$$

$$= P(y)$$

11

6	2
2	6
2	2
2	2
2	2
2	2

12

6	2
2	6
2	2
2	2
2	2
2	2

13

6	2
2	6
2	2
2	2
2	2
2	2

14

6	2
2	6
2	2
2	2
2	2
2	2

15

6	2
2	6
2	2
2	2
2	2
2	2

$$[x^c, x'] \rightarrow [y^o, x'] \rightarrow [y^o, y']$$

$$P([y^o, y'], [x^o, x']) = P(y^o | x') P(y' | \underline{x^o})$$

This will Not converge to P .

$$\text{This will have a stat dist } Q(x^o, x') = Q^o(x^o) Q^i(x')$$

"ignore" conditions in target dist.

"Polar" consequence in tetrahedral
but

$$(\text{CH}_3)_2\text{Si}^+ \text{Cl}^- = (\text{CH}_3)_2\text{Si}^+(\text{Cl})$$

for SiCl₃ SiCl₂ etc.

$$[\text{CH}_3]^+ \rightarrow [\text{CH}_3]^+ \text{Cl}^- = (\text{CH}_3)^+ \text{Cl}^-$$

$$[\text{CH}_3]^+ \rightarrow [\text{CH}_3]^+ \text{Cl}^- = (\text{CH}_3)^+ \text{Cl}^-$$

$D\beta$

$$P(x) \rho(x, x') = P(x') \rho(x, x')$$

stationarity

$$\int dx \underbrace{P(x)}_{\text{Lap}} \rho(x, x') = P(x')$$

current point new points

$D\beta$ is (usually) easier to check because local.
Stationarity, a global property, is harder.

جامعة

جامعة عجمان

جامعة

جامعة

جامعة

جامعة عجمان

جامعة

جامعة عجمان

جامعة عجمان

جامعة

جامعة عجمان

"Blurred" Gibbs sampler.

$$P(x^0, x^1, \underbrace{x^2, x^3}_{\theta^0} | \theta^1)$$

sample

$$P(\theta^0 | \theta^1)$$
$$P(\theta^1 | \theta^0)$$

either choose which it render to update
or sweep through updating all blocks in order.

Passive knee
stable

1

0.0
0.0

0.0
0.0

0.0
0.0

0.0
0.0

0.0
0.0

0.0
0.0

0.0
0.0

0.0
0.0

0.0
0.0

0.0
0.0

0.0
0.0

0.0
0.0

max

0.0
0.0

0.0
0.0

0.0
0.0

0.0
0.0

0.0
0.0

0.0
0.0

0.0
0.0

0.0
0.0

0.0
0.0

0.0

0.0
0.0

0.0
0.0

0.0
0.0


0.0
0.0

0.0
0.0

0.0
0.0

0.0
0.0

0.0
0.0



...
...

z

...

for someone familiar with MH

- have to choose Q (proposal)
- accept/reject $\alpha \notin \min\left(1, \frac{P(y)}{P(x)} \frac{\alpha(x|y)}{\alpha(y|x)}\right)$

Let $Q = w_k \mathcal{T}^{(d-1)}(x^{-k} - y^{-k}) P(x^k | x^{-k})$

$$\Rightarrow \alpha = 1 \quad \text{always.}$$

for someone familiar with Gibbs

other freedom to use any proposal (not just conditionals)

if you reject some points

HM. After writing was done, Sf

(unfilled) Q was sent by express.
((212) 1000-1) 205 = 205
((212) 1000-1) 205 = 205

(-x) $\frac{d}{dx} \left(e^{ax^2} \right) + b e^{ax^2} = Q$

205 = 205

for
205051902
JULY 19 1962
W.H. HARRIS
RECEIVED
CLERICAL WORKS
LIBRARY

205051902
JULY 19 1962
W.H. HARRIS
RECEIVED
CLERICAL WORKS
LIBRARY