

# Notes

Xinyu Zhong  
Wolfson College

November 16, 2022

# Contents

|          |   |          |
|----------|---|----------|
| <b>1</b> | <b>Spacetime Curvature</b>                      | <b>2</b> |
| 1.1      | Subsection 1 . . . . .                          | 2        |
| <b>2</b> | <b>Gravitation field equations</b>              | <b>2</b> |
| 2.1      | Energy Momentum Tensor . . . . .                | 2        |
| 2.1.1    | Properties of Energy-momentum tensor . . . . .  | 2        |
| 2.2      | Energy-momentum Tensor in Ideal Fluid . . . . . | 2        |
| 2.2.1    | Continuity Equation . . . . .                   | 2        |

## Abstract

Abstract of this course

# 1 Spacetime Curvature

*Definition Riemann Curvature Tensor:*

$$R_{abc}^d = \nabla_a \nabla_b v_c - \nabla_b \nabla_a v_c$$

The reason for doing so is that covariant derivative of dual vector field is not commutative

## 1.1 Subsection 1

# 2 Gravitation field equations

## 2.1 Energy Momentum Tensor

*Definition Energy Momentum Tensor:*

$$T^{\mu\nu}(x) = \rho_0(x) u^\mu(x) u^\nu(x)$$

$T^{00}$ : energy density

$T^{i0}$ : i-th component of 3-momentum density (times c)

$T^{ij}$ : flux of i-component of 3-momentum in j-direction

### 2.1.1 Properties of Energy-momentum tensor

- Always symmetric

## 2.2 Energy-momentum Tensor in Ideal Fluid

Consider Ideal Fluid:

We can find a local inertial frame so that  $T^{i0} = 0$ ; and the spatial components are isotropic:  $T^{ij} \propto \delta^{ij}$ .

Hence, in **instantaneous rest frame**:

$$T^{\mu\nu} = \text{diag}(\rho c^2, p, p, p)$$

where  $\rho c^2$  is the rest frame energy density and  $p$  is isotropic pressure. While in general:

$$T^{\mu\nu} = \left(\rho + \frac{p}{c^2}\right) u^\mu u^\nu - p g^{\mu\nu} \quad (1)$$

Equation 1 is the energy momentum tensor, valid in any coordinate system.

### 2.2.1 Continuity Equation

The energy-momentum tensor satisfies continuity equation:

$$\nabla_\mu T^{\mu\nu} = 0$$