

## SUPPLEMENTARY INFORMATION

Transitions between ground state hyperfine levels of a single  $^{171}\text{Yb}^+$  ion confined in a miniature Paul trap (diameter of 2 mm) are driven with microwave radiation close to 12.64 GHz [1] which is generated by mixing the signal from a fixed frequency source at 6.3 GHz with an rf signal whose frequency, amplitude, and phase are adjustable.

$^{171}\text{Yb}^+$  is produced from its neutral precursor by photoionization using a diode laser operating near 399 nm. Laser light near 369 nm driving resonantly the  $S_{1/2} F = 1 \leftrightarrow P_{1/2} F = 0$  transition in  $\text{Yb}^+$  is supplied by a frequency doubled Ti:Sa laser, and serves for cooling and state selectively detecting the ion. State-selective detection is achieved by collecting scattered fluorescence on the  $(S_{1/2}, F = 1 \leftrightarrow P_{1/2}, F = 0)$  resonance which allows for discriminating population in states  $S_{1/2}, F = 0$  (no fluorescence) and  $S_{1/2}, F = 1$  (resonance fluorescence is detected). Initialization in the state  $S_{1/2} F = 0$  ( $|0\rangle$ ) is done using 369 nm light tuned to the  $S_{1/2} F = 1 \leftrightarrow P_{1/2} F = 1$  transition. A diode laser delivers light near 935 nm and drives the  $D_{3/2} \leftrightarrow [3/2]_{1/2}$  transition to avoid optical pumping into the metastable  $D_{3/2}$  state during the cooling and detection periods.

## SUPPLEMENTARY LEGENDS

The STIRAP sequence carried out when varying the holding time  $T$  (Fig. 2) is characterized by these parameters: pulse separation of  $6/f_\Omega$ , pulse width of  $5/f_\Omega$  and  $\Delta t = \frac{1}{10f_\Omega}$ , where  $f_\Omega = \Omega/(2\pi) = 36.5$  kHz. The microwave frequency on the  $|+1\rangle - |0\rangle$  resonance was 12.6528121 GHz, and on the on the  $|-1\rangle - |0\rangle$  resonance it was 12.6328272 GHz; a static magnetic field  $B = 0.714$  mT defines a quantization axis. Each measurement point consists of 300 repetitions.

The parameters for the data shown in Fig. 3a) are as follows: The microwave frequency on the  $|+1\rangle - |0\rangle$  resonance was 12.6533088 GHz, and on the  $|-1\rangle - |0\rangle$  resonance it was 12.6323327 GHz; the microwave Rabi frequency  $\Omega = 31.8 \times 2\pi$  kHz; the rf frequency driving transitions between dressed states was set to 10.49676 MHz; a static magnetic field  $B = 0.749$  mT defines a quantization axis. Each datapoint is

the average of 50 (up to 40 ms) or 25 repetitions (over 500 ms).

For the Ramsey measurement shown in Fig. 3b),  $f_{\Omega} = \Omega/(2\pi) = 37.3$  kHz. The microwave frequency on the  $|+1\rangle - |0\rangle$  resonance was 12.6530938 GHz, and on the  $| -1\rangle - |0\rangle$  resonance it was 12.6325472 GHz; a static magnetic field  $B = 0.730$  mT defines a quantization axis. 0 ms - 30 ms: 20 repetitions per datapoint; 500 ms - 1000 ms: 40 repetitions.

- 
- [1] Hannemann, T. *et al.*, Self-learning estimation of quantum states. *Phys. Rev. A* **65**, 050303 1-4 (2002).