




Full-RIC joint bias: conditional variance and mean residuals




Whole covariance-shape resampling tails: 2 modes 0.17; 5 modes 0.76 (includes cross-mode correlations).
 C_single/25 is used only for mean-noise calibration. Linear contrasts are not posterior-tension significances.

FKP-only paired experiment: independent minus own n(z)

Joint relative to P-only







Joint relative to xi-only



Same data/random positions, per-branch I2 and Poisson subtraction, recomputed P windows and full IC.
Open dots show common-random block sensitivity. Pilot SEM is not an independent 25-phase uncertainty.

Mean paired increments in the four fitted components



The displayed diagonal sigma is a plotting scale; all response calculations use the full 74x74 covariance.
The unchanged 9.18 selection includes the BAO mask. These are paired increments, not new data or constraints.

Numerical checks and interpretation limits

Phase	LS blocks	I2 independent / own	IC scramble difference: q	Local joint b1 shift
ph000	2	0.999499	0.00011	+0.00088
ph012	2	0.996717	0.000196	+0.00027
ph024	2	0.999890	0.00035	-0.00280

Validated: same positions and frozen bins; analytic Poisson subtraction; independent differential IC kernels.

Mode construction: fixed fiducial derivatives and EZ covariance; no halo-residual-based selection.

Covariance: all 1000 original paired EZ vectors retained; no 74-D halo sample-covariance inversion.

Mean-noise interpretation assumes independent halo phases; finite mock covariance uncertainty remains.

FKP reference uses the other 24 phases. Shared reference tables make phase differences correlated.

The pilot uses 4N randoms for P and the listed common N-sized LS blocks, not production 25-block xi.

Model increments are evaluated at one fiducial. Constant-increment MAP is diagnostic only.

No new GSM parameter, RIC amplitude, formal covariance, or posterior is introduced.