STO+Marisa RSD | post-P0 recalibration

Physical p: old input-P0 1.308086 -> new post-P0 1.307926

p 68% interval: [0.6431, 2.0391] | 100 positive post-P0 realizations | calibration fNL fixed at +100

V = 1 (Gpc/h)^3 | mean100 data | unaveraged C_single500 | kP = 0.08 | no IR

Statistic	True fNL	Pre (unchanged)	Post: input-P0 p	Post: post-P0 p
P0 only	+0	0.01 / 95.54	13.83 / 88.44	11.33 / 88.36
P0 only	+100	101.78 / 96.37	110.06 / 88.70	110.20 / 87.95
P0 only	-100	-102.11 / 97.20	-86.30 / 88.66	-86.94 / 89.21
P+B kB=0.08	+0	-5.25 / 94.38	9.18 / 85.75	9.50 / 85.19
P+B kB=0.08	+100	97.23 / 97.64	102.08 / 86.65	100.65 / 87.47
P+B kB=0.08	-100	-104.74 / 95.96	-81.89 / 86.55	-83.04 / 87.68
P+B kB=0.14	+0	1.87 / 85.76	35.75 / 72.77	35.64 / 72.23
P+B kB=0.14	+100	105.20 / 84.62	114.38 / 73.40	115.22 / 73.15
P+B kB=0.14	-100	-97.20 / 88.78	-38.87 / 74.71	-37.39 / 73.98

Table entries: fNL posterior median / posterior standard deviation (not symmetric 68% intervals).

Only 4+4 P+B results are shown; P0-only has its original three active parameters.

Main post targets, full covariance, priors, reconstruction settings and binning are unchanged.

Post covariance-target and calibration-target overlaps are retained and explicitly accepted.

p uncertainty is not propagated. Convergence does not certify full STO noise closure or unbiased recovery.

Pre retains its own p and original covariance IDs; it is a historical control, not a pure reconstruction-only test.

Conditional fNL constraints after post-P0 calibration

Dots: posterior medians. Error bars: central 68% credible intervals. Dashed lines: simulation truth.



Same post data and C_single; only the calibration field and resulting fixed p changed. 4+4 P+B only.