STO+Marisa RSD | real-space p sensitivity

Fixed physical p: post-RSD calibration 1.307926 -> real-space calibrated value 1.282200

V = 1 (Gpc/h)^3 | mean100 data | unaveraged C_single500 | full P-B cross | kP = 0.08 | no IR

Statistic	True fNL	p = 1.307926	p = 1.282200	Median change
P0 only	+0	11.33 / 88.36	14.34 / 86.15	+3.01
P0 only	+100	110.20 / 87.95	106.89 / 85.25	-3.30
P0 only	-100	-86.94 / 89.21	-84.22 / 86.94	+2.72
P+B kB=0.08	+0	9.50 / 85.19	8.88 / 83.62	-0.61
P+B kB=0.08	+100	100.65 / 87.47	101.39 / 85.84	+0.74
P+B kB=0.08	-100	-83.04 / 87.68	-81.37 / 83.53	+1.67
P+B kB=0.14	+0	35.64 / 72.23	35.33 / 70.68	-0.31
P+B kB=0.14	+100	115.22 / 73.15	113.78 / 71.37	-1.45
P+B kB=0.14	-100	-37.39 / 73.98	-38.98 / 72.41	-1.59

Entries: fNL posterior median / posterior standard deviation; central 68% intervals are plotted on page 2.

Only fixed physical p changes; nine main cases use the previous RSD data, covariance, priors and binning.

P0-only keeps its three active parameters; P+B keeps the same four physical plus four noise parameters.

Real-space p is imported from the value actually used in real-space p-aware inference.

p uncertainty is not propagated. Accepted covariance-target overlap is retained.

This is a sensitivity comparison; sampler convergence does not certify unbiased recovery or complete STO noise closure.

RSD fNL constraints with real-space calibrated p

Dots: posterior medians. Bars: central 68% credible intervals. Dashed lines: simulation truth.



Only fixed physical p changed; all remaining RSD inference settings are the same.