

Investigating the molecular cause of hypermobile Ehlers-Danlos syndrome

MSB 1014 - Network Biology

Cay Rahn - 6255648

October 19, 2023

1 Introduction

1.1 Background

- Ehlers-Danlos syndromes are a group of heritable connective tissue disorders that can be classified into multiple subtypes
- current classification describes 13 subtypes [1]
- most common form: hypermobile EDS (hEDS) - only subtype with unknown molecular cause
- diagnosis based on clinical presentation
- identify molecular cause for: diagnosis, understanding, potential treatment [2]
- current state: ongoing study with aim to do this by analysing genes of 1000 affected individuals, results not expected before 2025 [3]
- many studies investigated genes but no clear molecular cause with connection to connective tissue established yet [4]
- until then use data known
- problem with existing data: criteria change in 2017, earlier diagnosed people might not classify anymore [5] + there is a unclear overlap with Hypermobility Spectrum Disorder (HSD) and terms often used interchangeable and diagnoses often grouped together (unclear whether there is a difference) [5, 6]
- we know the clinical representation and how other eds types work (kind of)

1.2 Aim

- aim: investigate molecular cause of hypermobile EDS
- studying differentially expressed genes in hEDS patients
- which biological processes and pathways are affected
- similarities to pathways/processes/function affected by other EDS genes
- TODO: what do we want to do in the big picture: find candidate genes? is this too ambitious?

2 Methods

2.1 Analysis of Differentially Expressed Genes and Network Creation

- data accessible at NCBI GEO database with the accession number GSE218012 [2]
- analysis with DeSeq2 in R based on the analysis exported from GEO2R [7] to identify up-regulated and down-regulated genes
- $|\log_2\text{FoldChange}| > 0.5$, $p\text{Value} < 0.05$, $p\text{Value}$ adjusted with Benjamini and Hochberg False Discovery Rate
- query differentially expressed genes from string db [8] (confidence cut off 0.4)
- query eds genes related to other eds types additionally: 21 genes retrieved from Disease Ontology with Disease Ontology ID 13359 [9], queried from string
- load data about differential expression, for EDS genes and differentially expressed genes

2.2 Enrichment analysis and clustering

- exploratory enrichment analysis on whole network
- large network, thus cluster before to get better insights for specific parts
- two different cluster methods used with different resulting cluster structure

2.2.1 MCODE

- MCODE finds "densely connected regions in large protein-protein interaction networks that may represent molecular complexes" [10]
- results in smaller clusters
- suited to analyse molecular function
- do we see clusters with other molecular functions than expected
- are genes clustered together with other eds genes
- are clusters generally mostly upregulated or downregulated

2.2.2 Community Clustering

- small cluster not helpful for biological processes and pathways
- therefore use second clustering method
- community clustering with GLay, "more suitable for functional interpretation" [11]

3 Results

3.1 Differentially Expressed Genes and Network creation

- 908 differentially expressed genes with chosen thresholds section 2.1
- String was able to query 828 of them [TODO: provide table of not queried in supplementary material or at least github?]
- after querying additional genes that are known to be related to other eds types: resulting network with 847 nodes and 6129 edges

4 Discussion

References

- [1] Malfait F, Francomano C, Byers P, Belmont J, Berglund B, Black J, et al. The 2017 international classification of the Ehlers–Danlos syndromes. *American Journal of Medical Genetics Part C: Seminars in Medical Genetics*. 2017 3;175:8-26.
- [2] Ritelli M, Colombi M. Molecular Genetics and Pathogenesis of Ehlers-Danlos Syndrome and Related Connective Tissue Disorders. *Genes*. 2020 5;11.
- [3] HEDGE (Hypermobile Ehlers-Danlos Genetic Evaluation) Study - The Ehlers Danlos Society;. Available from: <https://www.ehlers-danlos.com/hedge/>.
- [4] Caliozna L, Guerrieri V, Annunziata S, Bina V, Brancato AM, Castelli A, et al. Biomarkers for Ehlers-Danlos Syndromes: There Is a Role? *International Journal of Molecular Sciences* 2021, Vol 22, Page 10149. 2021 9;22:10149.
- [5] Gensemer C, Burks R, Kautz S, Judge DP, Lavallee M, Norris RA. Hypermobile Ehlers-Danlos syndromes: Complex phenotypes, challenging diagnoses, and poorly understood causes. *Developmental dynamics : an official publication of the American Association of Anatomists*. 2021 3;250:318.
- [6] Ritelli M, Chiarelli N, Cinquina V, Zoppi N, Bertini V, Venturini M, et al. RNA-Seq of Dermal Fibroblasts from Patients with Hypermobile Ehlers-Danlos Syndrome and Hypermobility Spectrum Disorders Supports Their Categorization as a Single Entity with Involvement of Extracellular Matrix Degrading and Proinflammatory Pathomechanisms. *Cells*. 2022 12;11.
- [7] Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology*. 2014 12;15:1-21.
- [8] Szklarczyk D, Kirsch R, Koutrouli M, Nastou K, Mehryary F, Hachilif R, et al. The STRING database in 2023: protein-protein association networks and functional enrichment analyses for any sequenced genome of interest. *Nucleic acids research*. 2023 1;51:D638-46.
- [9] Schriml LM, Mitraka E, Munro J, Tauber B, Schor M, Nickle L, et al. Human Disease Ontology 2018 update: Classification, content and workflow expansion. *Nucleic Acids Research*. 2019 1;47:D955-62.
- [10] Bader GD, Hogue CWV. An automated method for finding molecular complexes in large protein interaction networks. *BMC Bioinformatics*. 2003 1;4:1-27.
- [11] Su G, Kuchinsky A, Morris JH, States DJ, Meng F. GLayer: community structure analysis of biological networks. *Bioinformatics*. 2010 12;26:3135-7.