

# Element of Statistical Learning Note

Zichong Wang

December 30, 2025

## Contents

<b>1</b>	<b>Chap 3: Linear Methods for Regression</b>	<b>2</b>
1.1	Confidence region for $\hat{\beta}$	2
1.2	What is Linear	2
1.3	Gauss-Markov Theorem	2
1.4	QR decomposition	3
1.4.1	Application to Least Squared	3
1.4.2	About orthogonal matrix	4
1.5	Multiple testing in forward selection	5
1.6	Ridge regression	5
1.6.1	Equivalence of two forms	6
1.7	Bayesian view	7

# 1 Chap 3: Linear Methods for Regression

## 1.1 Confidence region for $\hat{\beta}$

We know that  $\hat{\beta} \sim \mathcal{N}(\beta, \sigma^2(\mathbf{X}^T \mathbf{X})^{-1})$ , thus  $\hat{\beta} - \beta \sim \mathcal{N}(0, \sigma^2(\mathbf{X}^T \mathbf{X})^{-1})$ .

Since for  $\mathbf{z} \sim \mathcal{N}(0, \mathbf{\Sigma})$ , we have  $\mathbf{z}^T \mathbf{\Sigma}^{-1} \mathbf{z} \sim \chi_k^2$ , where  $k = \text{rank}(\mathbf{\Sigma})$ , we have

$$(\hat{\beta} - \beta)^T (\sigma^2(\mathbf{X}^T \mathbf{X})^{-1})^{-1} (\hat{\beta} - \beta) \sim \chi_{p+1}^2$$

Thus,

$$\frac{1}{\sigma^2} (\hat{\beta} - \beta)^T \mathbf{X}^T \mathbf{X} (\hat{\beta} - \beta) \sim \chi_{p+1}^2$$

And have approx confidence region

$$(\hat{\beta} - \beta)^T \mathbf{X}^T \mathbf{X} (\hat{\beta} - \beta) \leq \hat{\sigma}^2 \chi_{p+1, 1-\alpha}^2$$

Actually,  $\hat{\sigma}^2 = \frac{1}{N-p-1} \sum_{i=1}^N (y_i - \hat{y}_i)^2 = \frac{1}{N-p-1} \text{RSS}$ , and  $\text{RSS}/\sigma^2 \sim \chi_{N-p-1}^2$ ,

$$\frac{(\hat{\beta} - \beta)^T \mathbf{X}^T \mathbf{X} (\hat{\beta} - \beta)/(p+1)}{\hat{\sigma}^2} \sim F_{p+1, N-p-1}$$

## 1.2 What is Linear

In the context of **linear model**, we are talking about linearity in parameters, meaning that the prediction  $\hat{y}$  is a linear combination of the parameters  $\beta_j$ . The  $\mathbf{X}$  itself can be non-linear transformations of the original features, e.g., polynomial terms, interaction terms, etc.  $y = 1/(\beta_0 + \beta_1 x)$  and  $y = \beta_0 e^{\beta_1 x}$  are not linear models, since they're not linear in parameters.

In the context of **Linear estimators**, we are talking about the estimator  $\hat{\theta}$  (e.g.  $\hat{\beta}$ ) can be written as a linear combination of the observed response values  $y_i$ , i.e.  $\hat{\theta} = \mathbf{c}^T \mathbf{y}$ . The weight  $\mathbf{c}$  depends only on  $\mathbf{X}$ , not on  $\mathbf{y}$ . A linear estimator **can be** a prediction at a new point, or the estimated coefficients  $\hat{\beta}$  themselves.

## 1.3 Gauss-Markov Theorem

Why assume only know  $\mathbf{X}$ , but not  $\mathbf{y}$ ?

Note that though  $y_i$  as sample responses, are observable, the following statements and arguments including assumptions, proofs and the others assume under the only condition of knowing  $\mathbf{X}_{i,j}$  but not  $y_i$ . — [1]

We have a *challenger* linear estimator  $\tilde{\beta} = \mathbf{C} \mathbf{y}$ , where  $\mathbf{C} = (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T + \mathbf{D}$ , a modification

of OLS estimator. Ensure it's unbiased:

$$\begin{aligned}
\mathbb{E}(\tilde{\beta}) &= \mathbb{E}(\mathbf{C}\mathbf{y}) \\
&= \mathbf{C}\mathbb{E}(\mathbf{y}) = ((\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T + \mathbf{D})\mathbf{X}\beta \\
&= \beta + \mathbf{D}\mathbf{X}\beta \\
&= \beta
\end{aligned}$$

Meaning  $\mathbf{D}\mathbf{X} = 0$ .

Now, compute the variance:

$$\begin{aligned}
\text{Var}(\tilde{\beta}) &= \text{Var}(\mathbf{C}\mathbf{y}) \\
&= \mathbf{C}\text{Var}(\mathbf{y})\mathbf{C}^T \\
&= \sigma^2 \mathbf{C}\mathbf{C}^T \\
&= \sigma^2 ((\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T + \mathbf{D})((\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T + \mathbf{D})^T \\
&= \sigma^2 ((\mathbf{X}^T \mathbf{X})^{-1} + \mathbf{D}\mathbf{D}^T) \\
&= \text{Var}(\hat{\beta}) + \sigma^2 \mathbf{D}\mathbf{D}^T \\
&\geq \text{Var}(\hat{\beta})
\end{aligned}$$

## 1.4 QR decomposition

Any real squared matrix  $\mathbf{A}$  can be decomposed as  $\mathbf{A} = \mathbf{Q}\mathbf{R}$ , where  $\mathbf{Q}$  is orthogonal matrix ( $\mathbf{Q}^T \mathbf{Q} = \mathbf{I}$ ), and  $\mathbf{R}$  is upper triangular matrix.

Any rectangular matrix  $\mathbf{A} \in \mathbb{R}^{m \times n}$  ( $m \geq n$ ), we can decompose it as  $\mathbf{A} = \mathbf{Q}\mathbf{R}$ , where  $\mathbf{Q}$  is  $m \times m$  orthogonal matrix, and  $\mathbf{R}$  is  $m \times n$  upper triangular matrix (the last  $m - n$  rows are all zero). It can be regarded as  $\mathbf{A} = \mathbf{Q}\mathbf{R} = [\mathbf{Q}_1 \quad \mathbf{Q}_2] \begin{bmatrix} \mathbf{R}_1 \\ \mathbf{0} \end{bmatrix} = \mathbf{Q}_1 \mathbf{R}_1$ , where  $\mathbf{Q}_1 \in \mathbb{R}^{m \times n}$ ,  $\mathbf{Q}_2 \in \mathbb{R}^{m \times (m-n)}$ ,  $\mathbf{R}_1 \in \mathbb{R}^{n \times n}$  which is upper triangular.

If  $\mathbf{A}$  have  $k$  linearly independent columns, then first  $k$  columns of  $\mathbf{Q}$  form an orthonormal basis of the column space of  $\mathbf{A}$ . The fact that any column  $k$  of  $\mathbf{A}$  only depends on the first  $k$  columns of  $\mathbf{Q}$  corresponds to the triangular form of  $\mathbf{R}$ .

QR decomposition can be calculated using Gram-Schmidt process, or using Householder reflections. In practice, Householder reflections are more stable and efficient.

### 1.4.1 Application to Least Squared

In linear least squares problems, we aim to find a vector  $\mathbf{x}$  that minimizes the Euclidean norm of the residual for an overdetermined system  $\mathbf{A}\mathbf{x} \approx \mathbf{b}$ , where  $\mathbf{A} \in \mathbb{R}^{m \times n}$  and  $m \geq n$ . The goal is to solve:

$$\min_{\mathbf{x}} \|\mathbf{A}\mathbf{x} - \mathbf{b}\|_2$$

Using the Full QR Decomposition, we substitute  $\mathbf{A} = \mathbf{Q} \begin{bmatrix} \mathbf{R}_1 \\ \mathbf{0} \end{bmatrix}$ :

$$\|\mathbf{A}\mathbf{x} - \mathbf{b}\|_2 = \left\| \mathbf{Q} \begin{bmatrix} \mathbf{R}_1 \\ \mathbf{0} \end{bmatrix} \mathbf{x} - \mathbf{b} \right\|_2$$

Since multiplying by an orthogonal matrix  $Q$  preserves the Euclidean norm, we can left-multiply the entire expression by  $Q^T$ :

$$\|Ax - b\|_2 = \left\| \begin{bmatrix} R_1 \\ 0 \end{bmatrix} x - Q^T b \right\|_2$$

If we partition  $Q^T b$  into two components— $c_1 \in \mathbb{R}^n$  and  $c_2 \in \mathbb{R}^{m-n}$ :

$$\left\| \begin{bmatrix} R_1 x - c_1 \\ -c_2 \end{bmatrix} \right\|_2^2 = \|R_1 x - c_1\|_2^2 + \|c_2\|_2^2$$

To minimize the total error, we must make the first term zero. The least squares solution is found by solving the square, upper-triangular system:

$$R_1 x = c_1$$

Since  $R_1$  is upper triangular, we can efficiently solve this system using back substitution, **that's how we solve linear equations manually in algebra class!** The remaining term  $\|c_2\|_2$  represents the minimum residual norm (the "error" of the fit).

#### 1.4.2 About orthogonal matrix

Let  $Q \in \mathbb{R}^{n \times n}$  be an orthogonal matrix. Write  $Q$  in terms of its column vectors:

$$Q = [q_1 \ q_2 \ \cdots \ q_n], \quad q_i^\top q_j = \delta_{ij}.$$

Then

$$Q^\top = \begin{bmatrix} q_1^\top \\ q_2^\top \\ \vdots \\ q_n^\top \end{bmatrix}.$$

Computation of  $Q^\top Q$ .

$$Q^\top Q = \begin{bmatrix} q_1^\top \\ q_2^\top \\ \vdots \\ q_n^\top \end{bmatrix} [q_1 \ q_2 \ \cdots \ q_n] = \begin{bmatrix} q_1^\top q_1 & q_1^\top q_2 & \cdots & q_1^\top q_n \\ q_2^\top q_1 & q_2^\top q_2 & \cdots & q_2^\top q_n \\ \vdots & \vdots & \ddots & \vdots \\ q_n^\top q_1 & q_n^\top q_2 & \cdots & q_n^\top q_n \end{bmatrix}.$$

Using  $q_i^\top q_j = \delta_{ij}$ ,

$$Q^\top Q = \begin{bmatrix} 1 & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 \end{bmatrix} = I.$$

Computation of  $QQ^\top$ .

$$QQ^\top = [q_1 \ q_2 \ \cdots \ q_n] \begin{bmatrix} q_1^\top \\ q_2^\top \\ \vdots \\ q_n^\top \end{bmatrix} = q_1 q_1^\top + q_2 q_2^\top + \cdots + q_n q_n^\top = \sum_{k=1}^n q_k q_k^\top.$$

For any  $\mathbf{x} \in \mathbb{R}^n$ ,

$$(\mathbf{Q}\mathbf{Q}^\top)\mathbf{x} = \sum_{k=1}^n \mathbf{q}_k(\mathbf{q}_k^\top \mathbf{x}) = \mathbf{x},$$

since  $\{\mathbf{q}_1, \dots, \mathbf{q}_n\}$  is an orthonormal basis of  $\mathbb{R}^n$ . Hence

$$\mathbf{Q}\mathbf{Q}^\top = \mathbf{I}.$$

Therefore, for an orthogonal matrix  $\mathbf{Q}$ ,

$$\mathbf{Q}^\top \mathbf{Q} = \mathbf{Q}\mathbf{Q}^\top = \mathbf{I}.$$

## 1.5 Multiple testing in forward selection

ESL page 60:

Other more traditional packages base the selection on F -statistics, adding “significant” terms, and dropping “non-significant” terms. These are out of fashion, since they do not take proper account of the multiple testing issues.

Assume we have  $p$  candidate features, and already selected  $k$  features. When considering adding a new feature, we are actually performing  $p - k$  hypothesis tests (each test for one feature). Even the rest  $p - k$  features are all noise, with significance level  $\alpha$ , we still have a probability of  $1 - (1 - \alpha)^{p-k}$  to incorrectly add at least one noise feature.

## 1.6 Ridge regression

Answer questions: why two forms are equivalent? Why not equivariant under scaling of the inputs? What is a good practice for it? df of ridge? In the case of orthogonal inputs, why  $\hat{\beta}^{\text{ridge}} = \hat{\beta}/(1 + \lambda)$ ?

Ridge regression shrinks the regression coefficients by imposing a penalty on their size:

$$\hat{\beta}^{\text{ridge}} = \arg \min_{\beta} \left\{ \sum_{i=1}^N (y_i - \beta_0 - \sum_{j=1}^p x_{ij} \beta_j)^2 + \lambda \sum_{j=1}^p \beta_j^2 \right\}$$

and is equivalent to

$$\begin{aligned} \hat{\beta}^{\text{ridge}} &= \arg \min_{\beta} \sum_{i=1}^N (y_i - \beta_0 - \sum_{j=1}^p x_{ij} \beta_j)^2 \\ &\text{subject to } \sum_{j=1}^p \beta_j^2 \leq t \end{aligned}$$

And there is a one-to-one correspondence between  $\lambda$  and  $t$ .

### 1.6.1 Equivalence of two forms

First, take a review of **KKT conditions**.

Consider a minimization problem with both equality and inequality constraints:

$$\begin{aligned} & \min_{\mathbf{x}} f(\mathbf{x}) \\ & \text{subject to } g_i(\mathbf{x}) \leq 0, & i = 1, \dots, m \\ & h_j(\mathbf{x}) = 0, & j = 1, \dots, l \end{aligned}$$

And the lagrangian function is:

$$L(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\mu}) = f(\mathbf{x}) + \sum_{i=1}^m \lambda_i g_i(\mathbf{x}) + \sum_{j=1}^l \mu_j h_j(\mathbf{x})$$

If  $\mathbf{x}^*$  is a local minimum, then there exist multipliers  $\lambda_i^* \geq 0$  and  $\mu_j^*$  such that the following conditions hold:

- **Stationary**

$$\nabla_{\mathbf{x}} L(\mathbf{x}^*, \boldsymbol{\lambda}^*, \boldsymbol{\mu}^*) = 0$$

- **Primal feasibility**

$$g_i(\mathbf{x}^*) \leq 0, \quad i = 1, \dots, m$$

$$h_j(\mathbf{x}^*) = 0, \quad j = 1, \dots, l$$

The gradient can be regarded as the force to push a particle, primal stationary means the force of  $\partial f(\mathbf{x}^*)$  is balanced by a linear sum of forces from constraints.

- **Dual feasibility**

$$\lambda_i^* \geq 0, \quad i = 1, \dots, m$$

All the  $\partial g_i(\mathbf{x}^*)$  forces must be one-sided, pointing inwards into the feasible set for  $\mathbf{x}$ .

- **Complementary slackness**

$$\lambda_i^* g_i(\mathbf{x}^*) = 0, \quad i = 1, \dots, m$$

The force only activated when the particle is on the boundary of feasible set.

In ridge regression, we have:

$$\mathcal{L} = \|\mathbf{Y} - \mathbf{X}\beta\|_2^2 + \alpha(\|\beta\|_2^2 - t)$$

According to stationary condition:

$$\nabla_{\beta} \mathcal{L} = -2\mathbf{X}^T(\mathbf{Y} - \mathbf{X}\beta) + 2\alpha\beta = 0$$

On the other hand, solving the unconstrained form is

$$\nabla_{\beta} (\|\mathbf{Y} - \mathbf{X}\beta\|_2^2 + \lambda\|\beta\|_2^2) = -2\mathbf{X}^T(\mathbf{Y} - \mathbf{X}\beta) + 2\lambda\beta = 0$$

Thus, if we set  $\lambda = \alpha$ , the two forms are equivalent.

One step further, according to complementary slackness:

$$\alpha(\|\beta\|_2^2 - t) = 0$$

**From unconstrained form, given  $\lambda$ ,** we can solve  $\beta$ , denote  $\beta(\lambda)$ , and define  $t(\lambda) = \|\beta(\lambda)\|_2^2$ . Then,  $\beta(\lambda)$  is also the solution of constrained form with  $t = t(\lambda)$  (apparently it's on the boundary, and the coefficient  $\alpha = \lambda$ ).

**From the constrained form, given  $t$ ,** we can also solve  $\beta$ , denote  $\beta(t)$ .

- If  $\|\beta(t)\|_2^2 < t$ , then according to complementary slackness,  $\alpha = 0$ , which means no penalty,  $\lambda = 0$ , back to OLS.
- If  $\|\beta(t)\|_2^2 = t$ , the boundary is effective, correspond to some  $\alpha > 0$ , and  $\lambda = \alpha$ .

## 1.7 Bayesian view

## References

- [1] Wikipedia contributors, "Gauss–Markov theorem," *Wikipedia, The Free Encyclopedia*, [https://en.wikipedia.org/wiki/Gauss%E2%80%93Markov\\_theorem](https://en.wikipedia.org/wiki/Gauss%E2%80%93Markov_theorem) (accessed Dec 29, 2025).