

Newton's Method and its Ramifications

Zichuan Wang

Ohio State University

April 6, 2021

History

We begin with a story.

Historical Part

Application I

Example I

References

-

We begin with a story.



Leibniz (1646 - 1716)



Oldenburg (1619 - 1677)

On May 2, 1676, Gottfried Wilhelm Leibniz wrote to Henry Oldenburg, the secretary of the Royal Society of London,

Historical Part

Application I

Example I

References

-

We begin with a story.



Leibniz (1646 - 1716)



Oldenburg (1619 - 1677)

On May 2, 1676, Gottfried Wilhelm Leibniz wrote to Henry Oldenburg, the secretary of the Royal Society of London, asking the English mathematician John Collins for a proof of

$$\sin(x) = x - \frac{1}{6}x^3 + \frac{1}{120}x^5 - \frac{1}{5040}x^7 + \frac{1}{362880}x^9 + \dots$$

We begin with a story.



Leibniz (1646 - 1716)



Oldenburg (1619 - 1677)

On May 2, 1676, Gottfried Wilhelm Leibniz wrote to Henry Oldenburg, the secretary of the Royal Society of London, asking the English mathematician John Collins for a proof of

$$\sin(x) = x - \frac{1}{6}x^3 + \frac{1}{120}x^5 - \frac{1}{5040}x^7 + \frac{1}{362880}x^9 + \dots$$

and information about how English mathematicians made use of infinite series in their research.

Sir Isaac Newton (1642 - 1727) answered Leibniz's request by writing a letter, now known as "Epistola Prior", to Oldenburg on June 13, 1676, in which Newton described his method of infinite series giving many examples thereof.

Sir Isaac Newton (1642 - 1727) answered Leibniz's request by writing a letter, now known as "Epistola Prior", to Oldenburg on June 13, 1676, in which Newton described his method of infinite series giving many examples thereof. In particular, he applied his generalized binomial theorem:

$$(x + y)^r = \sum_{k=0}^{\infty} \binom{r}{k} x^{r-k} y^k, \quad \forall r \in \mathbb{R}.$$

Sir Isaac Newton (1642 - 1727) answered Leibniz's request by writing a letter, now known as "Epistola Prior", to Oldenburg on June 13, 1676, in which Newton described his method of infinite series giving many examples thereof. In particular, he applied his generalized binomial theorem:

$$(x + y)^r = \sum_{k=0}^{\infty} \binom{r}{k} x^{r-k} y^k, \quad \forall r \in \mathbb{R}.$$



Leibniz



Oldenburg



Newton

Letter 1

We discuss portions of Epistola Prior and the next one (the “Epistola Posterior”).

Reading Classics
Math Seminar

Zichuan Wang

Historical Part

Application I

Example I

References

-

Letter 1

We discuss portions of Epistola Prior and the next one (the “Epistola Posterior”). In Epistola Prior, Newton approximates the real solutions of $f(y) = y^3 - 2y - 5 = 0$ to illustrate his method.

Letter 1

We discuss portions of Epistola Prior and the next one (the “Epistola Posterior”). In Epistola Prior, Newton approximates the real solutions of $f(y) = y^3 - 2y - 5 = 0$ to illustrate his method.

He guesses that a root is near $y = 2$ since, for example, $f(1) < 0$ but $f(3) > 0$.

Letter 1

We discuss portions of Epistola Prior and the next one (the “Epistola Posterior”). In Epistola Prior, Newton approximates the real solutions of $f(y) = y^3 - 2y - 5 = 0$ to illustrate his method.

He guesses that a root is near $y = 2$ since, for example, $f(1) < 0$ but $f(3) > 0$.

A root equals $y = 2 + p$ for some small p (at least, $|p| < 1$).

Letter 1

We discuss portions of Epistola Prior and the next one (the “Epistola Posterior”). In Epistola Prior, Newton approximates the real solutions of $f(y) = y^3 - 2y - 5 = 0$ to illustrate his method.

He guesses that a root is near $y = 2$ since, for example, $f(1) < 0$ but $f(3) > 0$.

A root equals $y = 2 + p$ for some small p (at least, $|p| < 1$). Substituting $y = 2 + p$ into $f(y) = 0$, he gets $g(p) = p^3 + 6p^2 + 10p - 1 = 0$.

Letter 1

We discuss portions of Epistola Prior and the next one (the “Epistola Posterior”). In Epistola Prior, Newton approximates the real solutions of $f(y) = y^3 - 2y - 5 = 0$ to illustrate his method.

He guesses that a root is near $y = 2$ since, for example, $f(1) < 0$ but $f(3) > 0$.

A root equals $y = 2 + p$ for some small p (at least, $|p| < 1$). Substituting $y = 2 + p$ into $f(y) = 0$, he gets $g(p) = p^3 + 6p^2 + 10p - 1 = 0$.

Neglecting the p^i terms for $i > 1$ since they are small, he knows $10p - 1 \approx 0$ so $p = 1/10 + q$ for some small q .

Letter 1

We discuss portions of Epistola Prior and the next one (the “Epistola Posterior”). In Epistola Prior, Newton approximates the real solutions of $f(y) = y^3 - 2y - 5 = 0$ to illustrate his method.

He guesses that a root is near $y = 2$ since, for example, $f(1) < 0$ but $f(3) > 0$.

A root equals $y = 2 + p$ for some small p (at least, $|p| < 1$). Substituting $y = 2 + p$ into $f(y) = 0$, he gets $g(p) = p^3 + 6p^2 + 10p - 1 = 0$.

Neglecting the p^i terms for $i > 1$ since they are small, he knows $10p - 1 \approx 0$ so $p = 1/10 + q$ for some small q .

Substituting $p = 1/10 + q$ into $g(p) = 0$, he gets $h(q) = q^3 + 6.3q^2 + 11.23q + 0.061 = 0$.

Letter 1

We discuss portions of Epistola Prior and the next one (the “Epistola Posterior”). In Epistola Prior, Newton approximates the real solutions of $f(y) = y^3 - 2y - 5 = 0$ to illustrate his method.

He guesses that a root is near $y = 2$ since, for example, $f(1) < 0$ but $f(3) > 0$.

A root equals $y = 2 + p$ for some small p (at least, $|p| < 1$). Substituting $y = 2 + p$ into $f(y) = 0$, he gets $g(p) = p^3 + 6p^2 + 10p - 1 = 0$.

Neglecting the p^i terms for $i > 1$ since they are small, he knows $10p - 1 \approx 0$ so $p = 1/10 + q$ for some small q .

Substituting $p = 1/10 + q$ into $g(p) = 0$, he gets $h(q) = q^3 + 6.3q^2 + 11.23q + 0.061 = 0$.

Neglecting higher order terms, he has $11.23q + 0.061 \approx 0$, so $q = -0.0054 + r$ for some r .

Letter 1

We discuss portions of Epistola Prior and the next one (the “Epistola Posterior”). In Epistola Prior, Newton approximates the real solutions of $f(y) = y^3 - 2y - 5 = 0$ to illustrate his method.

He guesses that a root is near $y = 2$ since, for example, $f(1) < 0$ but $f(3) > 0$.

A root equals $y = 2 + p$ for some small p (at least, $|p| < 1$). Substituting $y = 2 + p$ into $f(y) = 0$, he gets $g(p) = p^3 + 6p^2 + 10p - 1 = 0$.

Neglecting the p^i terms for $i > 1$ since they are small, he knows $10p - 1 \approx 0$ so $p = 1/10 + q$ for some small q .

Substituting $p = 1/10 + q$ into $g(p) = 0$, he gets $h(q) = q^3 + 6.3q^2 + 11.23q + 0.061 = 0$.

Neglecting higher order terms, he has $11.23q + 0.061 \approx 0$, so $q = -0.0054 + r$ for some r .

Similarly, he approximates that $r \approx -0.00004852$.

Letter 1

We discuss portions of Epistola Prior and the next one (the “Epistola Posterior”). In Epistola Prior, Newton approximates the real solutions of $f(y) = y^3 - 2y - 5 = 0$ to illustrate his method.

He guesses that a root is near $y = 2$ since, for example, $f(1) < 0$ but $f(3) > 0$.

A root equals $y = 2 + p$ for some small p (at least, $|p| < 1$). Substituting $y = 2 + p$ into $f(y) = 0$, he gets $g(p) = p^3 + 6p^2 + 10p - 1 = 0$.

Neglecting the p^i terms for $i > 1$ since they are small, he knows $10p - 1 \approx 0$ so $p = 1/10 + q$ for some small q .

Substituting $p = 1/10 + q$ into $g(p) = 0$, he gets $h(q) = q^3 + 6.3q^2 + 11.23q + 0.061 = 0$.

Neglecting higher order terms, he has $11.23q + 0.061 \approx 0$, so $q = -0.0054 + r$ for some r .

Similarly, he approximates that $r \approx -0.00004852$. So, $y \approx 2 + (0.1 + (-0.0054 + -0.00004852)) = 2.09455148$.

Historical Part

Application I

Example I

References

Letter 1

Indeed, Newton has the following diagram in his letter.

		$(+ 2,10000000)$ $(- 0,00544852)$ <hr/> $2,09455148$
$2 + p = y$	y^3 $- 2y$ $- 5$	$+ 8 + 12p + 6pp + p^3$ $- 4 - 2p$ $- 5$
	summa	$- 1 + 10p + 6pp + p^3$
$+ 0,1 + q = p$	$+ p^3$ $+ 6pp$ $+ 10p$ $- 1$	$+ 0,001 + 0,03q + 0,3qq + q^3$ $+ 0,06 + 1,2 + 6$ $+ 1 + 10,$ $- 1$
	summa	$0,061 + 11,23q + 6,3qq + q^3$
$- 0,0054 + r = q$	$+ q^3$ $+ 6,3qq$ $+ 11,23q$ $+ 0,061$	$- 0,0000001 + 0,000r \&c$ $+ 0,0001837 - 0,068$ $- 0,060642 + 11,23$ $+ 0,061$
	summa	$+ 0,0005416 + 11,162r$
$- 0,00004852 + s = r$		

Letter 1

There is a similar diagram below the previous diagram, where Newton solves $f(x, y) = y^3 + axy + a^2y - x^3 - 2x^3 = 0$ for y in "almost the same way", meaning that he expresses y as a power series in x .

Letter 1

There is a similar diagram below the previous diagram, where Newton solves $f(x, y) = y^3 + axy + a^2y - x^3 - 2x^3 = 0$ for y in "almost the same way", meaning that he expresses y as a power series in x . Observe how x^n divides each polynomial.

		$\left(a - \frac{x}{4} + \frac{xx}{64a} + \frac{131x^3}{512aa} + \frac{509x^4}{16384a^3} \&c \right)$
$a + p = y$	y^3 $+ axy$ $+ aay$ $- x^3$ $- 2a^3$	$a^3 + 3aap + 3app + p^3$ $+ aax + axp$ $+ a^3 + aap$ $- x^3$ $- 2a^3$
$-\frac{1}{4}x + q = p$	p^3 $+ 3app$ $+ axp$ $+ 4aap$ $+ aax$ $- x^3$	$-\frac{1}{64}x^3 + \frac{3}{16}xxq \&c$ $+ \frac{3}{16}axx - \frac{3}{2}axq + 3aqq$ $- \frac{1}{4}axx + axq$ $- axx + 4aaq$ $+ aax$ $- x^3$
$+\frac{xx}{64a} + r = q$	$3aqq$ $+ \frac{3}{16}xxq$ $- \frac{1}{2}axq$ $+ 4aaq$ $- \frac{65}{64}x^3$ $- \frac{1}{16}axx$	$+\frac{3x^4}{4096a} \&c$ $+\frac{3x^4}{1024a} \&c$ $-\frac{1}{128}x^3 - \frac{1}{2}axr$ $+\frac{1}{16}axx + 4aar$ $-\frac{65}{64}x^3$ $-\frac{1}{16}axx$

Leibniz's Response

Reading Classics

Math Seminar

Zichuan Wang

Historical Part

Application I

Example I

References

A transcribed copy of Epistola Prior was sent by Oldenburg on July 26, 1676, and Leibniz received it only a month later, on August 26 (16), 1676

Leibniz's Response

Reading Classics

Math Seminar

Zichuan Wang

Historical Part

Application I

Example I

References

A transcribed copy of Epistola Prior was sent by Oldenburg on July 26, 1676, and Leibniz received it only a month later, on August 26 (16), 1676 (England was using the Julian Calendar, while Germany was using the Gregorian Calendar).

Leibniz's Response

Reading Classics

Math Seminar

Zichuan Wang

Historical Part

Application I

Example I

References

-

A transcribed copy of *Epistola Prior* was sent by Oldenburg on July 26, 1676, and Leibniz received it only a month later, on August 26 (16), 1676 (England was using the Julian Calendar, while Germany was using the Gregorian Calendar). After a first reading, Leibniz did not understand how Newton got his p, q, r .

Leibniz's Response

A transcribed copy of *Epistola Prior* was sent by Oldenburg on July 26, 1676, and Leibniz received it only a month later, on August 26 (16), 1676 (England was using the Julian Calendar, while Germany was using the Gregorian Calendar). After a first reading, Leibniz did not understand how Newton got his p, q, r . But Leibniz did not want to delay his reply, so he quickly sent out his reply to Oldenburg on August 17, 1676 (Gregorian), in which he included some of his ideas on Newton's results, and a request:

Leibniz's Response

A transcribed copy of Epistola Prior was sent by Oldenburg on July 26, 1676, and Leibniz received it only a month later, on August 26 (16), 1676 (England was using the Julian Calendar, while Germany was using the Gregorian Calendar). After a first reading, Leibniz did not understand how Newton got his p, q, r . But Leibniz did not want to delay his reply, so he quickly sent out his reply to Oldenburg on August 17, 1676 (Gregorian), in which he included some of his ideas on Newton's results, and a request: "But I should like the very distinguished Newton also, to explain some matters more fully, such as such as the origin of the theorem that he places at the beginning, and also the way in which he finds the quantities p, q, r in his operations ...

Leibniz's Response

A transcribed copy of Epistola Prior was sent by Oldenburg on July 26, 1676, and Leibniz received it only a month later, on August 26 (16), 1676 (England was using the Julian Calendar, while Germany was using the Gregorian Calendar). After a first reading, Leibniz did not understand how Newton got his p, q, r . But Leibniz did not want to delay his reply, so he quickly sent out his reply to Oldenburg on August 17, 1676 (Gregorian), in which he included some of his ideas on Newton's results, and a request: "But I should like the very distinguished Newton also, to explain some matters more fully, such as such as the origin of the theorem that he places at the beginning, and also the way in which he finds the quantities p, q, r in his operations ... for he does not explain how it is derived from his method.

Leibniz's Response

A transcribed copy of Epistola Prior was sent by Oldenburg on July 26, 1676, and Leibniz received it only a month later, on August 26 (16), 1676 (England was using the Julian Calendar, while Germany was using the Gregorian Calendar). After a first reading, Leibniz did not understand how Newton got his p, q, r . But Leibniz did not want to delay his reply, so he quickly sent out his reply to Oldenburg on August 17, 1676 (Gregorian), in which he included some of his ideas on Newton's results, and a request: "But I should like the very distinguished Newton also, to explain some matters more fully, such as such as the origin of the theorem that he places at the beginning, and also the way in which he finds the quantities p, q, r in his operations ... for he does not explain how it is derived from his method. I have not yet had the chance to read his letter as carefully as it deserves, as I wished to reply to you forthwith."

Letter 2 on October 24, 1676

Reading Classics
Math Seminar

Zichuan Wang

Historical Part

Application I

Example I

References

-

In Epistola Posterior, Newton explained his way of extracting roots of an affected equation.


In Epistola Posterior, Newton explained his way of extracting roots of an affected equation. Say, we want to solve:

$$f(x, y) = y^6 - 5xy^5 + (x^3/a)y^4 - 7a^2x^2y^2 + 6a^3x^3 + b^2x^4 = 0.$$

Letter 2 on October 24, 1676

In Epistola Posterior, Newton explained his way of extracting roots of an affected equation. Say, we want to solve:


$$f(x, y) = y^6 - 5xy^5 + (x^3/a)y^4 - 7a^2x^2y^2 + 6a^3x^3 + b^2x^4 = 0.$$



Newton drew the diagrams above, the *Newton diagram* of $f(x, y)$, and wrote that

In Epistola Posterior, Newton explained his way of extracting roots of an affected equation. Say, we want to solve:

$$f(x, y) = y^6 - 5xy^5 + (x^3/a)y^4 - 7a^2x^2y^2 + 6a^3x^3 + b^2x^4 = 0.$$




Newton drew the diagrams above, the *Newton diagram* of $f(x, y)$, and wrote that “I apply the ruler DE to the lower of the places marked in the lefthand column, and rotate it from the lower to the higher to the right till it begins to reach likewise another or perhaps several of the remaining marked places.”

Letter 2

Fig. 1

<i>B</i>	x^4	x^4y	x^4y^2	x^4y^3	x^4y^4
x^3	x^3y	x^3y^2	x^3y^3	x^3y^4	
x^2	x^2y	x^2y^2	x^2y^3	x^2y^4	
x	xy	xy^2	xy^3	xy^4	
0	y	y^2	y^3	y^4	



So, a first approximation of a solution to

$$f(x, y) = y^6 - 5xy^5 + (x^3/a)y^4 - 7a^2x^2y^2 + 6a^3x^3 + b^2x^4 = 0$$

is a root of $y^6 - 7a^2x^2y^2 + 6a^3x^3 = 0$. Since the slope is $-1/2$, we want that $y = t(x)^{1/2}$ for some $t \in \mathbb{R}$. Then,

$$x^3(t^6 - 7a^2t^2 + 6a^3) = 0$$

Letter 2

<i>B</i>	x^4	x^4y	x^4y^2	x^4y^3	x^4y^4
x^3	x^3y	x^3y^2	x^3y^3	x^3y^4	
x^2	x^2y	x^2y^2	x^2y^3	x^2y^4	
x	xy	xy^2	xy^3	xy^4	
0	y	y^2	y^3	y^4	

Fig. 1




Fig. 2

So, a first approximation of a solution to

$$f(x, y) = y^6 - 5xy^5 + (x^3/a)y^4 - 7a^2x^2y^2 + 6a^3x^3 + b^2x^4 = 0$$

is a root of $y^6 - 7a^2x^2y^2 + 6a^3x^3 = 0$. Since the slope is $-1/2$, we want that $y = t(x)^{1/2}$ for some $t \in \mathbb{R}$. Then,

$$x^3(t^6 - 7a^2t^2 + 6a^3) = 0 \implies t \in \{\pm\sqrt{a}, \pm\sqrt{2a}\} \text{ could work.}$$

We make a choice $y = \sqrt{ax} + p$, substitute it back to $f(x, y)$ to get a new equation $g(\sqrt{x}, p)$, and continue likewise to get q, r , etc.

Historical Part

Application I

Example I

References

Theorem 1

Theorem

Let $f(x, y) \in \mathbb{C}[x, y]$ be y -general of order $n > 0$,

Theorem 1

Theorem

Let $f(x, y) \in \mathbb{C}[x, y]$ be y -general of order $n > 0$, which means that there exists $n \in \mathbb{Z}_{\geq 1}$ such that the coefficient of y^n is nonzero but the coefficients of y^0, \dots, y^{n-1} are zero.

Theorem 1

Theorem

Let $f(x, y) \in \mathbb{C}[x, y]$ be y -general of order $n > 0$, which means that there exists $n \in \mathbb{Z}_{\geq 1}$ such that the coefficient of y^n is nonzero but the coefficients of y^0, \dots, y^{n-1} are zero.

Then, there exists some $r \in \mathbb{Z}_{\geq 1}$ and a fractional power series $y = \sum_{k=0}^{\infty} c_k (x^{1/r})^k$ such that $f(x, y) = 0$ as a power series in x . □

Theorem 1

Theorem

Let $f(x, y) \in \mathbb{C}[x, y]$ be y -general of order $n > 0$, which means that there exists $n \in \mathbb{Z}_{\geq 1}$ such that the coefficient of y^n is nonzero but the coefficients of y^0, \dots, y^{n-1} are zero.

Then, there exists some $r \in \mathbb{Z}_{\geq 1}$ and a fractional power series $y = \sum_{k=0}^{\infty} c_k (x^{1/r})^k$ such that $f(x, y) = 0$ as a power series in x . □

Proof (popularized by Richard E. Borcherds on Youtube).

Theorem 1

Theorem

Let $f(x, y) \in \mathbb{C}[x, y]$ be y -general of order $n > 0$, which means that there exists $n \in \mathbb{Z}_{\geq 1}$ such that the coefficient of y^n is nonzero but the coefficients of y^0, \dots, y^{n-1} are zero.

Then, there exists some $r \in \mathbb{Z}_{\geq 1}$ and a fractional power series $y = \sum_{k=0}^{\infty} c_k (x^{1/r})^k$ such that $f(x, y) = 0$ as a power series in x . □

Proof (popularized by Richard E. Borcherds on Youtube).

For simplicity, let us assume that the y -order $n = 4$ and the Newton diagram of f looks like the following one:

Theorem 1


Theorem

Let $f(x, y) \in \mathbb{C}[x, y]$ be y -general of order $n > 0$, which means that there exists $n \in \mathbb{Z}_{\geq 1}$ such that the coefficient of y^n is nonzero but the coefficients of y^0, \dots, y^{n-1} are zero.

Then, there exists some $r \in \mathbb{Z}_{\geq 1}$ and a fractional power series $y = \sum_{k=0}^{\infty} c_k (x^{1/r})^k$ such that $f(x, y) = 0$ as a power series in x . □

Proof (popularized by Richard E. Borcherds on Youtube).

For simplicity, let us assume that the y -order $n = 4$ and the Newton diagram of f looks like the following one:



Theorem 1


Theorem

Let $f(x, y) \in \mathbb{C}[x, y]$ be y -general of order $n > 0$, which means that there exists $n \in \mathbb{Z}_{\geq 1}$ such that the coefficient of y^n is nonzero but the coefficients of y^0, \dots, y^{n-1} are zero.

Then, there exists some $r \in \mathbb{Z}_{\geq 1}$ and a fractional power series $y = \sum_{k=0}^{\infty} c_k (x^{1/r})^k$ such that $f(x, y) = 0$ as a power series in x . □

Proof (popularized by Richard E. Borcherds on Youtube).

For simplicity, let us assume that the y -order $n = 4$ and the Newton diagram of f looks like the following one:



Theorem 1


Theorem

Let $f(x, y) \in \mathbb{C}[x, y]$ be y -general of order $n > 0$, which means that there exists $n \in \mathbb{Z}_{\geq 1}$ such that the coefficient of y^n is nonzero but the coefficients of y^0, \dots, y^{n-1} are zero.

Then, there exists some $r \in \mathbb{Z}_{\geq 1}$ and a fractional power series $y = \sum_{k=0}^{\infty} c_k (x^{1/r})^k$ such that $f(x, y) = 0$ as a power series in x . □

Proof (popularized by Richard E. Borcherds on Youtube).

For simplicity, let us assume that the y -order $n = 4$ and the Newton diagram of f looks like the following one:




Proof

With a change of variable $x \rightarrow x^2$ ($\tilde{x} := \sqrt{x}$), all points on the dashed line with integer y -coordinates belong to $\mathbb{Z} \times \mathbb{Z}$.


Proof

With a change of variable $x \rightarrow x^2$ ($\tilde{x} := \sqrt{x}$), all points on the dashed line with integer y -coordinates belong to $\mathbb{Z} \times \mathbb{Z}$.



Proof


With a change of variable $x \rightarrow x^2$ ($\tilde{x} := \sqrt{x}$), all points on the dashed line with integer y -coordinates belong to $\mathbb{Z} \times \mathbb{Z}$.



Observe that the dashed line is given by $3y + x = 12$, and other red points (x, y) not on the dashed line must satisfy $3y + x > 12$.

Proof

With a change of variable $x \rightarrow x^2$ ($\tilde{x} := \sqrt{x}$), all points on the dashed line with integer y -coordinates belong to $\mathbb{Z} \times \mathbb{Z}$.



Observe that the dashed line is given by $3y + x = 12$, and other red points (x, y) not on the dashed line must satisfy $3y + x > 12$. (Define *weight* of (x, y) to be $3y + x$).

Proof

Reading Classics

Math Seminar

Zichuan Wang


Historical Part

Application I

Example I

References

With a change of variable $x \rightarrow x^2$ ($\tilde{x} := \sqrt{x}$), all points on the dashed line with integer y -coordinates belong to $\mathbb{Z} \times \mathbb{Z}$.



Observe that the dashed line is given by $3y + x = 12$, and other red points (x, y) not on the dashed line must satisfy $3y + x > 12$. (Define *weight* of (x, y) to be $3y + x$). Those terms in $f_0(x, y) = 0$ corresponding to red points of weight 12 form a homogeneous polynomial in y, x^3 .

Proof

Since \mathbb{C} is algebraically closed, we can factorize this homogeneous polynomial as follows.

Proof

Since \mathbb{C} is algebraically closed, we can factorize this homogeneous polynomial as follows.

With $\omega := y/x^3$, we have

$$y^4 + a_3(x^3)^3 y + a_2(x^3)^2 y^2 + a_1(x^3)y^3 + a_0(x^3)^4$$

Proof

Since \mathbb{C} is algebraically closed, we can factorize this homogeneous polynomial as follows.

With $\omega := y/x^3$, we have

$$\begin{aligned} & y^4 + a_3(x^3)^3y + a_2(x^3)^2y^2 + a_1(x^3)y^3 + a_0(x^3)^4 \\ &= (x^3)^4 (\omega^4 + a_3\omega^3 + a_2\omega^2 + a_1\omega^1 + a_0) \end{aligned}$$

Proof

Since \mathbb{C} is algebraically closed, we can factorize this homogeneous polynomial as follows.

With $\omega := y/x^3$, we have

$$\begin{aligned} & y^4 + a_3(x^3)^3y + a_2(x^3)^2y^2 + a_1(x^3)y^3 + a_0(x^3)^4 \\ &= (x^3)^4 (\omega^4 + a_3\omega^3 + a_2\omega^2 + a_1\omega^1 + a_0) \\ &= (x^3)^4 \prod_{i=1}^4 (\omega - \alpha_i) \end{aligned}$$

Proof

Since \mathbb{C} is algebraically closed, we can factorize this homogeneous polynomial as follows.

With $\omega := y/x^3$, we have

$$\begin{aligned} & y^4 + a_3(x^3)^3y + a_2(x^3)^2y^2 + a_1(x^3)y^3 + a_0(x^3)^4 \\ &= (x^3)^4 (\omega^4 + a_3\omega^3 + a_2\omega^2 + a_1\omega^1 + a_0) \\ &= (x^3)^4 \prod_{i=1}^4 (\omega - \alpha_i) \\ &= \prod_{i=1}^4 (y - \alpha_i x^3). \end{aligned}$$

Proof

Since \mathbb{C} is algebraically closed, we can factorize this homogeneous polynomial as follows.

With $\omega := y/x^3$, we have

$$\begin{aligned} & y^4 + a_3(x^3)^3y + a_2(x^3)^2y^2 + a_1(x^3)y^3 + a_0(x^3)^4 \\ &= (x^3)^4 (\omega^4 + a_3\omega^3 + a_2\omega^2 + a_1\omega^1 + a_0) \\ &= (x^3)^4 \prod_{i=1}^4 (\omega - \alpha_i) \\ &= \prod_{i=1}^4 (y - \alpha_i x^3). \end{aligned}$$

With a change of variable $y \rightarrow y + \alpha_1 x^3$ ($\tilde{y} := y - \alpha_1 x^3$) with $\alpha_1 \neq 0$,

Proof

Since \mathbb{C} is algebraically closed, we can factorize this homogeneous polynomial as follows.

With $\omega := y/x^3$, we have

$$\begin{aligned} & y^4 + a_3(x^3)^3y + a_2(x^3)^2y^2 + a_1(x^3)y^3 + a_0(x^3)^4 \\ &= (x^3)^4 (\omega^4 + a_3\omega^3 + a_2\omega^2 + a_1\omega^1 + a_0) \\ &= (x^3)^4 \prod_{i=1}^4 (\omega - \alpha_i) \\ &= \prod_{i=1}^4 (y - \alpha_i x^3). \end{aligned}$$

With a change of variable $y \rightarrow y + \alpha_1 x^3$ ($\tilde{y} := y - \alpha_1 x^3$) with $\alpha_1 \neq 0$, we get that

$$y \mid y^4 + b_3(x^3)^3y + b_2(x^3)^2y^2 + b_1(x^3)y^3 + b_0(x^3)^4$$

Proof

Since \mathbb{C} is algebraically closed, we can factorize this homogeneous polynomial as follows.

With $\omega := y/x^3$, we have

$$\begin{aligned} & y^4 + a_3(x^3)^3y + a_2(x^3)^2y^2 + a_1(x^3)y^3 + a_0(x^3)^4 \\ &= (x^3)^4 (\omega^4 + a_3\omega^3 + a_2\omega^2 + a_1\omega^1 + a_0) \\ &= (x^3)^4 \prod_{i=1}^4 (\omega - \alpha_i) \\ &= \prod_{i=1}^4 (y - \alpha_i x^3). \end{aligned}$$

With a change of variable $y \rightarrow y + \alpha_1 x^3$ ($\tilde{y} := y - \alpha_1 x^3$) with $\alpha_1 \neq 0$, we get that

$$y \mid y^4 + b_3(x^3)^3y + b_2(x^3)^2y^2 + b_1(x^3)y^3 + b_0(x^3)^4$$

since the change of variable preserves weight.

Proof

Since \mathbb{C} is algebraically closed, we can factorize this homogeneous polynomial as follows.

With $\omega := y/x^3$, we have

$$\begin{aligned} & y^4 + a_3(x^3)^3y + a_2(x^3)^2y^2 + a_1(x^3)y^3 + a_0(x^3)^4 \\ &= (x^3)^4 (\omega^4 + a_3\omega^3 + a_2\omega^2 + a_1\omega^1 + a_0) \\ &= (x^3)^4 \prod_{i=1}^4 (\omega - \alpha_i) \\ &= \prod_{i=1}^4 (y - \alpha_i x^3). \end{aligned}$$

With a change of variable $y \rightarrow y + \alpha_1 x^3$ ($\tilde{y} := y - \alpha_1 x^3$) with $\alpha_1 \neq 0$, we get that

$$y \mid y^4 + b_3(x^3)^3y + b_2(x^3)^2y^2 + b_1(x^3)y^3 + b_0(x^3)^4$$

since the change of variable preserves weight. Hence, $b_0 = 0$.

Proof

Historical Part


Application I

Example I

References


-

Geometrically, the vanishing of b_0 means that the lowest point on the dashed line is black.



Proof

Geometrically, the vanishing of b_0 means that the lowest point on the dashed line is black.



We are not sure whether the intermediate points are red or black, so let them be blue.

Historical Part

Application I

Example I

References

-

Proof

Do a change of variable $y \rightarrow x^3y$ ($\tilde{y} := y/x^3$).

[Historical Part](#)[Application I](#)[Example I](#)[References](#)

-

Proof

Do a change of variable $y \rightarrow x^3y$ ($\tilde{y} := y/x^3$).

Then, $y^4 \rightarrow x^{12}y^4$, $x^3y^3 \rightarrow x^{12}y^3$, $(x^3)^2y^2 \rightarrow x^{12}y^2$, and
 $(x^3)^3y \rightarrow x^{12}y$.

Proof

Do a change of variable $y \rightarrow x^3y$ ($\tilde{y} := y/x^3$).

Then, $y^4 \rightarrow x^{12}y^4$, $x^3y^3 \rightarrow x^{12}y^3$, $(x^3)^2y^2 \rightarrow x^{12}y^2$, and $(x^3)^3y \rightarrow x^{12}y$.

Dividing everything by x^{12} , the Newton diagram is obtained from the previous one by “left-shifting” w.r.t. y -coordinates.




Figure 1: y

Proof

Do a change of variable $y \rightarrow x^3 y$ ($\tilde{y} := y/x^3$).

Then, $y^4 \rightarrow x^{12} y^4$, $x^3 y^3 \rightarrow x^{12} y^3$, $(x^3)^2 y^2 \rightarrow x^{12} y^2$, and $(x^3)^3 y \rightarrow x^{12} y$.

Dividing everything by x^{12} , the Newton diagram is obtained from the previous one by “left-shifting” w.r.t. y -coordinates.




Figure 1: y




Figure 2: y/x^3 with x^{12} factored out.

If some of $(0, 1), (0, 2), (0, 3)$ is actually red, we repeat the previous procedure with a starting point whose y -coordinate is strictly less than 4.

Historical Part

Application I


Example I

References



-

Proof

If some of $(0, 1), (0, 2), (0, 3)$ is actually red, we repeat the previous procedure with a starting point whose y -coordinate is strictly less than 4. Since there are finitely many integers in $\{1, 2, 3, 4\}$, at one point we have to have that all points directly below the starting point are black:





If some of $(0, 1), (0, 2), (0, 3)$ is actually red, we repeat the previous procedure with a starting point whose y -coordinate is strictly less than 4. Since there are finitely many integers in $\{1, 2, 3, 4\}$, at one point we have to have that all points directly below the starting point are black:



Proof


If some of $(0, 1), (0, 2), (0, 3)$ is actually red, we repeat the previous procedure with a starting point whose y -coordinate is strictly less than 4. Since there are finitely many integers in $\{1, 2, 3, 4\}$, at one point we have to have that all points directly below the starting point are black:



, which means that Newton's ruler should continue to rotate after $y \rightarrow y + \alpha_1 x^3$.


Proof

Rotating the ruler and performing changes of variable, we eliminate the coefficients of $y^i x^j$ for all $i < 4$ and all j (up to arbitrarily large constant N).



Proof

Rotating the ruler and performing changes of variable, we eliminate the coefficients of $y^i x^j$ for all $i < 4$ and all j (up to arbitrarily large constant N).




Thus, $y \rightarrow y + c_1 x^3 + c_2 x^4 + c_3 x^5 + \dots$ makes

$$y^4 \mid \tilde{f} \left(x, y + \sum_{n \geq 1} c_n x^{n+2} \right).$$

So, $\tilde{f}(x, \sum_{n \geq 1} c_n x^{n+2}) = 0$.

Proof

Rotating the ruler and performing changes of variable, we eliminate the coefficients of $y^i x^j$ for all $i < 4$ and all j (up to arbitrarily large constant N).



Thus, $y \rightarrow y + c_1 x^3 + c_2 x^4 + c_3 x^5 + \dots$ makes

$$y^4 \mid \tilde{f} \left(x, y + \sum_{n \geq 1} c_n x^{n+2} \right).$$

So, $\tilde{f}(x, \sum_{n \geq 1} c_n x^{n+2}) = 0$.

Exercise: Finish the proof.

A Worked-out Example

We take the idea of the proof to solve the following equation



$$\begin{aligned}f(x, y) &= 4x^5 - 3x^4 - 4x^2(y - x) + 4(y - x)^2 \\&= 4x^5 - 3x^4 - 4x^2y + 4x^3 + 4y^2 - 8xy + 4x^2 = 0\end{aligned}$$

A Worked-out Example

We take the idea of the proof to solve the following equation

$$\begin{aligned}f(x, y) &= 4x^5 - 3x^4 - 4x^2(y - x) + 4(y - x)^2 \\&= 4x^5 - 3x^4 - 4x^2y + 4x^3 + 4y^2 - 8xy + 4x^2 = 0\end{aligned}$$

whose Newton diagram and graph are





A Worked-out Example

We take the idea of the proof to solve the following equation

$$\begin{aligned}f(x, y) &= 4x^5 - 3x^4 - 4x^2(y - x) + 4(y - x)^2 \\&= 4x^5 - 3x^4 - 4x^2y + 4x^3 + 4y^2 - 8xy + 4x^2 = 0\end{aligned}$$

whose Newton diagram and graph are



Historical Part


Application I

Example I


References

-

$$f(x, y) = 4x^5 - 3x^4 - 4x^2(y - x) + 4(y - x)^2.$$




$$f(x, y) = 4x^5 - 3x^4 - 4x^2(y - x) + 4(y - x)^2.$$



We solve $4y^2 - 8xy - 4x^2 = 4(y - x)^2$ and get $y = x$ as a root.

$$f(x, y) = 4x^5 - 3x^4 - 4x^2(y - x) + 4(y - x)^2.$$



We solve $4y^2 - 8xy - 4x^2 = 4(y - x)^2$ and get $y = x$ as a root. Then, we do $y \rightarrow y + x$ ($y_1 = y - x$) to get

$$4x^5 - 3x^4 - 4x^2y + 4y^2 = 0$$

Now, $y \rightarrow xy$ ($y_2 = y_1/x$) gives us


$$4x^5 - 3x^4 - 4x^2(xy) + 4(xy)^2 = x^2 \underbrace{(4x^3 - 3x^2 - 4xy + 4y^2)}_{g(x,y)}.$$

A Worked-out Example

The Newton diagram of

$$g(x, y) = 4x^3 - 3x^2 - 4xy + 4y^2$$

is



Historical Part

Application I

Example I

References


-

A Worked-out Example

The Newton diagram of

$$g(x, y) = 4x^3 - 3x^2 - 4xy + 4y^2$$

is



Then, we solve $4y^2 - 4xy - 3x^2 = 0$ to get approximate solutions $y = \frac{3}{2}x$ or $y = -\frac{1}{2}x$.

Historical Part

Application I

Example I

References


-

A Worked-out Example

The Newton diagram of

$$g(x, y) = 4x^3 - 3x^2 - 4xy + 4y^2$$

is



Then, we solve $4y^2 - 4xy - 3x^2 = 0$ to get approximate solutions $y = \frac{3}{2}x$ or $y = -\frac{1}{2}x$. Choosing $y = \frac{3}{2}x$, we do $y \rightarrow y + \frac{3}{2}x$ ($y_3 = y_2 - \frac{3}{2}x$).

Historical Part

Application I

Example I

References

-

A Worked-out Example

Substituting $y + \frac{3}{2}x$ into $g(x, y)$, we have

$$g(x, y + \frac{3}{2}x) = 4x^3 - 3x^2 - 4x(y + \frac{3}{2}x) + 4(y + \frac{3}{2}x)^2$$

A Worked-out Example

Substituting $y + \frac{3}{2}x$ into $g(x, y)$, we have

$$\begin{aligned} g(x, y + \frac{3}{2}x) &= 4x^3 - 3x^2 - 4x(y + \frac{3}{2}x) + 4(y + \frac{3}{2}x)^2 \\ &= 4x^3 - 3\cancel{x^2} - 4xy + 6\cancel{x^2} + 4y^2 + 12xy + 9\cancel{x^2} \end{aligned}$$

A Worked-out Example

Substituting $y + \frac{3}{2}x$ into $g(x, y)$, we have

$$\begin{aligned} g(x, y + \frac{3}{2}x) &= 4x^3 - 3x^2 - 4x(y + \frac{3}{2}x) + 4(y + \frac{3}{2}x)^2 \\ &= 4x^3 - 3\cancel{x^2} - 4xy + 6\cancel{x^2} + 4y^2 + 12xy + 9\cancel{x^2} \\ &= 4x^3 + 8xy + 4y^2 \end{aligned}$$

A Worked-out Example

Substituting $y + \frac{3}{2}x$ into $g(x, y)$, we have

$$\begin{aligned} g(x, y + \frac{3}{2}x) &= 4x^3 - 3x^2 - 4x(y + \frac{3}{2}x) + 4(y + \frac{3}{2}x)^2 \\ &= 4x^3 - 3\cancel{x^2} - 4xy + 6\cancel{x^2} + 4y^2 + 12xy + 9\cancel{x^2} \\ &= 4x^3 + 8xy + 4y^2 \end{aligned}$$

With $y \rightarrow xy$ ($y_4 = y_3/x$), we have

$$4x^3 + 8x(xy) + 4(xy)^2 = 4x^2 \underbrace{(x + 2y + y^2)}_{h(x,y)}.$$

A Worked-out Example

Substituting $y + \frac{3}{2}x$ into $g(x, y)$, we have

$$\begin{aligned} g(x, y + \frac{3}{2}x) &= 4x^3 - 3x^2 - 4x(y + \frac{3}{2}x) + 4(y + \frac{3}{2}x)^2 \\ &= 4x^3 - 3\cancel{x^2} - 4xy + 6\cancel{x^2} + 4y^2 + 12xy + 9\cancel{x^2} \\ &= 4x^3 + 8xy + 4y^2 \end{aligned}$$

With $y \rightarrow xy$ ($y_4 = y_3/x$), we have

$$4x^3 + 8x(xy) + 4(xy)^2 = 4x^2 \underbrace{(x + 2y + y^2)}_{h(x,y)}.$$

Here, we can just apply the quadratic formula:

$$y_4 = y = \frac{-2 \pm \sqrt{4 - 4x}}{2} = -1 \pm \sqrt{1 - x}.$$

A Worked-out Example

By Newton's generalized binomial theorem:

$$(x + y)^r = \sum_{k=0}^{\infty} \binom{r}{k} x^{r-k} y^k, \quad \forall r \in \mathbb{R}$$

, we have

$$\sqrt{1-x} = \sum_{k=0}^{\infty} (-1)^k \binom{1/2}{k} x^k$$

A Worked-out Example

By Newton's generalized binomial theorem:

$$(x + y)^r = \sum_{k=0}^{\infty} \binom{r}{k} x^{r-k} y^k, \quad \forall r \in \mathbb{R}$$

, we have

$$\begin{aligned}\sqrt{1-x} &= \sum_{k=0}^{\infty} (-1)^k \binom{1/2}{k} x^k \\ &= 1 - \frac{x}{2} - \frac{x^2}{8} - \frac{x^3}{16} - \frac{5x^4}{128} - \frac{7x^5}{256} + O(x^6)\end{aligned}$$

A Worked-out Example

By Newton's generalized binomial theorem:

$$(x+y)^r = \sum_{k=0}^{\infty} \binom{r}{k} x^{r-k} y^k, \quad \forall r \in \mathbb{R}$$

, we have

$$\begin{aligned}\sqrt{1-x} &= \sum_{k=0}^{\infty} (-1)^k \binom{1/2}{k} x^k \\ &= 1 - \frac{x}{2} - \frac{x^2}{8} - \frac{x^3}{16} - \frac{5x^4}{128} - \frac{7x^5}{256} + O(x^6)\end{aligned}$$


$$\begin{aligned}\text{So, } y_4 &= -\frac{x}{2} - \frac{x^2}{8} - \frac{x^3}{16} - \frac{5x^4}{128} - \frac{7x^5}{256} + O(x^6) \text{ or} \\ y_4 &= -2 + \frac{x}{2} + \frac{x^2}{8} + \frac{x^3}{16} + \frac{5x^4}{128} + \frac{7x^5}{256} + O(x^6).\end{aligned}$$

A Worked-out Example


So, $y = x + (x(\frac{3}{2}x + x(-1 \pm \sqrt{1-x})))$ are two solutions to $f(x, y) = 0$.

A Worked-out Example

So, $y = x + (x(\frac{3}{2}x + x(-1 \pm \sqrt{1-x})))$ are two solutions to $f(x, y) = 0$. Their graphs are:



$$+\sqrt{1-x}$$



$$-\sqrt{1-x}$$

Historical Part

Application I


Example I

References


-

A Worked-out Example


Comparing the following three graphs:



$$-\sqrt{1-x}$$



$$+\sqrt{1-x}$$



$$f(x, y) = 0$$

we see that the self-intersection of the curve $f(x, y) = 0$ at the origin is “resolved”

Historical Part

Application I


Example I

References


-

A Worked-out Example


Comparing the following three graphs:



$$-\sqrt{1-x}$$



$$+\sqrt{1-x}$$



$$f(x, y) = 0$$

we see that the self-intersection of the curve $f(x, y) = 0$ at the origin is “resolved”, whatever that means.

The End

Reading Classics
Math Seminar

Zichuan Wang

Historical Part

Application I

Example I

References

-

Thank you for your attention!

References

[4, 3, 5, 2, 1]



Richard E. Borcherds.

Algebraic geometry 48: Newton's rotating ruler.

2020.



Egbert Brieskorn and Horst Knörrer.

Plane Algebraic Curves.

Modern Birkhäuser Classics, 1986.



Chris Christensen.

Newton's Method for Resolving Affected Equations.

The College Mathematics Journal: Vol. 27, No. 5, pp. 330-340,
1996.



George Chrystal.

Algebra, Vol 2: An Elementary Text-Book for the Higher Classes of Secondary Schools and for Colleges (Classic Reprint).

Forgotten Books, 2018.



Issac Newton.

The Correspondence of Isaac Newton, Volume 2.

Cambridge University Press, 2008.

A Gap in the Proof

In the last step of the proof, we did not show why Newton's ruler could stop only at points (x, y) where $\frac{y-4}{x-0} \in (\mathbb{Z}_{<0})^{-1}$.

A Gap in the Proof

In the last step of the proof, we did not show why Newton's ruler could stop only at points (x, y) where $\frac{y-4}{x-0} \in (\mathbb{Z}_{<0})^{-1}$. We denote the equation of the ruler by $\mu y + x = \nu = 4\mu$, where $\mu = p/q \in \mathbb{Q}_{>0}$ and $p, q \in \mathbb{Z}_{>0}$.

A Gap in the Proof

In the last step of the proof, we did not show why Newton's ruler could stop only at points (x, y) where $\frac{y-4}{x-0} \in (\mathbb{Z}_{<0})^{-1}$. We denote the equation of the ruler by $\mu y + x = \nu = 4\mu$, where $\mu = p/q \in \mathbb{Q}_{>0}$ and $p, q \in \mathbb{Z}_{>0}$.

Then, $\tilde{x} = x^{1/q}$ (so, $x = \tilde{x}^q$) and $\tilde{y} = (y - c\tilde{x}^p)/\tilde{x}^p$ (so, $y = \tilde{x}^p(\tilde{y} + c)$) with $c \neq 0$.

A Gap in the Proof

In the last step of the proof, we did not show why Newton's ruler could stop only at points (x, y) where $\frac{y-4}{x-0} \in (\mathbb{Z}_{<0})^{-1}$. We denote the equation of the ruler by $\mu y + x = \nu = 4\mu$, where $\mu = p/q \in \mathbb{Q}_{>0}$ and $p, q \in \mathbb{Z}_{>0}$.

Then, $\tilde{x} = x^{1/q}$ (so, $x = \tilde{x}^q$) and $\tilde{y} = (y - c\tilde{x}^p)/\tilde{x}^p$ (so, $y = \tilde{x}^p(\tilde{y} + c)$) with $c \neq 0$.

$$f(x, y) = \underbrace{\sum_{\mu\beta+\alpha=4\mu} b_{\alpha\beta} x^\alpha y^\beta}_{\text{on the line}} + \underbrace{\sum_{\mu\beta+\alpha>4\mu} b_{\alpha\beta} x^\alpha y^\beta}_{\text{above the line}}$$

A Gap in the Proof

In the last step of the proof, we did not show why Newton's ruler could stop only at points (x, y) where $\frac{y-4}{x-0} \in (\mathbb{Z}_{<0})^{-1}$. We denote the equation of the ruler by $\mu y + x = \nu = 4\mu$, where $\mu = p/q \in \mathbb{Q}_{>0}$ and $p, q \in \mathbb{Z}_{>0}$.

Then, $\tilde{x} = x^{1/q}$ (so, $x = \tilde{x}^q$) and $\tilde{y} = (y - c\tilde{x}^p)/\tilde{x}^p$ (so, $y = \tilde{x}^p(\tilde{y} + c)$) with $c \neq 0$.

$$\begin{aligned} f(x, y) &= \underbrace{\sum_{\mu\beta+\alpha=4\mu} b_{\alpha\beta} x^\alpha y^\beta}_{\text{on the line}} + \underbrace{\sum_{\mu\beta+\alpha>4\mu} b_{\alpha\beta} x^\alpha y^\beta}_{\text{above the line}} \\ &= \tilde{x}^{4p} \underbrace{\left(\sum_{\mu\beta+\alpha=4\mu} b_{\alpha\beta} (\tilde{y} + c)^\beta + \sum_{\mu\beta+\alpha>4\mu} b_{\alpha\beta} \tilde{x}^{p\alpha+q\beta-4p} (\tilde{y} + c)^\beta \right)}_{\tilde{f}(\tilde{x}, \tilde{y})} \end{aligned}$$

A Gap in the Proof

In the last step of the proof, we did not show why Newton's ruler could stop only at points (x, y) where $\frac{y-4}{x-0} \in (\mathbb{Z}_{<0})^{-1}$. We denote the equation of the ruler by $\mu y + x = \nu = 4\mu$, where $\mu = p/q \in \mathbb{Q}_{>0}$ and $p, q \in \mathbb{Z}_{>0}$.

Then, $\tilde{x} = x^{1/q}$ (so, $x = \tilde{x}^q$) and $\tilde{y} = (y - c\tilde{x}^p)/\tilde{x}^p$ (so, $y = \tilde{x}^p(\tilde{y} + c)$) with $c \neq 0$.

$$\begin{aligned} f(x, y) &= \underbrace{\sum_{\mu\beta+\alpha=4\mu} b_{\alpha\beta} x^\alpha y^\beta}_{\text{on the line}} + \underbrace{\sum_{\mu\beta+\alpha>4\mu} b_{\alpha\beta} x^\alpha y^\beta}_{\text{above the line}} \\ &= \tilde{x}^{4p} \underbrace{\left(\sum_{\mu\beta+\alpha=4\mu} b_{\alpha\beta} (\tilde{y} + c)^\beta + \sum_{\mu\beta+\alpha>4\mu} b_{\alpha\beta} \tilde{x}^{p\alpha+q\beta-4p} (\tilde{y} + c)^\beta \right)}_{\tilde{f}(\tilde{x}, \tilde{y})} \end{aligned}$$

So, $\tilde{f}(0, \tilde{y}) = \sum_{\mu\beta+\alpha=4\mu} b_{\alpha\beta} (\tilde{y} + c)^\beta$ is a quartic in \tilde{y} .

A Gap in the Proof

So, $\tilde{f}(0, \tilde{y}) = \sum_{\mu\beta+\alpha=4\mu} b_{\alpha\beta}(\tilde{y} + c)^\beta$ is a quartic in \tilde{y} .

Historical Part

Application I

Example I

References

-

A Gap in the Proof

So, $\tilde{f}(0, \tilde{y}) = \sum_{\mu\beta+\alpha=4\mu} b_{\alpha\beta}(\tilde{y} + c)^\beta$ is a quartic in \tilde{y} . By our assumption that the smallest power of \tilde{y} with nonzero coefficient is 4, the coefficients of $\tilde{y}^0, \tilde{y}^1, \tilde{y}^2, \tilde{y}^3$ in $\tilde{f}(0, \tilde{y})$ are 0.

Historical Part

Application I

Example I

References

A Gap in the Proof

Historical Part

Application I

Example I

References

So, $\tilde{f}(0, \tilde{y}) = \sum_{\mu\beta+\alpha=4\mu} b_{\alpha\beta}(\tilde{y} + c)^\beta$ is a quartic in \tilde{y} . By our assumption that the smallest power of \tilde{y} with nonzero coefficient is 4, the coefficients of $\tilde{y}^0, \tilde{y}^1, \tilde{y}^2, \tilde{y}^3$ in $\tilde{f}(0, \tilde{y})$ are 0. So, with $\tilde{y} = t - c$,

$$g(t) := \sum_{\mu\beta+\alpha=4\mu} b_{\alpha\beta}t^\beta = b_{04}(t - c)^4.$$

A Gap in the Proof

Historical Part

Application I

Example I

References

So, $\tilde{f}(0, \tilde{y}) = \sum_{\mu\beta+\alpha=4\mu} b_{\alpha\beta}(\tilde{y} + c)^\beta$ is a quartic in \tilde{y} . By our assumption that the smallest power of \tilde{y} with nonzero coefficient is 4, the coefficients of $\tilde{y}^0, \tilde{y}^1, \tilde{y}^2, \tilde{y}^3$ in $\tilde{f}(0, \tilde{y})$ are 0. So, with $\tilde{y} = t - c$,

$$g(t) := \sum_{\mu\beta+\alpha=4\mu} b_{\alpha\beta}t^\beta = b_{04}(t - c)^4.$$

Compare the coefficient of t^3 on both sides:

$$b_{\mu 3} = -4b_{04}c$$

A Gap in the Proof

So, $\tilde{f}(0, \tilde{y}) = \sum_{\mu\beta+\alpha=4\mu} b_{\alpha\beta}(\tilde{y} + c)^\beta$ is a quartic in \tilde{y} . By our assumption that the smallest power of \tilde{y} with nonzero coefficient is 4, the coefficients of $\tilde{y}^0, \tilde{y}^1, \tilde{y}^2, \tilde{y}^3$ in $\tilde{f}(0, \tilde{y})$ are 0. So, with $\tilde{y} = t - c$,

$$g(t) := \sum_{\mu\beta+\alpha=4\mu} b_{\alpha\beta}t^\beta = b_{04}(t - c)^4.$$

Compare the coefficient of t^3 on both sides:

$$b_{\mu 3} = -4b_{04}c \neq 0 \quad (\text{since } \text{char}(\mathbb{C}) = 0).$$

A Gap in the Proof

So, $\tilde{f}(0, \tilde{y}) = \sum_{\mu\beta+\alpha=4\mu} b_{\alpha\beta}(\tilde{y} + c)^\beta$ is a quartic in \tilde{y} . By our assumption that the smallest power of \tilde{y} with nonzero coefficient is 4, the coefficients of $\tilde{y}^0, \tilde{y}^1, \tilde{y}^2, \tilde{y}^3$ in $\tilde{f}(0, \tilde{y})$ are 0. So, with $\tilde{y} = t - c$,

$$g(t) := \sum_{\mu\beta+\alpha=4\mu} b_{\alpha\beta}t^\beta = b_{04}(t - c)^4.$$

Compare the coefficient of t^3 on both sides:

$$b_{\mu 3} = -4b_{04}c \neq 0 \quad (\text{since } \text{char}(\mathbb{C}) = 0).$$

So, $\mu \in \mathbb{Z}_{>0}$.



A Gap in the Proof

So, $\tilde{f}(0, \tilde{y}) = \sum_{\mu\beta+\alpha=4\mu} b_{\alpha\beta}(\tilde{y} + c)^\beta$ is a quartic in \tilde{y} . By our assumption that the smallest power of \tilde{y} with nonzero coefficient is 4, the coefficients of $\tilde{y}^0, \tilde{y}^1, \tilde{y}^2, \tilde{y}^3$ in $\tilde{f}(0, \tilde{y})$ are 0. So, with $\tilde{y} = t - c$,

$$g(t) := \sum_{\mu\beta+\alpha=4\mu} b_{\alpha\beta}t^\beta = b_{04}(t - c)^4.$$

Compare the coefficient of t^3 on both sides:

$$b_{\mu 3} = -4b_{04}c \neq 0 \quad (\text{since } \text{char}(\mathbb{C}) = 0).$$

So, $\mu \in \mathbb{Z}_{>0}$.



Kiran Kedlaya: <https://arxiv.org/abs/math/9810142>