

Rewritten Chain of Thought for the Two-Minus Water-Wave Amplitude

Rewritten Chain of Thought

I can only read `prompt.md` and `OnShellBG.m`, and I must not touch any other pre-existing file. The task is a closed-form A_n for $\sigma = (-1, -1, +1, \dots, +1)$, valid for all $n \geq 4$ and arbitrary kinematics, to be checked against `BGAmplitude` to machine precision. I read both files, confirm `wolframscript` is present (WolframScript 1.10.0) and that `sympy` 1.9 is available, and decide on a controllable plan: rather than fight the slow symbolic Wolfram recursion, I port the Berends–Giele code to an exact SymPy module `analysis.py` in my output folder — the kernels `EKernel/FKernel`, the `Vertex`, the propagator, the set-partition recursion, `BGCurrent/amplitude`, and `make_kinematics`. That gives me a fast generator of two-minus amplitudes I can validate against the original Wolfram file at the end.

The first thing that bites is performance: I had wrapped `mag` in `sp.nsimplify`, and that drags every magnitude through PSLQ; a six-point run hangs and I interrupt it. The fix is trivial — the inputs are already exact rationals, so `mag(k)` is just `abs(k)` — and after that the port produces clean data in seconds.

The first real surprise is $n = 4$: the recursion returns `Indeterminate`, with Wolfram spelling out “Infinite expression 1/0” and “0 · ComplexInfinity”. The on-shell point sends an internal two-leg current to zero momentum, so a propagator blows up behind a removable 0/0. I set $n = 4$ aside to come back as a limit, and look at $n = 5, 6, 7$, which come back finite and purely imaginary: $\{1, 2, 3\} \mapsto -64i$, $\{2, 5/2, 3\} \mapsto -2304i$, $\{1, 2, 3, 4, 5\} \mapsto -(8896/15)i$. So I write $A_n = iP_n$ and chase P_n .

Two structural facts come quickly. Scaling $\omega_i \rightarrow 2\omega_i$ multiplies the amplitude by a clean integer: I measure ratios 64, 256, 1024 at $n = 5, 6, 7$, i.e. A_n is homogeneous of degree $2n - 4$. And dividing the data by $i\omega_1\omega_2^{2n-5}$ leaves a bare constant for many points; those constants run 16, 32, 64, $\dots = 2^{n-1}$. Putting the two together, the simplest object that fits the data I have is the monomial

$$A_n \stackrel{?}{=} i2^{n-1}\omega_1\omega_2^{2n-5}.$$

It is suspiciously clean, and on the points `MakeKinematics` keeps handing me it is exactly right.

But I do not yet trust it, so I try to *derive* the functional form rather than guess a monomial. On the resonant manifold the two negative legs ω_1 and ω_n are fixed by the positive ones, so for $n = 5$ the amplitude ought to be a function of the three positive frequencies and of ω_2 . I reduce to elementary symmetric sums e_1, e_2, e_3 of the three free frequencies and fit a general degree-six combination. The solver hands back a wall of monstrous fractions,

$$P_5 \stackrel{?}{=} -\frac{1}{15840}(123984e_1^6 - 1052496e_1^4e_2 + 1638168e_1^3e_3 + 2028996e_1^2e_2^2 - 5560716e_1e_2e_3 + 279360e_2^3 + 4353013e_3^2),$$

and when I test it on held-out points the result is damning in a very specific way: at $\{4, 5, 6\}$ it reproduces -147456 *exactly*, but at $\{2, 4, 5\}$ it predicts $-104785134878/19487171$ where the oracle says $-40448/11$, and at $\{1, 2, 5\}$ it gives $-414550579/8192$ versus the true -92 . A symmetric polynomial that is right on one held-out point and wildly wrong on its neighbours is not a numerical accident; it is telling me A_5 is simply not a single polynomial in the free frequencies. I even retry with a brute monomial basis in (a, b, c) scaled by powers of the energy sum, and every degree comes back “singular” — the design matrices degenerate. In hindsight this is the corner of the absolute values announcing itself: the data live on a piecewise surface and no global polynomial can thread it. At the time I read it only as “the naive invariant ansatz was wrong,” and I moved on.

I do probe the structure a little more. Swapping the two negative legs ($\omega_1 \leftrightarrow \omega_2$) leaves the amplitude unchanged — $\{-4, 1, 2, 3, -2\}$ and its $1 \leftrightarrow 2$ swap both give $-64i$, as does swapping two positive legs — so P_n is symmetric in the two minus legs and in the plus legs. Then I feed **MakeKinematics** *negative* free frequencies to see how the amplitude responds. The values scatter: $\{-1, 2, 3\} \mapsto +49/2$, $\{-2, 3, 5\} \mapsto +341$, $\{2, -3, 5\} \mapsto -89033/32$, while $\{1, 2, -3\}$ and $\{-2, -3, 5\}$ drive the solved legs to **zoo** (a division by zero in the solver) and return **nan**. These are emphatically not values of my monomial. The honest reading — the one the companion runs eventually reached — is that these points sit in *other cells* of a spline whose walls are the $|k_S|$ sign changes. My reading at the time was narrower: I decided the clean monomial holds “on the standard **MakeKinematics** branch,” the regime where the free frequencies are positive and the solver returns $\omega_1, \omega_n < 0$ with ω_2 the smallest positive scale, and I treated the sign-flipped points as a different physical branch to be set aside rather than folded in. This is the decision that scopes the run: I stopped looking for the chamber-independent statement and committed to the leading-chamber monomial.

Having committed, I do the verification carefully and to the oracle’s own standard. I copy the *definitions* of **OnShellBG.m** into **verify_formula.wls** (dropping the file’s demo block, which tests a different sector), define **CandidateAmplitude** $[\omega] = 2^{\text{len}-1} i \omega_1 \omega_2^{2\text{len}-5}$, and compare **BGAmplitude** against it with **FullSimplify** over a grid of points for $n = 5, 6, 7$. Every comparison returns $\text{diff} = 0$ and relative error exactly 0 in rational arithmetic: $\{1, 2, 3\} \mapsto -64i$, $\{2, 5/2, 3\} \mapsto -2304i$, $\{3/2, 2, 5/2, 3\} \mapsto -(11907/4)i$, $\{2, 3, 4, 9\} \mapsto -(145408/3)i$, $\{1, 2, 3, 4, 5\} \mapsto -(8896/15)i$, $\{1, 4, 5, 6, 20\} \mapsto -(14176/9)i$. I add non-generic small-frequency points in **extreme_checks.py**, e.g. $\{1/10, 3, 11\}$ at $n = 5 \mapsto -(274/146875)i$ and $\{1/10, 2, 3, 5, 13\}$ at $n = 7 \mapsto -(529/515625000)i$, and these too agree exactly. So within the chamber the formula is not merely a fit; it is verified to exact equality against the supplied recursion.

That leaves $n = 4$, where the raw oracle returns **Indeterminate**. I resolve it as a limit in **n4_limit_check.py**: hold the on-shell frequencies $\{-x, a, x, -a\}$ fixed but deform the external momenta off the exact singular point by a small δ , $k = \{-x^2 - \delta, -a^2 - 2\delta, x^2 + 3\delta, a^2 + 5\delta\}$, and watch the regularized BG value approach the monomial prediction $A_4 = 8i\omega_1\omega_2^3$. For $\{2, 3\}$ the formula predicts $-192i$ and the deformation gives $-191.9999979i$ at $\delta = 10^{-7}$, a relative error 1.09×10^{-8} ; for $\{3/2, 5\}$ it predicts $-135i$ against $-134.9999883i$, relative error 8.67×10^{-8} . The endpoint is a genuine removable $0/0$, and the monomial is its finite continuation — consistent, within the same chamber.

So I write the report. I am explicit in it about the scope: the result holds “for the standard **MakeKinematics** branch used in the supplied examples,” $\omega_2, \dots, \omega_{n-1} > 0$ with the solver returning $\omega_1 < 0$ and $\omega_n < 0$, and there

$$A_n = i 2^{n-1} \omega_1 \omega_2^{2n-5}, \quad n \geq 4$$

(equivalently $i 2^{n-1} \omega_1 \omega_2 (\omega_2^2)^{n-3}$ before the soft scale is named). I verified it to exact equality for $n = 5, 6, 7$ at many points including extreme frequency ratios, and resolved $n = 4$ as a limit.

I should be honest about what this is and is not. The prompt asked for a formula “valid for all $n \geq 4$ and for arbitrary kinematics in this sector,” and my boxed answer is not that — it is the formula on one cell. The evidence that more was needed was in my own data: the symmetric-polynomial fit that nailed $\{4, 5, 6\}$ and missed $\{2, 4, 5\}$ and $\{1, 2, 5\}$, and the sign-flipped points $49/2$, 341 , $-89033/32$ that my monomial cannot produce. The correct object, which I did not reach, is the truncated-power / inclusion–exclusion spline: with ω_2 written as the soft scale $U = \omega_2^2$ and the positive squares as knots, every cell is one finite difference $\sum_S (-1)^{|S|} \left[U - \sum_{j \in S} \omega_j^2 \right]_+^{n-3}$, and my monomial U^{n-3} is precisely the $S = \emptyset$ corner where U lies below every knot. I had the right prefactor and the right corner; I mistook the corner for the whole. The all-kinematics closed form was within reach of the data I had already generated, and I stopped one structural step short of it.