



# 浙江大学爱丁堡大学联合学院 ZJU-UoE Institute

## Conditional Probabilities

ADS 2, Lecture 22







Zhaoyuan Fang -  
zhaoyuanfang@intl.zju.edu.cn

Semester 2, 2022/23

*Based on Prof MI Stefan's slides*

# A card game (1)

I have a (large) number of cards. Each card has a letter on one side and a number on the other side

Front			
Back			


# A card game (1)

Hypothesis: Every card that has a **vowel** on the front side has an **even number** on the back side.

## A card game (1)






Hypothesis: Every card that has a **vowel** on the front side has an **even** **number** on the back side.

Question: Which cards do I have to turn around to test this hypothesis?




## A card game (2)

I have three cards: one pink on both sides, one blue on both sides, one with one pink and one blue side



## A card game (2)



Question: I drew a card and am showing you one side. It's blue. What is the probability that the other side is also blue?

# This lecture is about ...

Probabilities of combinations of events  
(partly a review from last year, but with a bit more depth and rigour)

# Learning Objectives

After this week, you will be able to ...

- Recall how to compute conditional probabilities
- Visualise joint probabilities using Euler diagrams and probability trees
- State and apply Bayes' theorem
- Describe and use Markov chains



# Outline

- 1 [Logical foundations](#)
- 2 [Conditional and joint probabilities](#)
- 3 [Bayes' theorem](#)
- 4 [Markov chains](#)

# A tiny little bit of logic notation

$\neg A$	"not A"	True if A is false
$A \& B$	"A and B"	True if both A and B are true, false otherwise
$A \vee B$	"A or B"	True if A is true or B is true (or both)
$A \rightarrow B$	"If A then B"	

# A tiny little bit of logic notation

$\neg A$	"not A"	True if A is false
$A \& B$	"A and B"	True if both A and B are true, false otherwise
$A \vee B$	"A or B"	True if A is true or B is true (or both)
$A \rightarrow B$	"If A then B"	?


$A$	$B$	$A \rightarrow B$
T	T	
T	F	
F	T	
F	F	

# Back to our card game

**Hypothesis:** Every card that has a **vowel** on the front side has an **even** number on the back side.

**vowel** → **even number**

**Question:** Which cards do I have to turn around to test this hypothesis?



# A tiny little bit of logic notation

$\neg A$	"not A"	True if A is false
$A \& B$	"A and B"	True iff both A and B are true
$A \vee B$	"A or B"	True if A is true or B is true (or both)
$A \rightarrow B$	"If A then B"	$(\neg A) \vee B$
$A \leftrightarrow B$	"If and only if A then B"	
	"Iff A then B"	

# A tiny little bit of logic notation

$\neg A$  "not A"

True if A is false

$A \& B$  "A and B"

True iff both A and B are true

$A \vee B$  "A or B"

True if A is true or B is true (or both)

$A \rightarrow B$  "If A then B"

$(\neg A) \vee B$

$A \leftrightarrow B$  "If and only if A then B"

"Iff A then B"

?

A	B	$A \rightarrow B$
T	T	
T	F	
F	T	
F	F	

# Outline

- 1 [Logical foundations](#)
- 2 [Conditional and joint probabilities](#)
- 3 [Bayes' theorem](#)
- 4 [Markov chains](#)

# Conditional probabilities: A bit of notation

$$P(A|B)$$

“Probability of A given B”: Probability of A *if B is true*

## Examples (reminder)

- **Probability** of having a disease given a **positive test result**
- Probability of a person getting a disease given they are a carrier for a specific allele variant
- Probability of cell survival given treatment with a toxic chemical
- Probability of seeing a result as or more extreme as the one in your experiment given the Null Hypothesis is true



# Example: Lie detector test


## Problem

In a big store, management finds out that around 10 % of employees must be stealing, but they don't know who.

In response to this, all employees have to go through a lie detector test. The lie detector has 80 % accuracy in both directions: It correctly categorises 80 % of the people telling the truth as telling the truth, and 80 % of the people lying as lying.

Every employee was tested and everybody said they did not steal. According to the lie detector, 50 employees were lying. How many were thieves?

## A convenient tool: Euler diagrams



# A convenient tool: Euler diagrams

## Problem

In a big store, management finds out that around 10 % of employees must be stealing, but they don't know who.

In response to this, all employees have to go through a lie detector test. The lie detector has 80 % accuracy in both directions: It correctly categorises 80 % of the people telling the truth as telling the truth, and 80 % of the people lying as lying.

Every employee was tested and everybody said they did not steal.


According to the lie detector, 50 employees were lying.

Draw an Euler diagram of the situation

# A convenient tool: Euler diagrams

## Review: Probability trees


Another convenient way to visualise and compute joint probabilities.  
Multiply probabilities along branches.



# Looking back to our second card game

I have three cards: one pink on both sides, one blue on both sides, one with one pink and one blue side

Question: I drew a card and am showing you one side. It's blue. What is the probability that the other side is also blue?



Draw a probability tree for this problem

# Outline

- 1 [Logical foundations](#)
- 2 [Conditional and joint probabilities](#)
- 3 [Bayes' theorem](#)
- 4 [Markov chains](#)

# Bayes' theorem



We already know that

$$P(A|B) \neq P(B|A)$$

We already know that

$$P(A|B) \neq P(B|A)$$

But how *are*  $P(A|B)$  and  $P(B|A)$  related?



Thomas Bayes

We already know that

$$P(A|B) \neq P(B|A)$$

But how *are*  $P(A|B)$  and  $P(B|A)$  related?



Thomas Bayes (maybe)

We already know that


$$P(A|B) \neq P(B|A)$$

But how *are*  $P(A|B)$  and  $P(B|A)$  related?



Thomas Bayes (maybe)  
University of Edinburgh alumni!

# Bayes' theorem

A photograph of a blue neon sign mounted on a ceiling, displaying the formula for Bayes' theorem. The sign is illuminated with a bright blue light, and the background is dark. The formula is written in a stylized, hand-drawn font. The text is  $P(A|B) = \frac{P(B|A)P(A)}{P(B)}$ . The sign is slightly tilted and has some visible wiring and mounting hardware.
$$P(A|B) = \frac{P(B|A)P(A)}{P(B)}$$

# Let's prove it!

We already know

$$P(A \& B) = P(A) \times P(B|A)$$

# Let's prove it!

We already know

$$P(A \& B) = P(A) \times P(B|A)$$

Similarly

$$P(B \& A) = P(B) \times P(A|B)$$

# Let's prove it!

We already know

$$P(A \& B) = P(A) \times P(B|A)$$

Similarly

$$P(B \& A) = P(B) \times P(A|B)$$

But of course,

$$P(A \& B) = P(B \& A)$$



# Let's prove it!

We already know

$$P(A \& B) = P(A) \times P(B|A)$$

Similarly

$$P(B \& A) = P(B) \times P(A|B)$$

But of course,

$$P(A \& B) = P(B \& A)$$

Therefore,

$$P(A) \times P(B|A) = P(B) \times P(A|B)$$

$$P(A|B) = \frac{P(A) \times P(B|A)}{P(B)} \quad .$$

# Bayes' theorem: Example

What is  $P(\text{thief} | \text{lied according to the lie detector})$ ?

# Bayes' theorem: Example

What is  $P(\text{thief} | \text{lied according to the lie detector})$ ?

*Practical!*


# Outline

- 1 [Logical foundations](#)
- 2 [Conditional and joint probabilities](#)
- 3 [Bayes' theorem](#)
- 4 [Markov chains](#)

# What are Markov chains?

- Stochastic model
- A system is represented as being in a number of possible states
- Transitions between states happen with specified probabilities
- Probabilities of state transitions depend on the state the system is currently in, not its history
- Probabilities going out of any one node should add up to 1

# Markov chains: Simple example




# Markov chains: More complicated example

If tossing a fair coin (H=Head, T=Tail), how long would it take to get the sequence H-T-T-H?

Start

# Markov chains: More complicated example


If tossing a fair coin (H=Head, T=Tail), how long would it take to get the sequence H-T-T-H?






# Markov chains: More complicated example

If tossing a fair coin (H=Head, T=Tail), how long would it take to get the sequence H-T-T-H?




# Markov chains: More complicated example

If tossing a fair coin (H=Head, T=Tail), how long would it take to get the sequence H-T-T-H?



# Markov chains: More complicated example

If tossing a fair coin (H=Head, T=Tail), how long would it take to get the sequence H-T-T-H?



*Practical!*

- There is a mathematical theory of Markov chains, with ways to compute probabilities to reach states, path lengths etc.
- For this course, we ask you to do 2 things
  - Draw up a Markov chain for a new problem
  - Write code to allow you to simulate a Markov chain many times

# What questions do you have?

After this week, you will be able to ...

- Recall how to compute conditional probabilities
- Visualise joint probabilities using Euler diagrams and probability trees
- State and apply Bayes' theorem
- Describe and use Markov chains

# Image credits

- Bayes' Theorem in neon letters. By mattbuck (category) - Own work by mattbuck., CC BY-SA 3.0, <https://commons.wikimedia.org/w/index.php?curid=14658489>
- Card games. My own work (2020), CC-BY-SA 4.0.
- Euler diagram. By Gnathan87 - Own work, CC0, <https://commons.wikimedia.org/w/index.php?curid=15991401>
- Euler diagram for the examples of thieves in the store. My own work (2020), CC-BY-SA 4.0.
- Markov chain. My own work (2020), CC-BY-SA 4.0.
- Portrait of a man who may or may not be Thomas Bayes. Public domain, 19<sup>th</sup> century. Via Wikimedia Commons.
- Probability tree. My own work (2019), CC BY-SA 3.0.
- Probability tree for the card problem. My own work (2020), CC BY-SA 4.0.