

Homework 3

Ziming Ji

PHY 539: Introduction to String Theory

October 31, 2018

1 Problem 1

a) It is not hard to show that under $z \rightarrow \frac{az+b}{cz+d}$, the cross ratio becomes

$$\frac{\left(\frac{az_1+b}{cz_1+d} - \frac{az_2+b}{cz_2+d}\right) \left(\frac{az_3+b}{cz_3+d} - \frac{az_4+b}{cz_4+d}\right)}{\left(\frac{az_1+b}{cz_1+d} - \frac{az_3+b}{cz_3+d}\right) \left(\frac{az_2+b}{cz_2+d} - \frac{az_4+b}{cz_4+d}\right)} = \frac{(z_1 - z_2)(z_3 - z_4)(bc - ad)^2}{(z_1 - z_3)(z_2 - z_4)(bc - ad)^2} = \frac{(z_1 - z_2)(z_3 - z_4)}{(z_1 - z_3)(z_2 - z_4)}. \quad (1)$$

b) We set the transformation to be $z \rightarrow \frac{az+b}{cz+d}$. Solving the set of equations

$$az_1 + b = 0 \wedge az_2 + b = cz_2 + d \wedge cz_3 + d = 0 \wedge ad - bc = 1, \quad (2)$$

we have

$$\begin{aligned} a &\rightarrow -\frac{\sqrt{z_3 - z_2}}{\sqrt{(z_1 - z_2)(z_1 - z_3)}}, b \rightarrow \frac{z_1 \sqrt{z_3 - z_2}}{\sqrt{(z_1 - z_2)(z_1 - z_3)}}, \\ c &\rightarrow \frac{z_2 - z_1}{\sqrt{(z_1 - z_2)(z_1 - z_3)}\sqrt{z_3 - z_2}}, d \rightarrow \frac{(z_1 - z_2)z_3}{\sqrt{(z_1 - z_2)(z_1 - z_3)}\sqrt{z_3 - z_2}} \end{aligned} \quad (3)$$

or

$$\begin{aligned} a &\rightarrow \frac{\sqrt{z_3 - z_2}}{\sqrt{(z_1 - z_2)(z_1 - z_3)}}, b \rightarrow -\frac{z_1 \sqrt{z_3 - z_2}}{\sqrt{(z_1 - z_2)(z_1 - z_3)}}, \\ c &\rightarrow \frac{z_1 - z_2}{\sqrt{(z_1 - z_2)(z_1 - z_3)}\sqrt{z_3 - z_2}}, d \rightarrow \frac{(z_2 - z_1)z_3}{\sqrt{(z_1 - z_2)(z_1 - z_3)}\sqrt{z_3 - z_2}}. \end{aligned} \quad (4)$$

2 Problem 2

We refer to the b c system as a $2D$ field theory with a Grassmann action

$$S = \frac{1}{2\pi} \int d^2 z \, b \bar{\partial} c \quad (5)$$

a)

$$\begin{aligned}
T(z)T(w) &= (: (\partial_z b)c(z) : -\lambda \partial_z : bc(z) :) (: (\partial_w b)c(w) : -\lambda \partial_w : bc(w) :) \\
&=: (\partial_z b)c(z) :: (\partial_w b)c(w) : -\lambda \partial_z : bc(z) :: (\partial_w b)c(w) : \\
&\quad -\lambda : (\partial_z b)c(z) : \partial_w : bc(w) : +\lambda^2 \partial_z : bc(z) : \partial_w : bc(w) :
\end{aligned} \tag{6}$$

The full contraction(quartic) part is

$$\lambda^2 \frac{\partial}{\partial z} \frac{\partial}{\partial \omega} \frac{1}{(z-\omega)(z-\omega)} - \lambda \frac{\partial}{\partial \omega} \frac{\frac{\partial}{\partial z} \frac{1}{z-\omega}}{z-\omega} - \lambda \frac{\partial}{\partial z} \frac{\frac{\partial}{\partial \omega} \frac{1}{z-\omega}}{z-\omega} + \frac{\partial}{\partial z} \frac{1}{z-\omega} \frac{\partial}{\partial \omega} \frac{1}{z-\omega} = \frac{-6(\lambda-1)\lambda-1}{(z-\omega)^4} \tag{7}$$

The quadratic part is

$$123 \tag{8}$$