

Urban biogeography of fungal endophytes across San Francisco

Emma Gibson¹ and Naupaka Zimmerman¹

¹Department of Biology, University of San Francisco

Corresponding author:

Naupaka Zimmerman¹

Email address: nzimmerman@usfca.edu

ABSTRACT

In natural and agricultural systems, the plant microbiome—the microbial organisms associated with plant tissues—has been shown to have important effects on host physiology and ecology, yet we know little about how these plant-microbe relationships play out in urban environments. Here we characterize the composition of fungal communities associated with leaves of one of the most common sidewalk trees in the city of San Francisco, California. We focus our efforts on endophytic fungi (asymptomatic microfungi that live inside healthy leaves), which have been shown in other systems to have large ecological effects on the health of their plant hosts. Specifically, we characterized the foliar fungal microbiome of *Metrosideros excelsa* (Myrtaceae) trees growing in a variety of urban environmental conditions. We used high-throughput culturing, PCR, and Sanger sequencing of the ITS nrDNA region to quantify the composition and structure of fungal communities growing within healthy leaves of 30 *M. excelsa* trees from 6 distinct sites, which were selected to capture the range of environmental conditions found within city limits. Sequencing resulted in 854 high-quality ITS sequences. These sequences clustered into 85 Operational Taxonomic Units (97% VSEARCH OTUs). We found that these communities encompass relatively high alpha (within) and beta (between-site) diversity. Because the communities are all from the same host tree species, and located in relatively close geographical proximity to one another, these analyses suggest that urban environmental factors such as heat islands or differences in vegetation or traffic density (and associated air quality) may potentially be influencing the composition of these fungal communities. These biogeographic patterns provide evidence that plant microbiomes in urban environments can be as dynamic and complex as their natural counterparts. As human populations continue to transition out of rural areas and into cities, understanding the factors that shape environmental microbial communities in urban ecosystems stands to become increasingly important.

INTRODUCTION

Although major cities and urban centers only cover a small portion the Earth's total geographic area, more than 50% of the human population lives in these urban centers, and the impact that these cities have on the environment can be seen worldwide (Schneider, Friedl, and Potere 2009). With increasing urbanization, understanding the ecology of cities and urban settings has become critical to human health and well being. Despite their comparatively small geographic size, the high density of human populations in these environments makes them distinct ecosystems with their own unique dynamics (Sukopp 1998). Urban environments represent the convergence of humans from around the world, any plant or animal species those humans might have brought with them, and infrastructures such as roads, sewers, and tall buildings. Despite the complexity that these ecosystems present, they are often overlooked by ecologists because more traditional ecology has focused on 'natural' systems. In urban systems, however, human influence is a primary ecological factor (McDonnell and Niemelä 2011). In recent years, ecologists have begun studying the urban environment just as they would a natural environment, in order to understand the novel environmental conditions this setting presents to the organisms that live there (Wu 2014).

In this study, we focus on the urban ecology of trees in San Francisco, California. Urban trees can play a major role in shaping the ecosystem of a city. Just as the trees in a forest have a considerable impact on its climate and ecology, trees in cities can have notable effects on a city's environment. For example, plant life in cities can impact temperature, air quality, and other aspects of human health (Willis

48 and Petrokofsky 2017). The urban heat island effect, which occurs when ‘islands’ of heat form as heat
49 gets trapped between tall buildings, is one of the most well-documented unique urban anthropogenic
50 environmental conditions (Oke 1973). Trees in urban environments have been shown to interact with
51 these city-specific environmental factors (Kong et al. 2014). For example, trees in cities have been
52 shown to improve urban air quality by taking up significant amounts of carbon dioxide from city air
53 (Nowak et al. 2014). As pollution generated in urban centers is one of the major contributing factors to
54 worldwide pollution, trees in urban environments may have a role in managing the environmental impacts
55 of urbanization (Alberti et al. 2003).

56 Due to their importance, there is growing interest in understanding and maintaining sustainable urban
57 ecologies. One potentially major factor influencing health of plants both in nature and cities is the plant
58 microbiome. As the widespread availability of DNA sequencing has made it possible to characterize
59 microbial communities more easily and comprehensively, the microbiome has become a major area of
60 interest in numerous organisms (Kyrpides, Elie-Fadrosh, and Ivanova 2016). Just as the emerging field of
61 human microbiome study has revealed that symbiotic, non-pathogenic microbes can have major impacts
62 on human health (David et al. 2014), plants also host numerous symbiotic microbes. Similar to the
63 human microbiome, the plant microbiome contains great diversity and is comprised of multiple distinct
64 communities in various tissue systems such as the roots (rhizosphere), leaf surface (phyllosphere), and leaf
65 interior (endosphere) (Turner, James, and Poole 2013). All of these microbiomes can have an impact on
66 their host’s physiology. For instance, bacterial root microbes have been shown to play a role in the growth
67 of various plant species (Gaiero et al. 2013). They can also play a variety of roles in host physiology,
68 depending on their location within the plant (Schlaeppli and Bulgarelli 2015). These communities can
69 be quite dynamic, and vary with factors such as plant age (Cavaglieri, Orlando, and Etcheverry 2009).
70 While each of these sets of interactions is important, here we focus on the microbial ecology of fungal
71 microorganisms living in the leaf endosphere. The endosphere is an ideal system for biogeographic
72 studies, both because it is highly diverse, well replicated, and because it is a major interface between the
73 host plant and its environment (Meyer and Leveau 2012).

74 Endophytic microbes are naturally found in the interior of leaves, whether introduced by natural
75 wounds or openings in the leaf surface, or through penetrating the plant surface with hydrolytic enzymes
76 (Hallmann et al. 1997). Although some of these microbes may be latent pathogens or decomposers
77 waiting for the leaf to die, others are mutualists that may confer a benefit to their host (Carroll 1988).
78 In wild grasses, symbiotic fungi have been shown to protect their hosts by discouraging herbivory, and
79 can even affect host reproductive viability in those same systems (Clay 1988). In controlled settings,
80 inoculation experiments have shown that specific species of endophytes can have an impact on their
81 host’s overall health, including factors such as resistance and susceptibility to disease (Busby, Ridout,
82 and Newcombe 2016). In nature, the fungal microbial communities can have impacts on their host’s
83 physiology, such as limiting pathogen damage (Arnold et al. 2003).

84 In the wild, endophytic communities display species diversity comparable to that of any macroscopic
85 community, even among individual trees from the same species (Gazis, Rehner, and Chaverri, n.d.).
86 However, what factors influence this diversity across diverse environments remains an area of active
87 research. The biodiversity of these communities can be quite high, especially in areas like tropical forests
88 (Arnold and Lutzoni 2007). In such natural settings, plant-associated microbial community compositions
89 can show clear biogeographic structure (Andrews and Harris 2000). The urban setting, however, may
90 be distinct, because factors such as rainfall and elevation will likely be less apparent over a smaller
91 geographic area, but new factors such as proximity to roads and tall buildings may introduce effects of
92 their own. Studies of suburban forests in Japan have indicated that an urban setting has a notable impact
93 on endophytic diversity (Matsumura and Fukuda 2013). However, the full impact of urban environmental
94 factors on endophytic communities has yet to be completely understood.

95 In this study, we used culturing and barcode gene sequencing to identify the species makeup of endo-
96 phytic communities in the New Zealand Christmas Tree, *Metrosideros excelsa* (Myrtaceae), throughout
97 San Francisco, CA to quantify the diversity and biogeographic patterns of foliar fungal communities
98 in this urban environment. Based on studies in natural systems, we expected to find high community
99 diversity both within and between sites, as well as a degree of biogeographic structure to these community
100 compositions. We also anticipated that urban environmental factors may play a role in shaping the
101 biogeography in these endophytic communities. We found that leaves from trees sampled at different
102 sites across the city did indeed harbor distinct communities, and also that culture-based fungal isolation

103 frequencies differed among sites as well.

104 METHODS

105 Host Selection

106 We chose to focus on *Metrosideros excelsa* individuals. *Metrosideros excelsa* was an ideal species for
107 our study because it is widely planted throughout San Francisco and as such we were able to obtain
108 samples from a large number of trees in a variety of locations throughout the city. Although *M. excelsa*'s
109 endophytic communities have been characterized in its native home of New Zealand, there have been
110 few studies about these communities outside of its native environment or in an urban setting (McKenzie,
111 Buchanan, and Johnston 1999). In a related Hawaiian species, *Metrosideros polymorpha*, the species
112 makeup of fungal endophyte communities has been shown to vary greatly with environmental factors
113 such as elevation and rainfall (Zimmerman and Vitousek 2012).

114 Site Selection

115 We selected six sites across the city for leaf sample collection (Fig. 1). When selecting sites, we
116 took factors such as traffic levels, estimated temperature, proximity to tall buildings, elevation, and
117 proximity to the ocean and San Francisco Bay into account. In selecting the sites, we wanted to
118 capture the wide range of potential urban climates. We used street tree location data available from
119 the City of San Francisco (permalink: <https://data.sfgov.org/City-Infrastructure/Street-Tree-List/tkzw-k3nq>), which documents the location and species of every tree in San
120 Francisco, to choose unique locations around the city with sufficient densities of *Metrosideros excelsa*
121 individuals (Fig. 1).
122

123 Sample Collection

124 We collected small branches from 5 trees in each of these sites using a clipper pole, collecting at least 3
125 sun-exposed outer branches from each tree. Because *M. excelsa* is an evergreen tree and the newer leaves
126 are likely to contain fewer fungi, we controlled for leaf age by only collecting branches which contained
127 dark green leaves that appeared to be at least one year old. We collected all samples on the same day,
128 August 26, 2017, to ensure that daily weather patterns and seasonal effects would not have an impact on
129 the microbial community composition. Once collected, leaves were stored in labeled plastic bags and
130 stored at 4°C until culturing. All leaves were processed within 48 hours of collection. Permission to
131 sample leaves was granted by the San Francisco Bureau of Urban Forestry.

132 Culturing Methods

133 From each sample, we selected a subset of dark green asymptomatic leaves for culturing. We surface-
134 sterilized the leaves by first rinsing them with distilled water, then immersing them in a sterile petri dish
135 with 95% ethanol for 10 seconds, 10% NaOCl for 2 minutes, then 70% ethanol for another 2 minutes
136 (following (Arnold et al. 2007)), emptying the dish between rinses and leaving it closed until inside
137 a sterile biosafety cabinet after the last rinse. We then cut the leaves into small (2mm x 2mm) pieces
138 with flame-sterilized scissors and placed them into 1.5 mL microcentrifuge slant tubes filled with 2%
139 VWR-brand malt extract agar (MEA). For each tree, we prepared 6 leaves and made 100 tubes, except for
140 the trees from the downtown site. For the trees at this site, we prepared 140 tubes per tree because we
141 found that they had low isolation frequencies in a preliminary sampling bout. All leaves were prepared
142 this way within 48 hours of the initial leaf sampling, to prevent death of the leaf tissue from altering the
143 fungal community composition.

144 After two weeks, we evaluated the tubes for fungal growth and subcultured the emergent fungi from
145 tubes with growth onto 35mm plates with 2% MEA in order to better evaluate their morphotypes and
146 accumulate sufficient tissue for future barcode gene sequencing and water voucher preparation. We
147 re-evaluated and subcultured these tubes in the following months to capture any late-growing fungi. Each
148 pure culture was vouchered in sterile dI water and stored in a living culture collection in the Zimmerman
149 Lab at the University of San Francisco.

150 Molecular Methods

151 We extracted DNA from fungal mycelium using the Sigma RED Extract 'n Amp DNA extraction kit and
152 following previously published protocols (U'Ren et al. 2012). First, we added fungal tissue to sterile

tubes filled with 1 mm zirconium oxide beads, then added 100 μ L of Extract 'n' Amp DNA extraction solution. Next, we put the tubes in a bead-beater (Mini-Beadbeater-96, Bio Spec Products Inc.) for one minute. The samples were then placed on a heat block at 95°C for 10 minutes. After the heating step, we added a dilution buffer to each tube and stored them at 4°C until PCR.

We performed PCR using fungal-specific primers for the Internal Transcribed Spacer region, a commonly-accepted fungal barcode locus (Schoch et al. 2012), using the ITS1F forward primer (5'-CTT GGT CAT TTA GAG GAA GTA A-3') and ITS4 reverse primer (5'-TCC TCC GCT TAT TGA TAT GC-3'). For each PCR reaction, we used 1 μ L of template DNA, 10 μ L Extract 'n' Amp Taq polymerase, 6.4 μ L PCR-grade water, 1 μ L bovine serum albumin, 0.8 μ L ITS1F forward primer, and 0.8 μ L ITS4 reverse primer (U'Ren et al. 2012). For the PCR reaction, we used a BioRAD T100 thermal cycler with the following cycles: 95°C for 3 minutes; 35 cycles of 95°C for 30 seconds, 54°C for 30 seconds, 72°C for 30 seconds; then 72°C for 10 minutes. To ensure that the fungal DNA successfully amplified, and that the master mix was not contaminated, we ran 5 μ L of each sample and a negative PCR control on a 1% agarose gel with 1X Tris-acetate-EDTA (TEA) buffer and SYBR Safe. We ran the gel at 120 volts for 20 minutes and visualized bands using UV transillumination. Successful PCR samples with clean negative controls were kept at 4°C until sequencing preparation.

To prepare successfully amplified samples for Sanger sequencing, we first cleaned each sample with 1 μ L Thermo Fisher Shrimp Alkaline Phosphatase Exonuclease (ExoSAP-IT). To clean the samples, we used the following cycle on a BioRAD T100 thermal cycler: 37°C for 15 minutes, 80°C for 15 minutes, then hold at 4°C. After cleaning, samples were kept at 4°C until they were ready to be sent for sequencing. Directly before being sent for sequencing, cleaned samples that showed bright bands on their gels were diluted with an additional 15 μ L PCR water, although a small number of samples that had faint bands on the gels were not diluted. Cleaned samples were sent to MCLabs (South San Francisco, CA) for Sanger sequencing.

Computational and Statistical Methods

We used Geneious 11.1 to manually clean and trim the Sanger sequencing data, and to identify and remove failed and low-quality sequences (Kearse et al. 2012). Sequences that appeared to have multiple strong signals at a given position were discarded because they may have originated from from a mixed culture. Usable sequences were cleaned by trimming the ends of low-quality nucleotides, clarifying any ambiguity codes, and resolving incorrect base assignment due to dye blobs. Cleaned sequences are available on GenBank <>.

After processing the cleaned sequences to add relevant metadata, we used mothur version 1.39.5 to determine Operational Taxonomic Units (OTUs), based on 97% ITS sequence similarity and abundance-based greedy clustering (Schloss et al. 2009). Next, we used R 4.1.1 (R Core Team 2021) to analyze the resulting OTU table. We used vegan (Oksanen et al. 2018) to calculate species accumulation curves to determine species richness from the OTU data, create a Non-Metric Multidimensional Scaling (NMDS) ordination, and perform a PERMANOVA to assess differences in communities between sites (Anderson 2001). We used the Tree-Based Alignment Selector (TBAS) toolkit to assign taxonomic context to each ITS sequence (Carbone et al. 2017). This toolkit matches unknown ITS sequences to the most similar ITS sequences in a large multi-gene phylogeny of confidently-assigned taxa. We used the simple features (Pebesma 2018) and ggmap (Kahle and Wickham 2013) packages to produce maps of study sites.

Significant differences in fungal leaf isolation frequency and tree diameter were determined using a nonparametric Kruskal–Wallis one-way test of variance. All analyses, figures, and the final manuscript were generated using a makefile, bash scripts, and R and Rmarkdown. All relevant metadata, as well as analysis and manuscript code, are available on GitHub: <https://github.com/ZimmermanLab/SF-metrosideros-endophytes/> and archived at <>. The complete computational environment for the analyses is documented in a Dockerfile based on rocker containers (Boettiger 2014) and renv renv.lock (Ushey 2021) in that same repository.

Methods for Comparing Fungi in San Francisco Trees to New Zealand Trees

In order to compare the foliar microbial populations found in *M. excelsa* throughout San Francisco to those in the tree's native range (New Zealand), we surveyed public GenBank records and performed a literature search. First, we searched Google Scholar for papers mentioning “*Metrosideros excelsa* fungal endophytes” and then, more generally, “foliar fungi in *Metrosideros excelsa* leaves.” After this search, we looked through the papers cited by those we originally found in order to find any additional studies.

207 Then, we searched GenBank for fungal ITS sequences that had *M. excelsa* listed as their host organism.
208 We performed this search using the parameters “host=“*Metrosideros excelsa*”, organism=fungi”, in order
209 to limit our results to fungal sequences. This search yielded several new sequences that weren’t included
210 in the previous search because they were unpublished. Using both of these methods, we were able to find
211 about 30 taxa to include as a point of comparison, although many of these were described as pathogenic
212 instead of endophytic.

213 RESULTS

214 Overall, we found high diversity in these urban endophytic communities. From 30 different host trees,
215 we successfully cultured and sequenced 854 fungal isolates, which made up 85 distinct 97% ITS OTUs.
216 Some communities encompassed a low number of fungal isolates; species richness analyses showed that
217 in nearly all of the trees analyzed, our sampling did not encompass the complete microbial diversity (Fig.
218 3). While we documented species diversity within some sites that exceeds 20 distinct fungal OTUs, there
219 appears to be greater species diversity in these fungal endophytic communities than this study was able to
220 document.

221 Isolation Frequency

222 The isolation frequency, or the percentage of leaf pieces that yielded fungal isolates, varied significantly
223 between sites (Kruskal Wallis $p < 0.01$, Fig. 2A). In most sites, the isolation frequency also varied
224 between trees, especially in the Bay site. Trees within the Bay site also show the greatest variation in
225 diameter at breast height (DBH) (Fig. 2B). The only site that does not show as consistent variation in
226 isolation frequency is the Downtown site, which is also the site with the lowest isolation frequencies
227 overall.

228 Diversity Patterns

229 We found 85 total 97% nrITS OTUs among the 30 different trees. Both isolation frequency and number
230 of fungal species found varied notably between trees. Species accumulation curves showed that the total
231 amount of fungal diversity was not completely sampled in any of the sites (Fig. 3).

232 The most abundant fungal class that we found in our samples was Dothideomycetes, followed by
233 Sordariomycetes; each had over 300 sequences assigned to them (Table 1). Dothideomycetes was the
234 predominant class in most sites except for the Downtown and Bay sites (Fig. 4). In both of these sites,
235 Sordariomycetes was the most common class instead. There are several classes that are either absent
236 or present in small numbers in most sites, but more abundant in one or several sites. For example,
237 Eurotiomycetes are more common in the Downtown and Freeway sites, and Leotiomycetes are only
238 abundant in the Mt. Davidson site. Only two sequences were not identified by T-BAS. We queried these
239 two sequences against the NCBI nt database via BLASTn and found them to match with high confidence
240 to the Pezizomycetes, based on similarity to a strain recently identified in Australia (Catcheside, Qaraghuli,
241 and Catcheside 2017).

242 Biogeographic Patterns

243 While the NMDS ordination group ellipses (standard error around the centroid for each site) suggest that
244 there are differences in dispersion among sites, the multivariate dispersion of these communities was not
245 significantly different ($p = 0.76$; Fig. 5). Sites that were closer together geographically tended to also be
246 closer to one another in ordination space.

247 DISCUSSION

248 In this study, we sought to characterize the foliar microbiome of *Metrosideros excelsa* trees across the
249 city of San Francisco. We found that the microbial composition of these urban trees’ leaves varied in
250 many aspects, from number of fungal isolates to the identities of said isolates. While we did not quantify
251 environmental variation in this study, the community variation we observed could be explained by either
252 environmental or host physiological factors.

253 Isolation Frequency and Tree size

254 Both isolation frequency and DBH varied greatly from one site to the other, and some sites show
255 considerable within-site differences in these factors as well. The trees in the Bay site show the greatest

range in both isolation frequency and DBH (Fig. 2), indicating that the size of a tree may have a correlation with the number of fungal endophytes found within its leaves. The Mt. Davidson trees have a larger median and range of DBH than the ocean or freeway trees (Fig. 2B), and also have a higher median isolation frequency, which could indicate that trees with a larger DBH may also have a higher number of fungal endophytes.

Just as it shows the greatest range of isolation frequencies and tree sizes, the Bay site also has some of the least similarity between its fungal communities on the NMDS ordination, and the two trees with vastly different communities are also the smallest (Fig. 5). In this instance, it appears that the size of the trees has the largest impact on the endophytic communities. Furthermore, the trees in the Mt. Davidson site, which have some of the most similar communities (Fig. 5), also have fairly large trees with similar DBH. Additionally, it is notable that trees with a wider trunk are presumably older than those with thinner trunks, and therefore trees with a larger DBH are likely older and have had more time to accumulate fungal communities. The fact that these trees have similar sizes in addition to similar community composition indicates that a tree's size, and by extension its age, may have an impact on its endophytic communities.

Although there appears to be a general pattern with larger trees hosting a greater number of fungal endophytes, DBH cannot explain all of the variation in isolation frequencies, as demonstrated by the Downtown and Balboa sites. Although the trees from the downtown site have a higher median DBH (Fig. 2B), the trees from the Balboa site have considerably more endophytes (Fig. 2A), which suggests that larger trees do not necessarily have a higher number of fungal endophytes.

Diversity Patterns

Endophytic microbiomes in tree leaves are known to be highly diverse in natural systems, and that also appears to be the case in these urban trees. The species accumulation curves indicate that even in the most well-represented sites, there are still many potentially undiscovered OTUs within these endophytic communities (Fig. 3). Additionally, there is a considerable amount of taxonomic variation between certain sites, in addition to the generally high diversity of each site (Fig. 4). In such cases, it is likely that environmental factors play a role in shaping the endophytic communities of these trees. While the impact of environmental factors may be less evident when focusing on fungal endophyte community richness, it becomes more apparent when looking at the identities of these endophytes. The composition of these communities can vary greatly among trees with similar isolation frequencies, as demonstrated by the taxonomic composition of the Mt. Davidson and Bay sites (Fig. 4). In general, there was higher compositional similarity between trees from the same location than between trees of a similar size (Fig. 4). (fig:nmds-plot) and Table (tab:permanova-table)).

Biogeographic Patterns

There appears to be a degree of biogeographic structure to the endophytic communities within these urban trees. Communities of trees from the same site generally cluster closer together than trees from different sites (Fig. 5). Although the Balboa site appears to have a high isolation frequency (Fig. 2A), many of the slant tubes that showed fungal growth failed to grow into a larger culture, leading to a lower number of usable sequences (Fig. 3). This low number of sequences per tree may indicate that the divergence in community composition that appears to be present might be due to the low sample size, rather than actual difference between the microbial communities between the Balboa trees (Fig. 5).

However, even sites with low within-site diversity appear to be more similar to other trees from geographically close sites. For example, the Downtown trees do not cluster with the Mt. Davidson, Balboa, or Ocean trees in the ordination, indicating that they have fairly different fungal communities (Fig. 5). Generally, the Westernmost sites (Ocean and Balboa) cluster to one side of the ordination, while the Easternmost sites (Bay and Downtown) cluster to the other, and the sites closer to the middle of the city (Mt. Davidson and Freeway) cluster between them (Fig. 5). Notably, the Downtown and Freeway sites cluster fairly close together (Fig. 5) despite being fairly distant geographically (Fig. 1). In this case, it is possible that there are environmental factors that cause the communities to be more compositionally similar. One of the most likely environmental similarities between these two sites is that they are both exposed to high traffic levels, with one site located in a bustling downtown and the other located near two large freeways. Carbon dioxide emissions, particulates, or both of these pollutants could possibly be having an effect on both these trees and their microbiomes. Endophytic bacteria have been shown to have a phytoremediating effect (Afzal, Khan, and Sessitsch 2014), so it is possible that endophytic fungi may be playing a similar role in helping these plants persist in high-traffic and thus lower air quality conditions.

Community composition compared to related trees in other locations

In widely-distributed plant species, fungal endophyte composition can differ quite considerably in a species from its native home to the places it has been introduced (Taylor, Hyde, and Jones 1999). *M. excelsa* is native to New Zealand, but it and its close relatives in the *Myrtaceae* family are widespread across the world. Across all locations, the most abundant classes found within these *M. excelsa* fungal communities were Dothideomycetes and Sordariomycetes by a notable margin (Table 1). At broad taxonomic levels such as Class, the endophytes found in this study appear to be similar to those found in other *Myrtaceae* trees elsewhere. For instance, a study of *Myrtaceae* trees in South America, the first and second most common fungal isolates were also identified as Dothideomycetes followed by Sordariomycetes (Vaz et al. 2014). Dothideomycetes and Sordariomycetes are also the first and second most abundant classes in the endophytic communities of *M. excelsa*'s close relative in Hawaii, *Metrosideros polymorpha*, as well (Zimmerman and Vitousek 2012). This could indicate that these fungal classes are well-adapted to living as endophytes within these trees, but it is also possible that these communities are actually quite different when considered at a finer taxonomic resolution.

Comparison of trees in San Francisco and New Zealand

When comparing *M. excelsa*'s endophytic taxa found in the literature to those that we collected, we found that the taxa between the regions were somewhat similar on an order level, and even more so on a class level. Only one of the orders that was extremely common in New Zealand (Mycosphaerales) was absent in San Francisco, and only two of the most common orders in San Francisco (Capnodiales and Pezizales) were absent in New Zealand. However, most of the families that are found in New Zealand are absent in San Francisco, and vice versa. We only found four families in common: Mycosphaerellaceae, Glomerellaceae, Nectriaceae, and Helotiaceae. Of these, Mycosphaerellaceae is the only family that is present in New Zealand and was very prevalent (over 50 isolates) in San Francisco. The other families that they had in common were only present in very small numbers in San Francisco, and we were unable to find documentation of three most prevalent families in San Francisco (Xylariaceae, Botryosphaeriaceae, and Cladosporiaceae) being isolated from leaves in New Zealand. Although this could be due to the small sample size, it suggests that *M. excelsa*'s endophytic populations may be influenced primarily by their environment. However, additional research into both the endophytic populations of other trees in San Francisco and *M. excelsa* in its native range would be required to draw further conclusions.

It is important to note the difference in methods between how we collected our samples and how others studying *M. excelsa* in New Zealand collected theirs. Whereas we took care to only collect samples from unfallen, asymptomatic leaves, most of the papers referenced in this comparison use either leaf litter, dead leaves, or obviously symptomatic leaves. Therefore, it is reasonable to infer that the data we are comparing our results to is biased towards pathogenic and decomposing fungi, and underrepresents the non-harmful endophytic populations in *M. excelsa*'s native range. Additionally, the number of fungal isolates we were able to find using literature and GenBank searches was much lower than the number of samples we collected ourselves, making quantitative comparison unfeasible. However, we think that such comparisons are still worthwhile as a preliminary look into the extent that *M. excelsa* in San Francisco may maintain its native endophytic microbiome.

CONCLUSIONS

These findings indicate that the endophytic microbiomes of urban trees are complex and diverse, and may show a degree of biogeographic structure that reflects their natural counterparts. Additionally, it is likely that urban environmental factors play a considerable role in shaping the endophytic communities of these trees. This study has demonstrated that the urban endophytic microbiome is highly diverse and has the potential to be structured by urban environmental factors. Our species accumulation curves indicate that the full diversity of these endophytic communities has yet to be sampled (Fig. 3). Even in the small geographic area of San Francisco, we found notable trends in microbiome composition that may vary with geographic factors, host physical factors such as DBH and age, and uniquely urban environmental factors, such as traffic. A combination of environmental factors and perhaps host physiology therefore appear to be the driving force behind the diversity of these microbiomes. While it is difficult to determine, given the data we have, the exact mechanisms that influencing the composition of these communities, the amount of species diversity and biogeographic structure we present here indicate that the foliar microbiomes of

urban trees may be just as complex and dynamic as those of trees in nature, and are worth of further concerted study.

ACKNOWLEDGEMENTS

This study was conducted as an undergraduate Biology honors thesis project by EG under the guidance of NZ. Joshua Copeland and Julian Murdzek assisted with culture processing, vouchering, and DNA extractions. Jeff Oda contributed supplies and equipment expertise.

REFERENCES

- Afzal, Muhammad, Qaiser M Khan, and Angela Sessitsch. 2014. "Endophytic Bacteria: Prospects and Applications for the Phytoremediation of Organic Pollutants." *Chemosphere* 117: 232–42.
- Alberti, Marina, John M Marzluff, Eric Shulenberger, Gordon Bradley, Clare Ryan, and Craig Zumbrunnen. 2003. "Integrating Humans into Ecology: Opportunities and Challenges for Studying Urban Ecosystems." *AIBS Bulletin* 53 (12): 1169–79.
- Anderson, Marti J. 2001. "A New Method for Non-Parametric Multivariate Analysis of Variance." *Austral Ecology* 26 (1): 32–46.
- Andrews, John H, and Robin F Harris. 2000. "The Ecology and Biogeography of Microorganisms on Plant Surfaces." *Annual Review of Phytopathology* 38 (1): 145–80.
- Arnold, A Elizabeth, Daniel A Henk, Rebecca L Eells, Francois Lutzoni, and Rytas Vilgalys. 2007. "Diversity and Phylogenetic Affinities of Foliar Fungal Endophytes in Loblolly Pine Inferred by Culturing and Environmental PCR." *Mycologia* 99 (2): 185–206.
- Arnold, A Elizabeth, and Francois Lutzoni. 2007. "Diversity and Host Range of Foliar Fungal Endophytes: Are Tropical Leaves Biodiversity Hotspots?" *Ecology* 88 (3): 541–49.
- Arnold, A Elizabeth, Luis Carlos Mejia, Damond Kylo, Enith I Rojas, Zuleyka Maynard, Nancy Robbins, and Edward Allen Herre. 2003. "Fungal Endophytes Limit Pathogen Damage in a Tropical Tree." *Proceedings of the National Academy of Sciences* 100 (26): 15649–54.
- Boettiger, Carl. 2014. "An Introduction to Docker for Reproducible Research, with Examples from the r Environment." *arXiv(2015) ACM SIGOPS Operating Systems Review, Special Issue on Repeatability and Sharing of Experimental Artifacts*. 49(1), 71-79, 1410.0846v1. <http://arxiv.org/abs/1410.0846v1> <http://arxiv.org/pdf/1410.0846v1>.
- Busby, Posy E., Mary Ridout, and George Newcombe. 2016. "Fungal Endophytes: Modifiers of Plant Disease." *Plant Molecular Biology* 90 (6): 645–55. <https://doi.org/10.1007/s11103-015-0412-0>.
- Carbone, Ignazio, James B. White, Jolanta Miadlikowska, A. Elizabeth Arnold, Mark A. Miller, Frank Kauff, Jana M. U'Ren, Georgiana May, and François Lutzoni. 2017. "T-BAS: Tree-Based Alignment Selector Toolkit for Phylogenetic-Based Placement, Alignment Downloads and Metadata Visualization: An Example with the Pezizomycotina Tree of Life." *Bioinformatics* 33 (8): 1160–68. <https://doi.org/10.1093/bioinformatics/btw808>.
- Carroll, George. 1988. "Fungal Endophytes in Stems and Leaves: From Latent Pathogen to Mutualistic Symbiont." *Ecology* 69 (1): 2–9.
- Catcheside, Pamela S, Samra Qaraghuli, and David EA Catcheside. 2017. "A New Species of Small Black Disc Fungi, Smardaea Australis (Pezizales, Pyronemataceae), Is Described from Australia." *Swainsona* 31: 17–101.
- Cavaglieri, Lilia, Julieta Orlando, and Miriam Etcheverry. 2009. "Rhizosphere Microbial Community Structure at Different Maize Plant Growth Stages and Root Locations." *Microbiological Research* 164 (4): 391–99.
- Clay, Keith. 1988. "Fungal Endophytes of Grasses: A Defensive Mutualism Between Plants and Fungi." *Ecology* 69 (1): 10–16.
- David, Lawrence A, Corinne F Maurice, Rachel N Carmody, David B Gootenberg, Julie E Button, Benjamin E Wolfe, Alisha V Ling, et al. 2014. "Diet Rapidly and Reproducibly Alters the Human Gut Microbiome." *Nature* 505 (7484): 559.
- Gaiero, Jonathan R, Crystal A McCall, Karen A Thompson, Nicola J Day, Anna S Best, and Kari E Dunfield. 2013. "Inside the Root Microbiome: Bacterial Root Endophytes and Plant Growth Promotion." *American Journal of Botany* 100 (9): 1738–50.

- Gazis, Romina, Stephen Rehner, and Priscila Chaverri. n.d. "Species Delimitation in Fungal Endophyte Diversity Studies and Its Implications in Ecological and Biogeographic Inferences." *Molecular Ecology* 20 (14): 3001–13. <https://doi.org/10.1111/j.1365-294X.2011.05110.x>.
- Hallmann, J, A Quadt-Hallmann, WF Mahaffee, and JW Kloepper. 1997. "Bacterial Endophytes in Agricultural Crops." *Canadian Journal of Microbiology* 43 (10): 895–914.
- Kahle, David, and Hadley Wickham. 2013. "Ggmap: Spatial Visualization with Ggplot2." *The R Journal* 5 (1): 144–61. <https://journal.r-project.org/archive/2013-1/kahle-wickham.pdf>.
- Kearse, Matthew, Richard Moir, Amy Wilson, Steven Stones-Havas, Matthew Cheung, Shane Sturrock, Simon Buxton, et al. 2012. "Geneious Basic: An Integrated and Extendable Desktop Software Platform for the Organization and Analysis of Sequence Data." *Bioinformatics* 28 (12): 1647–49. <https://doi.org/10.1093/bioinformatics/bts199>.
- Kong, F, H Yin, P James, LR Hutyra, and HS He. 2014. "Effects of Spatial Pattern of Greenspace on Urban Cooling in a Large Metropolitan Area of Eastern China." *Landscape and Urban Planning* 128: 35–47. <https://doi.org/10.1016/j.landurbplan.2014.04.018>.
- Kyrpides, Nikos C, Emiley A Eloë-Fadrosh, and Natalia N Ivanova. 2016. "Microbiome Data Science: Understanding Our Microbial Planet." *Trends in Microbiology* 24 (6): 425–27.
- Matsumura, Emi, and Kenji Fukuda. 2013. "A Comparison of Fungal Endophytic Community Diversity in Tree Leaves of Rural and Urban Temperate Forests of Kanto District, Eastern Japan." *Fungal Biology* 117 (3): 191–201. <https://doi.org/https://doi.org/10.1016/j.funbio.2013.01.007>.
- McDonnell, Mark J, and J Niemelä. 2011. "The History of Urban Ecology." *Urban Ecology*, 9.
- McKenzie, E. H. C., P. K. Buchanan, and P. R. Johnston. 1999. "Fungi on Pohutukawa and Other *Metrosideros* Species in New Zealand." *New Zealand Journal of Botany* 37 (2): 335–54. <https://doi.org/10.1080/0028825X.1999.9512637>.
- Meyer, Katrin M, and Johan HJ Leveau. 2012. "Microbiology of the Phyllosphere: A Playground for Testing Ecological Concepts." *Oecologia* 168 (3): 621–29.
- Nowak, David J., Satoshi Hirabayashi, Allison Bodine, and Eric Greenfield. 2014. "Tree and Forest Effects on Air Quality and Human Health in the United States." *Environmental Pollution* 193: 119–29. <https://doi.org/https://doi.org/10.1016/j.envpol.2014.05.028>.
- Oke, Tim R. 1973. "City Size and the Urban Heat Island." *Atmospheric Environment (1967)* 7 (8): 769–79.
- Oksanen, Jari, F. Guillaume Blanchet, Michael Friendly, Roeland Kindt, Pierre Legendre, Dan McGlinn, Peter R. Minchin, et al. 2018. *Vegan: Community Ecology Package*. <https://CRAN.R-project.org/package=vegan>.
- Pebesma, Edzer. 2018. "Simple Features for R: Standardized Support for Spatial Vector Data." *The R Journal* 10 (1): 439–46. <https://doi.org/10.32614/RJ-2018-009>.
- R Core Team. 2021. *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing. <https://www.R-project.org/>.
- Schlaeppli, Klaus, and Davide Bulgarelli. 2015. "The Plant Microbiome at Work." *Molecular Plant-Microbe Interactions* 28 (3): 212–17.
- Schloss, Patrick D, Sarah L Westcott, Thomas Ryabin, Justine R Hall, Martin Hartmann, Emily B Hollister, Ryan A Lesniewski, et al. 2009. "Introducing Mothur: Open-Source, Platform-Independent, Community-Supported Software for Describing and Comparing Microbial Communities." *Applied and Environmental Microbiology* 75 (23): 7537–41.
- Schneider, Annemarie, Mark A Friedl, and David Potere. 2009. "A New Map of Global Urban Extent from MODIS Satellite Data." *Environmental Research Letters* 4 (4): 044003.
- Schoch, Conrad L, Keith A Seifert, Sabine Huhndorf, Vincent Robert, John L Spouge, C Andre Levesque, Wen Chen, et al. 2012. "Nuclear Ribosomal Internal Transcribed Spacer (ITS) Region as a Universal DNA Barcode Marker for Fungi." *Proceedings of the National Academy of Sciences* 109 (16): 6241–46.
- Sukopp, Herbert. 1998. "Urban Ecology—Scientific and Practical Aspects." In *Urban Ecology*, 3–16. Springer.
- Taylor, J. E., K. D. Hyde, and E. B. G. Jones. 1999. "Endophytic Fungi Associated with the Temperate Palm, *Trachycarpus Fortunei*, Within and Outside Its Natural Geographic Range." *New Phytologist* 142 (2): 335–46.

469 Turner, Thomas R, Euan K James, and Philip S Poole. 2013. "The Plant Microbiome." *Genome Biology*
470 14 (6): 209.

471 U'Ren, Jana M, F Lutzoni, J Miadlikowska, A D Laetsch, and A Elizabeth Arnold. 2012. "Host and
472 Geographic Structure of Endophytic and Endolichenic Fungi at a Continental Scale." *American*
473 *Journal of Botany* 99 (5): 898–914.

474 Ushey, Kevin. 2021. *Renv: Project Environments*. <https://rstudio.github.io/renv/>.

475 Vaz, Aline BM, Sonia Fontenla, Fernando S Rocha, Luciana R Brandao, Mariana LA Vieira, Virginia De
476 Garcia, Aristoteles Goes-Neto, and Carlos A Rosa. 2014. "Fungal Endophyte β -Diversity Associated
477 with Myrtaceae Species in an Andean Patagonian Forest (Argentina) and an Atlantic Forest (Brazil)." *Fungal Ecology* 8: 28–36.

478 Willis, Katherine J., and Gillian Petrokofsky. 2017. "The Natural Capital of City Trees." *Science* 356
479 (6336): 374–76. <https://doi.org/10.1126/science.aam9724>.

480 Wu, Jianguo. 2014. "Urban Ecology and Sustainability: The State-of-the-Science and Future Directions." *Landscape and Urban Planning* 125: 209–21.

481 Zimmerman, Naupaka B., and Peter M. Vitousek. 2012. "Fungal Endophyte Communities Reflect
482 Environmental Structuring Across a Hawaiian Landscape." *Proceedings of the National Academy of*
483 *Sciences* 109 (32): 13022–27. <https://doi.org/10.1073/pnas.1209872109>.

Table 1. Overall sequenced fungal isolate counts by Class and Order. All fungi successfully sequenced in this study were placed within the Ascomycota.

Class	Order	Number of sequences
Dothideomycetes	Capnodiales	266
Dothideomycetes	Botryosphaeriales	127
Dothideomycetes	Pleosporales	52
Dothideomycetes	unclassified,Capnodiales	3
Dothideomycetes	Trypetheliales	1
Dothideomycetes	unclassified	1
Eurotiomycetes	Eurotiales	57
Leotiomycetes	Helotiales	19
Pezizomycetes	Pezizales	54
Sordariomycetes	Xylariales	237
Sordariomycetes	Diaporthales	49
Sordariomycetes	Coniochaetales	38
Sordariomycetes	Sordariales	9
Sordariomycetes	Hypocreales	7
Sordariomycetes	Glomerellales	1

Table 2. PERMANOVA model output based on Bray-Curtis distance, with permutations stratified by site. Foliar fungal community composition in leaves of *Metrosideros excelsa* across San Francisco is significantly related to both site of sampling and tree diameter at breast height (DBH).

Term	df	Sum of Squares	Mean of Squares	F	R ²	p-value
trees\$Site.ID	5	3.642	0.728	3.015	0.373	0.022
trees\$DBH.cm	1	0.556	0.556	2.302	0.057	0.022
Residuals	23	5.557	0.242	NA	0.570	NA
Total	29	9.755	NA	NA	1.000	NA




Figure 1. A map of the locations sampled across the city of San Francisco, California, USA. Five trees from each site were sampled. Sites were selected to span the range of environmental conditions across the city. Map data © 2021 Google.




Figure 2. Isolation frequencies (A) and tree diameters (B) at each site. Isolation frequency (A) is a measure of how many slant tubes showed signs of fungal growth, out of how many total slant tubes were made. Approximately 100 slant tubes were made for each tree except for the trees in the downtown site, which had 140 slant tubes per tree because they had low isolation frequencies during the initial sampling. Sites along x axis are arranged left to right in order of their geographic location from west (left) to east (right).




Figure 3. Species accumulation curves showing fungal community OTU richness. Each line in A represents the OTU richness for a single host tree. Each line in B represents the combined OTU richness of all trees ($n = 5$) in one site.




Figure 4. Normalized relative abundances of taxonomic groupings (Classes in A and Orders in B) in each site. Sites along x axis are arranged left to right in order of their geographic location from west (left) to east (right).

NMDS ordination of fungal community composition




Figure 5. NMDS ordination of community compositions. Each point represents the endophytic community of one tree, and the size of the point corresponds to that tree's DBH, while the color of said point corresponds to the site that tree is from. Points that are closer together indicate that the trees they represent have similar community compositions. The ellipses show the standard error around the centroid of all points within a site, and are also color-coded according to which site they represent.