



IDX G9 Chemistry S+
Study Guide Issue Semester 1 Monthly 2
By Allison and Desheng, Edited by Jillian

NOTE: This is an official document by Indexademics. Unless otherwise stated, this document may not be accredited to individuals or groups other than the club IDX, nor should this document be distributed, sold, or modified for personal use in any way.

Contents:

Chapter 6: Electronic Structure of Atoms

6.1 Wave Nature of Light

- Light is electromagnetic radiation (EMR): oscillating electric and magnetic fields.
- Important wave properties:
 - **Wavelength (λ)** – distance between peaks (units: m, nm).
 - **Frequency (ν)** – number of waves per second (units: $\text{s}^{-1} = \text{Hz}$).
- All EM waves in vacuum travel at: **Speed of light: $c = 3.00 \times 10^8 \text{ m/s}$.**
- Relationship between λ and ν : **$c = \lambda\nu$.**
- **Visible Spectrum(from shortest λ / highest energy to longest λ / lowest energy): Violet \rightarrow Indigo \rightarrow Blue \rightarrow Green \rightarrow Yellow \rightarrow Orange \rightarrow Red.**
 - So: Wavelength: $\lambda(\text{violet}) < \dots < \lambda(\text{red})$; Frequency & energy: $\nu(\text{violet}) > \dots > \nu(\text{red})$.
- **Typical Question Types**
 - Given λ , find ν (or vice versa) using $c = \lambda\nu$.
 - Compare colors or regions of EM spectrum in terms of λ , ν , and energy.
 - Conceptual questions about how changing λ affects energy.

- **Common Traps**
 - **Forgetting to convert nm \rightarrow m (Very Important!)** when using formulas.
 - Confusing brightness with energy per photon (**energy per photon depends on frequency**, not brightness).

6.2 Quantized Energy and Photons

- **Planck: Energy is quantized** — it comes in packets called quanta.
 - **Planck's constant: $h = 6.626 \times 10^{-34} \text{ J}\cdot\text{s}$.**
- The energy of one photon: **$E = h\nu$.**
- **Photoelectric effect:**
 - Light shining on a metal can eject electrons from the surface.
 - **Ejection only happens if photon energy \geq threshold energy of the metal (frequency must be \geq threshold frequency ν_0).**
 - If you **increase intensity (brightness) at the same frequency**: more photons \rightarrow **more electrons ejected**, but same KE for each electron.
 - If you increase frequency (above ν_0): each photon has more energy \rightarrow electrons have higher KE.
 - **Remember that brightness doesn't change the frequency!**
- **Important Formulas:**
 - Photon energy: **$E_{\text{photon}} = h\nu = hc/\lambda$.**
 - Threshold energy: **$E_{\text{threshold}} = h\nu_0$.**
 - Kinetic energy of an ejected electron: **$KE = E_{\text{photon}} - E_{\text{threshold}} = h\nu - h\nu_0$.**
- **Typical Question Types**
 - Calculate the energy of a photon given ν or λ .
 - Find the total energy of N photons.
 - Given λ of light and the metal's threshold frequency ν_0 , find the KE of ejected electrons or decide if electrons are ejected.

- **Common Traps**

- Using $E = h\nu_0$ when the question is asking for KE. Remember: $KE = h(\nu - \nu_0)$.
- Thinking very intense low-frequency light can eject electrons. If $\nu < \nu_0$, no electrons are ejected, **no matter how bright**.

6.3 Line Spectra and the Bohr Model

- Continuous vs Line Spectra:

- **Continuous spectrum:** contains all wavelengths in a range (like white light through a prism \rightarrow smooth rainbow).
- **Atomic emission spectrum:** only specific discrete lines (certain wavelengths/frequencies), each element has a unique line spectrum.

- **Bohr Model of the Hydrogen Atom:**

- Electrons occupy specific energy levels (orbits) labeled by $n = 1, 2, 3, \dots$
- $n = 1$: lowest energy level \rightarrow ground state.
- $n \geq 2$: higher energy levels \rightarrow excited states.
- When electrons absorb energy, they jump from lower n to higher n (excited state).
- When they fall back to lower n , they emit photons whose energy equals the energy gap between the levels.
 - Any state $n > 3$ that goes back to $n = 3$ level exhibit infrared rays
 - Any state $n > 2$ that goes back to $n = 2$ exhibit visible light
 - Any state $n > 1$ that goes back to $n = 1$ (ground state) exhibit UV light

- Energy change in a transition: $\Delta E = E_{\text{higher}} - E_{\text{lower}} = h\nu$.
- Because only specific n levels exist, only specific energy differences are allowed \rightarrow only specific frequencies (lines) appear.

- **Strengths and Limits of Bohr Model**
 - **Strengths:** explains line spectrum of hydrogen and other one-electron ions (He^+ , Li^{2+} , etc.); introduces quantized energy levels.
 - **Limits:** doesn't work well for multi-electron atoms; electrons are not actually moving in simple circular orbits; cannot fully explain chemical behavior trends.
- **Typical Question Types**
 - Explain why hydrogen shows a line spectrum instead of a continuous one.
 - Define ground state and excited state in terms of energy levels n .
 - Relate a spectral line to an electron transition (higher $n \rightarrow$ lower n).
- **Common Traps**
 - Mixing up absorption (electron moves up in n) and emission (electron falls down in n).
 - Forgetting that a specific line = a specific pair of energy levels.

6.4 The Wave Behavior of Matter

- **de Broglie Matter Waves:**
 - Particles of matter have **wave properties**.
 - Wavelength of a particle: $\lambda = h/(mv)$.
 - **For large objects** (like a baseball or a car), mass is huge $\rightarrow \lambda$ is extremely small \rightarrow **wave behavior is negligible**.
 - **For very small particles** (like electrons), mass is tiny $\rightarrow \lambda$ can be comparable to atomic dimensions \rightarrow **wave nature is important**.
- **Heisenberg Uncertainty Principle:**
 - Fundamentally impossible to know both the exact position and exact momentum/velocity of an electron at the same time.
 - Measuring an electron (for example, by hitting it with a photon) disturbs it; this is a built-in limit of nature, not just bad equipment.

- de Broglie wavelength: $\lambda = h/(mv)$.
- **Typical Question Types**
 - Calculate or compare de Broglie wavelengths of particles with different masses and speeds.
 - Explain why we see wave behavior for electrons but not for macroscopic objects like soccer balls.
 - Explain the uncertainty principle qualitatively.
- **Common Traps**
 - Thinking the uncertainty principle is just due to bad measurements instead of a fundamental physical limit.
 - Forgetting that matter waves matter most for very small, light particles (like electrons).

6.5 Quantum Mechanics and Atomic Orbitals

- **From Bohr to Quantum Mechanics:**
 - The quantum mechanical model combines quantized energies, wave nature of matter, and the uncertainty principle.
 - Schrödinger developed a wave equation that describes electrons in atoms.
- **Schrödinger's Wave functions and Orbitals:**
 - Solutions to Schrödinger's equation are wave functions that describe atomic orbitals.
- **Atomic orbital:** a 3D region of space around the nucleus where there is a high probability of finding an electron (often drawn as enclosing about 90% of the electron's probability).
- **Orbit vs Orbital:**

- **Bohr orbit:** fixed circular path with defined radius (like a mini planet orbiting the nucleus).
- **Quantum orbital:** no fixed path, a probability cloud showing where the electron is likely to be found.
- Energy levels are still quantized, but electrons are not in neat circular paths.
- **Typical Question Types**
 - Define atomic orbital (must mention region of space and probability).
 - **Compare orbit vs orbital. (Remember they are different!)**
 - Explain why we can't state the exact location of an electron in an atom.
- **Common Traps**
 - Saying that an orbital is a "path" or "track" instead of a region of high probability.
 - Forgetting that even though we use probability clouds, the energy levels are still quantized.

6.6 Representations of Orbitals

- Different atomic orbitals denoted by **s, p, d, f**
 - s: Spherical shape
 - p: Dumbell shape
 - d: Four leaf clover shape
- For a given principle energy greater than 1, there's **1s orbital, 3p, 5d, 7f**, due to the different orientations of the orbitals.
- The principle quantum number n always equals the number of sublevels within that principle energy level.
- **For example:** 1s 2s 3s orbitals are identical in size and orientation, **but their sizes are increasing due to electrons with increasing energy.**

6.7 Many Electron Atoms

- In one electron atom, subshells all have the same energy.
- In many electron atoms, electron-electron repulsions cause different subshells at different energy. **Energies of subshells follow order $ns < np < nd < nf$** (e.g. $n=3$ orbitals increase in energy in the order $3s < 3p < 3d$.)
- In many electron atoms, orbital sets in the same sublevel have the same energy. Chemists call them **degenerate orbitals**.
- The range of energy levels within a principal energy level can overlap the energy levels of another principal level.
- **Aufbau Principle**
 - According to the Aufbau principle, **electrons occupy the orbitals of lowest energy first.**
 - Orbitals of the same sublevel always have the same energy.
 - **On the same principle energy level, $s < p < d < f$**
 - The range of energy levels within a principal energy level can **overlap** the energy levels of another principal level. (e.g. **4s and 3d**)
 - **When writing electron configurations, look out for s and d! For example, when 4s (2) and 3d (4), 4s should give one electron to 3d to make it more stable, making it 4s (1) 3d (5)**
- **Energy Ranking of electrons : 1s 2s 2p 3s 3p 4s 3d 4p 5s 4d 5p 6s 4f 5d 6p 7s 5f 6d 7p...**
- **Pauli Exclusion Principle** : an atomic orbital can hold a maximum of two electrons and they must have opposite spin.
 - According to the Pauli exclusion principle, when electrons with **opposite spins** occupy an orbital, the electrons are said to be **paired**; a **single electron** in one orbital is **unpaired**.

6.8 - 6.9 Electron Configurations and the Periodic Table

- **Orbital Diagrams**
 - Each box in the diagram represents one orbital.
 - Half-arrows represent the electrons.
 - The direction of the arrow represents the relative spin of the electron.
- **Electron Configurations**
 - A number denoting the energy level
 - A letter denoting the type of orbital
 - A superscript denoting the number of electrons in those orbitals.
- **Hund's Rule**
 - For degenerate orbitals, the lowest energy is attained when the number of electrons with the same spin is maximized.
 - This means that, for a set of orbitals in the same sublevel, there **must be one electron in each orbital before pairing and the electrons have the same spin, as much as possible.**
- **Abbreviated Electron Configurations (Condensed Electron Configurations)**
 - In writing the Abbreviated electron configuration of an element, **the electron configuration of the nearest noble-gas element of lower atomic number(core electrons) is represented by its chemical symbol in brackets.**
Eg: [Ne]
- **Inner shell electrons, Outer shell electrons and Valence electrons**
 - The inner-shell electrons are referred to as the **core electrons.**
 - The electrons given after the noble-gas core are called the outer-shell electrons. **The outer-shell electrons include the electrons involved in chemical bonding, which are called the valence electrons.**
 - **For the elements with atomic number 30 or less, all of the outer-shell electrons are valence electrons.** (The v.e. number for transition elements may vary. e.g. Fe [Ar]4s23d6)

- **Valence electrons are involved in chemical bonding** and are largely responsible for an atom's chemical behavior.
- **Electron Configuration of Excited State**
 - The most stable organization is the lowest possible energy, called the ground state. The electron configuration for an excited state electron shows one or more electrons in a higher energy level.
- **Electron Configuration for Ions**
 - **Transition metals will lose outmost shell electrons first, then lose the inner d level electrons.**
- **Exceptional Electron Configurations**
 - Some actual electron configurations differ from those assigned using the aufbau principle because although half-filled sublevels are not as stable as filled sublevels, they are more stable than other configurations.
- **Electron Configuration and the Periodic Table**
 - Chemists **fill orbitals in increasing order of energy.**
 - Different blocks on the periodic table correspond to different types of orbitals: **s and p-block are representative elements; d block are transition elements; f block are lanthanides and actinides, or inner transition elements**

s,p,d,f blocks in the periodic table.

