



IDX G10 AP Precalculus **Study Guide Issue S1 M1** **By Samuel, Edited by Darwyn**

NOTE: This is an official document by Indexademics. Unless otherwise stated, this document may not be accredited to individuals or groups other than the club IDX, nor should this document be distributed, sold, or modified for personal use in any way.

2.1 Polynomials

2.2 Long Divison; the Remainder and Factor Theorems

2.3 Graphing Polynomial Functions

2.4 Finding Maximums and Minimums of Polynomial

2.6 Solving Polynomial Equations by Factoring

2.7 General results for polynomial equations

3.1 Linear Inequalities; Absolute Value

2.1 Polynomials

Polynomials is a function of a form:

- $f(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + 2x^2 + a_1 x + a_0$ (n is a nonnegative integer and a are real numbers)
- Polynomial Function of degree n
- **Terms:** $ax \dots$; **Coefficients:** $a \dots$
- **Leading term:** the term containing the highest power of x : $a_n x^n$; **Leading Coefficient:** a_n

Degree	Name	Example (please try to write one by yourself)
0	constant	
1	linear	
2	quadratic	
3	cubic	
4	quartic	
5	quintic	

- **zeros of a function:** the roots of the equation $f(x)=0$

2.2 Long Division; the Remainder and Factor Theorems

Long Division: dividing $f(x) = 6x^3 - 2x^2 + 5x + 1$ by $x - 1$ yields


$$\text{Thus, } 6x^3 - 2x^2 + 5x + 1 = (x - 1)(6x^2 + 4x + 9) + 10.$$

$$\begin{array}{r}
 \begin{array}{c} 6x^2 + 4x + 9 \quad \leftarrow \text{Quotient} \\ \hline \end{array} \\
 \text{Divisor} \rightarrow x - 1 \overline{) 6x^3 - 2x^2 + 5x + 1} \leftarrow \text{Dividend} \\
 \begin{array}{r}
 6x^3 - 6x^2 \\
 \hline
 4x^2 + 5x \\
 4x^2 - 4x \\
 \hline
 9x + 1 \\
 9x - 9 \\
 \hline
 10 \leftarrow \text{Remainder}
 \end{array}
 \end{array}$$

- **Remainder Theorem:** When polynomial $P(x)$ is divided by $x - a$, the remainder is $P(a)$
 - $P(x) = (x - a) \cdot Q(x) + R \implies P(a) = R$
- **Factor Theorem:** A number a is a solution of the polynomial equation $f(x) = 0$ if and only if $x - a$ is a factor of $f(x)$


2.3 Graphing Polynomial Functions

The graphs of polynomial functions have certain fundamental shapes.




Quadratic function

$$y = x^2$$




Cubic function

$$y = x^3$$



Quartic function



$$y = x^4$$





Quintic function

$$y = x^5$$

Graph of $f(x) = ax^3 + bx^2 + cx + d$




Graph of $f(x) = ax^4 + bx^3 + cx^2 + dx + e$




➤ Effect of a Squared or Cubed Factor

- A polynomial function will have a **zero** that is a real number at each point at which it crosses the x-axis.

* If a polynomial $P(x)$ has a factor such as $(x-c)$ **only**, the graph crosses the x-axis **once**, at $x=c$ straight.




* If a polynomial $P(x)$ has a factor such as $(x-c)^2$, then $x=c$ is a **double root** of $P(x)=0$ and the graph is **tangent** to the x-axis at $x=c$.




Cubic

$$y = (x-1)(x-3)^2$$



Quartic


$$y = (x-1)(x-3)^2(x-4)$$



Quintic


$$y = x^2(x-1)(x-3)^2$$

* If a polynomial $P(x)$ has a factor such as $(x-c)^3$, then $x=c$ is a **triple root** of $P(x)=0$ and the graph flattens out around $(c,0)$ and crosses the x-axis at the point.




Cubic

$$y = (x-2)^3$$



Cubic

$$y = -(x-2)^3$$



Quartic

$$y = (x-1)^3(x-3)$$

2.4 Finding Maximums and Minimums of Polynomial

- Quadratic Functions:** the **Maximum or Minimum value** of the quadratic function $y = ax^2 + bx + c$ occurs at $x = -b/2a$
- Cubic functions**
 - Point of inflection: **Points of inflection** of a polynomial function occur at input values where the **rate of change** of the function changes from **increasing** to **decreasing** or from **decreasing** to **increasing**. This occurs where the graph of a polynomial function changes from **concave down** to **concave up** or from **concave up** to **concave down**
- Local maximum and local minimum**

- When the graph of a cubic function has a “**peak**” and a “**valley**”, the function has a **local maximum** at the **highest point** of the **peak** and a **local minimum** at the **lowest point** of the **valley**
- the **highest point** among the **local maximum** is the **global maximum**; the **lowest point** among the **local minimum** is the **global minimum**
- **Local Extrema Theorem:** A polynomial function of degree n has at most $n-1$ relative maxima/minima
- **Extreme Value Theorem:** If a polynomial function f is on a closed interval $[a,b]$, then f has both a maximum and a minimum value on $[a,b]$
- **Intermediate Value Theorem:** If $P(x)$ is a polynomial function such that $a < b$ and $P(a)$ is not equal to $P(b)$, then $P(x)$ takes on every value between $P(a)$ and $P(b)$ in the interval $[a,b]$
- **The Location Principle:** if $P(a) \cdot P(b) < 0$, then $P(x)$ has at least one zero between a and b .

2.6 Solving Polynomial Equations by Factoring

- **Rational Root Theorem**

➤ **Rational Root Theorem**

Use factoring to solve the general polynomial equation,

$$a_n x^n + a_{n-1} x^{n-1} + a_{n-2} x^{n-2} + \cdots + a_1 x + a_0 = 0$$

$$(qx - p)(\quad) = 0$$

If $qx - p$ is a factor (so that $x = \frac{p}{q}$ is a root), then q must divide a_n , and p must divide a_0 .

Let $P(x)$ be a polynomial of degree n with integral coefficients and a nonzero constant terms:

$$P(x) = a_n x^n + a_{n-1} x^{n-1} + a_{n-2} x^{n-2} + \cdots + a_1 x + a_0, \text{ where } a_0 \neq 0$$

If one of the roots of the equation $P(x) = 0$ is $x = \frac{p}{q}$ where p and q are nonzero integers with no common factor other than 1, then p must be a factor of the **constant term** a_0 and q must be a factor of the **leading coefficient** a_n .

- **Other ways**

- Cross multiplication
- use one variable to represent one complicated element
 - E.g. $t = 2x^2$
- $a^3 - b^3 = (a-b)(a^2 + ab + b^2)$
- $a^3 + b^3 = (a+b)(a^2 - ab + b^2)$

2.7 General results for polynomial equations

- **The Fundamental Theorem of Algebra:** In the **complex number system** consisting of all real and imaginary numbers, if $P(x)$ is a polynomial function of degree n ($n > 0$) with **complex coefficients**, then the equation $P(x)=0$ has **exactly n roots** (provided a double root is counted as 2 roots. a triple root is counted as 3 roots, and so on)
- **Complex Conjugates Theorem:** If $P(x)$ is a polynomial with **real coefficients**, and $a+bi$ is an imaginary root of the equation $P(x)=0$, the **$a-bi$** is also a root
- **Theorem 3:** Suppose $P(x)$ is a polynomial with **rational coefficients**, and a and b are rational numbers such that \sqrt{b} is irrational, if $a+\sqrt{b}$ is an imaginary root of the equation $P(x)=0$, then **$a-\sqrt{b}$** is also a root
- **Theorem 4:** If $P(x)$ is a polynomial of **odd degree with real coefficients**, then the equation $P(x)=0$ has **at least one real root**
- **Vieta's Formula:** For the equation $a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0 = 0$, with $a_n \neq 0$: the sum of the roots is $-a_{n-1}/a_n$; the product of the roots is a_0/a_n if n is even, and $-a_0/a_n$ if n is odd.

3.1 Linear Inequalities; Absolute Value

Quick Review

1. You can add the same number to (or subtract the same number from) both sides of an inequality.

$$a < b \Rightarrow a + c < b + c$$

2. You can multiply (or divide) both sides of an inequality by the same **positive** number.

$$a < b, c > 0 \Rightarrow ac < bc$$

3. You can multiply (or divide) both sides of an inequality by the same **negative** number if you *reverse the inequality sign*.

$$a < b, c < 0 \Rightarrow ac > bc$$

4. Another definition of absolute value of $x \in \mathbb{R}$ is: $|x| = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0 \end{cases}$

➤ Notations

- $1 < x < 5$ is equivalent to $(1, 5)$
- $0 \leq x \leq 4$ is equivalent to $[0, 4]$
- $10 \leq x < 40$ is equivalent to $[10, 40)$
- $x > 1$ is equivalent to $(1, \infty)$
- $x \leq 9$ is equivalent to $(-\infty, 9]$.

➤ Graphing

- A solid dot (\bullet) \rightarrow an endpoint is included in the graph;
- An open dot (\circ) \rightarrow the endpoint is not included in the graph
- **Combined inequality** \rightarrow its graph is the intersection of the graphs

Wrong combined inequalities: for example, $6 < x < 1$, it is the **empty set**. And we use \emptyset to describe.

➤ Absolute Value

- The absolute value of a number x , denoted $|x|$, is interpreted geometrically the **distance** from x to zero in either direction on the number line.

Sentences Involving $ x $ $c \geq 0$			
Sentence	Meaning	Graph	Solution
$ x = c$	The distance from x to 0 is <i>exactly</i> c units.		$x = c$ or $x = -c$
$ x < c$	The distance from x to 0 is <i>less than</i> c units.		$-c < x < c$
$ x > c$	The distance from x to 0 is <i>greater than</i> c units.		$x < -c$ or $x > c$

- $|x - k|$ is interpreted the distance from x to k in either direction on the number line

Sentences Involving $ x - k $			
Sentence	Meaning	Graph	Solution
$ x - 5 = 3$	The distance from x to 5 is 3 units.		$x = 2$ or $x = 8$
$ x - 1 < 2$	The distance from x to 1 is <i>less than</i> 2 units.		$-1 < x < 3$
$ x + 3 > 2$ ($ x - (-3) > 2$)	The distance from x to -3 is <i>greater than</i> 2 units.		$x < -5$ or $x > -1$

Sentence	Equivalent Sentence	For example
$ ax + b = c$	$ax + b = \pm c$	$ 2x - 3 = 4 \Rightarrow 2x - 3 = \pm 4$
$ ax + b < c$	$-c < ax + b < c$	$ 2x - 3 < 4 \Rightarrow -4 < 2x - 3 < 4$
$ ax + b > c$	$ax + b < -c$ or $ax + b > c$	$ 2x - 3 > 4 \Rightarrow 2x - 3 < -4$ OR $2x - 3 > 4$