



IDX G10 Chemistry H
Study Guide S1 Midterms
By TaeYun Kang, Edited by Edward Chen

NOTE: This is an official document by Indexademics. Unless otherwise stated, this document may not be accredited to individuals or groups other than the club IDX, nor should this document be distributed, sold, or modified for personal use in any way.

Contents:

1. Formula Weights
2. Avogadro's Number (N)
3. Empirical Formulas from Analyses
4. Quantitative Info from Equations
5. Limiting Reactants
6. Electrolyte

Formula Weights

- Definition: sum of the atomic weights/masses of the atoms in the chemical formula
- Molecular Weight: formula weight of a molecule
 - Unit: amu

Percentage Composition

- percentage of mass contributed by each element in the substance
- $\% \text{ composition} = \frac{\text{mass of element}}{\text{mass of substance}} \times 100\%$

Avogadro's Number (N)

- 1 mole (mol): a unit for measuring the number of microscopic particles
- Value of Avogadro's number: 6.022×10^{23}
 - 1 mol H₂O molecules = 6.022×10^{23} H₂O molecules

Masses, Molar Mass, and Numbers of Particles

- 1 mole of different substances will contain the same number of particles but different masses proportional to formula weights
- The concept of moles provides a bridge between mass and number of particles:

$$\begin{array}{ccccc} & \div M & & \times N & \\ m & \longleftrightarrow & n & \longleftrightarrow & \# \\ & \times M & & \div N & \end{array}$$

-
- m: mass, n: number of moles, #: number of particles, N: Avogadro's number
- Molar mass (M): The mass in grams of 1 mol of a substance
 - Numerically equal to molecular weight, but in grams per mol (g/mol)

Empirical Formulas from Analyses

- Empirical Formulas: chemical formulas that only give the ratio of moles (percentage composition) of each element in a molecule
- Thus, the EF of a substance can be found through moles

$$\begin{array}{ccccccc} & \text{assume} & & \div M & & \text{compare to} & \\ \% & \xrightarrow{100\text{-g sample}} & m & \longrightarrow & n & \xrightarrow{\text{the smallest } n} & \text{EF} \end{array}$$

- The calculated values for mole ratio should be rounded to appropriate whole numbers
 - Non-whole number ratios should be multiplied by a factor that converts them into whole numbers
 - E.g., 1:2.5 = 2:5, 1:1.33 \approx 3:4
- E.g., 11.1% H, 88.9% O by mass in the sample
 - 11.1g H, 88.9g O → 11.1mol H, 5.6mol O
 - H:O = 11.1:5.6 = 11.1/5.6:5.6/5.6 = 1.98:1 \approx 2:1
 - H₂O (empirical formula)

Molecular Formulas from Empirical Formulas

- The subscripts of MF are always whole-number multiples of the subscripts of EF
 - This multiple is found by dividing the molecular weight by the EF weight
 - E.g., molecular weight = 60 g/mol, empirical formula CH₂O (EF weight = 30 g/mol).
 - whole-number multiple = 60/30 = 2
 - C₂H₄O₂ (molecular formula)

Combustion Analysis

- Technique for determining EF in the lab (commonly used for hydrocarbon compounds)


- $C_xH_y + O_2 \rightarrow H_2O + CO_2$
- Calculate the # of moles of C and H in the original sample with the moles of H_2O and CO_2
- If a third element is present ($C_xH_yO_z$), its mass can be determined by subtracting the measured masses of C and H from original sample mass

Quantitative Info from Equations

- $2H_2 + O_2 \rightarrow 2H_2O$
 - 2 moles of H_2 reacts with 1 mole of O_2 to produce 2 moles of H_2O
- Stoichiometrically equivalent quantities: the quantity of moles given by coefficient
 - E.g. **2 mol H_2 , 1 mol O_2 , 2 mol H_2O**
 - Their relationship can be represented as $2 \text{ mol } H_2 \simeq 1 \text{ mol } O_2 \simeq 2 \text{ mol } H_2O$
 - \simeq : “is stoichiometrically equivalent to”

Stoichiometric Relationships

- For a balanced equation involving X and Y:



- These relationships are used to solve mass-mass problems

Limiting Reactants

- Definition: reactant that is completely consumed in a reaction
- Excess reactants: other reactants that are consumed
- Determining the limiting reactant by finding this ratio for each reactant:
 - Number of moles divided by coefficient
 - The substance with the smallest ratio is the limiting reactant

Theoretical Yields

- Definition: quantity of product calculated to form when all of a limiting reactant is consumed
- Actual yield: amount of product actually obtained
 - It is almost always less than but never greater than the theoretical yield because:

- Reactants do not fully react
- Side reactions other than the desired reaction occurs
- Not all of the product can be recovered from mixtures
- Chemistry is a natural science
- Percent yield: $\frac{\text{actual yield}}{\text{theoretical yield}} \times 100\%$

Electrolyte

- Solutions
 - Homogeneous mixtures of two or more pure substances
 - Solvent — the substance present in greatest abundance
 - Solute — the substance dissolved in the solvent
- Dissociation: when an ionic substance dissolves in water, the polar water molecules pull individual ions from the crystal and solvates them
- Electrolyte: a substance that, when dissolved in water, results in a solution containing ions and conducts electricity
- Non-electrolyte: substance that doesn't form ions and doesn't conduct electricity in a solution
 - Most molecular substances
- Strong electrolytes fully ionize in water, dissociating to ions completely or nearly completely
 - Good conductors
 - All water-soluble ionic compounds are strong electrolytes
 - Some molecular compounds are strong electrolytes (e.g., strong acids like)
- Weak electrolytes only partially ionize, and exist in solutions mostly in the form of molecules, with only a fraction as ions
- Solvation: process in which an ion is surrounded by water molecules arranged in a specific manner
- Double arrow: reversible reactions, showing a chemical reaction that can go both forwards and backwards