



## IDX G9 Physics S Study Guide Issue 1

By Yenny, Edited by Claire Kong

**NOTE: This is an official document by Indexademics. Unless otherwise stated, this document may not be accredited to individuals or groups other than the club IDX, nor should this document be distributed, sold, or modified for personal use in any way.**

### Table of contents:

1. Picturing Motion
2. Position
3. Graphs to Represent Motion
4. How fast
5. Accelerated Motion
6. Acceleration
7. Constant Acceleration

### **Picturing Motion**

- Picturing Motion
    - o Motion: Changing position over time to a frame of reference
    - o Motion Diagram is used to show the position of a moving object at equal time intervals
      - Purpose to visualize displacement and velocity
      - Direction represents the way object is moving
      - Particle model is a simplified version of a motion diagram. Particle is used to represent the object when its size is smaller than the distance and when the internal motion of it is not crucial
- Where and When

- Position represents 'where' and time represents 'when'
  - o Position can be negative when it opposes from the positive value but time always stays positive
  
- Coordinate system: Demonstrates the position of the zero point and the direction of the variable's increase or decrease value
  
- Time interval is the difference between two times
  - o  $\Delta t = t(\text{final}) - t(\text{initial})$
  
- Vector describes both magnitude and direction while scalar only describes magnitude
  - o Vector: Displacement, velocity, acceleration, force
  - o Scalar: Distance, speed, time, interval, mass, volume, density, temperature, energy
  
- Vector is represented by a line and an arrow
  - o Use scale to represent the length as magnitude (ex. 1cm=10m) and arrows to represent directions
  - o Use the symbol of an letter with arrow drawn above
  
- Resultant vector is the conclusion of two or more vectors
  - o How to find: Move the tail (end of the arrow) of one vector to the tip (head of the arrow) of another vector without the change of magnitude and direction. Always remember head-tail
  - o Represent the result vector as letter R with an arrow above it
  - o Subtracting vectors are done with two ways:
    - Method 1: Put tails of A and B together and find the opposing value which adds together
    - Method 2: Add -B to A
  
- Distance, position, and displacement is different
  - o Distance: How far apart objects are with no direction
  - o Displacement: The object's overall change in position with direction
  - o Position: Where is an object at specific time with direction

## **Position-Time Graphs**

- Position-time graph, also called p-t graph, is done with plotting the time on x axis and position on y axis
  - o When the object is at rest, the slope is a horizontal line parallel to x axis
  - o When the object is at uniform motion (no change in speed and direction) the slope is a slant straight line
- Information about object and its motion are shown
  - o y axis: instantaneous position (The position object is at a specific time)
  - o y intercept: initial position
  - o Position: Where
  - o Time: When
  - o Intersection of graph lines: where two objects meet

## **How Fast**

- Velocity is position divided by time
  - o Average velocity is the change in displacement over change in time
    - The slope of p-t graph indicates the average velocity
- Average speed is the absolute value of the slope of p-t graph
- Steeper slope means larger velocity
- Instantaneous velocity is the speed and direction of an object at a particular instant
- Motion diagram can be used to find average velocity
  - o Average velocity is the same direction as displacement and the proportional of displacement
  - o Distance equals the same of average velocity multiply time and initial displacement

## **Acceleration**

- Acceleration: Rate at which object's velocity changes

- o When the object is at uniform motion, the constant velocity is represented by a straight line.
- o When the object is at nonuniform motion, the velocity changes with acceleration
- o Large acceleration - large change in velocity and short time interval
- o Small acceleration - small change in large time interval
- Velocity-Time graph represents displacement with the area
- o Slant straight line: Uniformly accelerated motion
- o Rising graph line: acceleration is positive
- o Descending graph line: acceleration is negative
- o Horizontal line: velocity is constant and acceleration is 0
- o Steeper acceleration means larger absolute value of acceleration
- Average Acceleration: The change in velocity during some measurable time interval divided by that time interval
- Instantaneous Acceleration: The change in velocity at an instant of time by using the slope of the line or the tangent line
- Sign of the acceleration does not determine whether the object is speeding up or slowing down
- o Positive acceleration: positive direction
- o Negative acceleration: negative direction
- o Speed up: same direction with initial velocity
- o Slow down: opposite direction with initial velocity

#### Motion with Constant Acceleration

- Final velocity equals the sum of initial velocity and the multiple of average acceleration and change in time
- Position, velocity, and acceleration of d-t graph is considered when turning to v-t graph

- Formula to remember:

$$v_f = v_i + a\Delta t$$

$$\Delta d = \frac{(v_i + v_f)}{2} \Delta t = \bar{v} \times t$$

$$\Delta d = v_i t + \frac{1}{2} a t^2$$

$$\Delta d = \frac{(v_f^2 - v_i^2)}{2\bar{a}}$$