



IDX G9 Physics S
Study Guide Issue S1 Midterms
By Yenny, Edited by Claire

NOTE: This is an official document by Indexademics. Unless otherwise stated, this document may not be accredited to individuals or groups other than the club IDX, nor should this document be distributed, sold, or modified for personal use in any way.

Contents

1. Chapter 3: Accelerated Motion -3.3 Free Fall
2. Chapter 4: Forces in One Dimension

Chapter 3:

Free Fall

- What is Free Fall?
 - The motion of a falling object when air resistance is negligible and the action can be considered due to gravity alone
- History
 - Aristotle: a heavier object falls faster than a lighter object
 - - Galileo argued that neglecting the effect of air resistance, all objects in free fall have the same constant acceleration.
- Acceleration due to gravity
 - The acceleration of an object in free fall that results from the influence of **Earth's gravity**

- No matter what the falling object is made of, weighed, positioned, dropped down, or threw up
- On earth, $g = 9.8 \text{ m/s}^2$
- On moon, $g = \frac{1}{6}$ of Earth (About 1.63 m/s^2)
- Direction points towards the center of the planet, downward
- Small variations in g at different places on earth
- About Free fall
 - Falling downward: initial velocity is 0 m/s (If down is $+$, v_i is larger than 0 m/s)
 - Throwing upward: initial velocity is not 0 m/s (If up is $+$, v_i is smaller than 0 m/s)
- Thrown Upward
 - Upward is $+$
 - Acceleration is same all along (rise+fall)
 - Initial velocity is $+$, initial distance is 0
 - When it is at highest position, $v=0$, displacement is the largest
 - If the object is above initial position, displacement is $+$
 - If the object is below initial position, displacement is $-$
 - Rising: velocity is $+$, displacement is $+$
 - Falling: velocity is $-$, displacement is $+$

- When it returns to original position, $v=-v_i$, displacement is 0

Chapter 4:

Introduction of Forces

- What is a force?
 - A force is a push or pull. A force cannot be seen but it can show how a force affects on an object
- Effect of Force
 - Cause acceleration
 - a is not 0
 - Increase, decrease, change direction
 - Deformation (Change in dimension, shape)
- System and the external world
 - System is to whom the force is exerted
 - The internal world is everything around the system that exerts force on the system
- Agents by whom the force is exerted
- System and agent has interaction between only 2 objects
- Contact forces and field forces

- Contact force: an object from the external world touches a system and thereby exerts a force on it
- Field force: exerted without contact (ex. gravitational force, electric force, magnetic force)
- Types of forces
 - a. Gravity: A field force due to gravitational attraction between two objects, generally Earth and the object
 - Direction: straight down towards the center of the Earth
 - Gravity= mg
 - Agent: Earth, System: object
 - b. Normal: a perpendicular contact force exerted by a surface on an object
 - Direction: perpendicular to the surface and away from the surface
 - Agent: Surface, System: object
 - c. Thrust: a general term for the forces that move objects such as rockets, planes, cars, and people
 - Direction: in the same direction as the acceleration of the object
 - Agent: Gas expelled by the engine, System: rocket
 - d. Tension: the pull exerted by a string, rope, or cable when attached to a body and pulled taut
 - Direction: away from the object and parallel to the string

- Agent: Rope, System: object
- e. Spring: a restoring force that is the push or pull a spring exerts on an object
- Direction: opposite the displacement of the object at the end of the spring
- Agent: spring, System: object attached to the spring
- f. Friction: the contact force that acts to oppose sliding motion between surfaces
- Direction: Parallel to the surface and opposite the direction of sliding
- Agent: ground, System: object

Free-body diagram

- Pictorial model:
 - First circle the system, find the contact with external world, identify contact force, and finally identify field forces
- Free-body diagram:
 - Apply the particle model to the pictorial model
 - Arrow to represent a force, label it (Mind direction, length)

Combining Forces in One Dimension

- Net force: the vector sum of all the forces on an object
- Combining forces in one dimension

- Same direction: $F_{\text{net}} = F_1 + F_2$
- Opposite direction: $F_{\text{net}} = F_1 - F_2$

Force and Acceleration

- One way to find out is by doing experiments
 - Controlled force exerted horizontally
 - Very smooth surface to reduce the resistance

Newton's Laws

- Galileo's Thought Experiment
 - If friction could be emitted,
 - Initial height equals final height
 - Thought experiment since it can't become reality
- First Law
 - Galileo concluded that in the ideal case of 0 resistance, horizontal motion would never stop
 - Newton: an object that is at rest will remain at rest, and an object that is moving will continue to move in a straight line with constant speed, if and only if the net force acting on that object is zero
- Inertia
 - Law of inertia = another name of Newton's first law

- Inertia is the tendency of an object to resist change
 - Not a force
- Equilibrium: The state when the net force on an object is zero
 - Two states of equilibrium: $a = 0$
 - At rest
 - Moving at a constant velocity
 - Net force = cause of velocity change
 - No net force = equilibrium
 - Newton's first law identifies a net force as something that disturbs a state of equilibrium
- Second Law
 - The acceleration of an object is equal to the sum of the forces acting on the object, divided by the mass of the object
 - $F_{\text{net}} = ma$
 - Steps:
 - Identify all the forces acting on the object
 - Find the net force
 - Use Newton's second law to calculate the acceleration

- Mass and Weight
 - Mass is the amount of matter that makes up an object
 - Mass of an object is always the same everywhere in the universe
 - Unit: kilograms
 - Weight is the force due to the pull of gravity on an object
 - The weight of an object will very depend on its location in the universe
 - Unit: Newtons