



IDX G10 CHEM H

Study Guide Issue #1

By TaeYun Kang, Edited by Edward Chen

**NOTE: This is an official document by Indexademics. Unless otherwise stated, this document may not be accredited to individuals or groups other than the club IDX, nor should this document be distributed, sold, or modified for personal use in any way.**

## Contents:

1. [Intermolecular Forces: Liquids and Solids](#)
2. [Stoichiometry](#)

## **Intermolecular Forces: Liquids and Solids**

### Review: Intermolecular Forces

- **Intermolecular forces (IMFs)** are attractive forces between molecules
- **Intramolecular forces** are attractive forces within molecules
- Three types of IMFs:
  - **London Dispersion Forces (LDF)** – attraction force between induced dipoles
    - A symmetrical, spherical atom loses its spherical shape and becomes a **temporary dipole**, which has a separation of opposite charges, creating **induced dipoles** in adjacent atoms
    - LDF increases with increasing molecular/atomic size and mass, since larger electron clouds are more easily distorted (more polarizable)
    - Present in both nonpolar and polar molecules
  - **Dipole-Dipole Forces** – attraction between permanent dipoles
  - **Hydrogen Bonds** – strong dipole-dipole force between the H of one molecule (bonded to N/O/F) and the N/O/F of another molecule
    - In a covalent bond of H-N/O/F, the partial charges are extremely large due to the high difference in electronegativity
    - Large partial charges result in strong dipole-dipole forces

Use IMF to explain properties

- Physical properties of halogen (7A)
  - Colored Gas:
    - Fluorine – Pale Yellow
    - Chlorine – Green
    - Bromine – Red
    - Iodine – Purple/Black
  - Poisonous
  - Diatomic
  - Density of gas increases down the group, leading to deeper color and higher b.p.


Select Properties of Liquids

- **Surface tension** is the amount of energy needed to stretch/increase the surface of a liquid by a unit area
  - Stronger IMFs → higher surface tension
- **Cohesion** is the intermolecular attraction between like molecules
- **Adhesion** is the intermolecular attraction between different molecules
- **Viscosity** is a measure of a fluid's resistance to flow
  - Stronger IMFs → higher viscosity
- **Volatility** is a measure of a fluid's tendency to vaporize
- **Vapor pressure** of a liquid is the pressure exerted by its vapor when the liquid and vapor are in dynamic equilibrium
  - **Dynamic equilibrium** is the condition where two opposing processes occur at the same time and rate
  - Stronger IMFs → Lower Vapor Pressure at a certain temperature
  - **Normal boiling point:** b.p. of a liquid at 1atm (760 torr)

Phase Change and Phase Diagram

- **Melting** – solid to liquid
- **Freezing** – liquid to solid
- **Vaporization** – liquid to gas
- **Condensation** – gas to liquid
- **Sublimation** – solid to gas
- **Deposition** – gas to solid
- **Endothermic** (absorb energy) – melting, vaporization, sublimation
- **Exothermic** (release energy) – freezing, condensation, deposition

- **Phase diagram** summarizes the conditions at which a substance exists as a solid, liquid, or gas
  - The curves represent boundaries of phase change
  - **Triple Point:** the temperature and pressure where three phases coexist in equilibrium
  - **Critical Point:** beyond this point, liquid and gas are indistinguishable



## Structure of Solids

- **Crystalline Solid** possesses rigid and long-range order
  - Atoms, molecules, or ions occupy specific positions
- **Amorphous Solid** does not possess a well-defined arrangement and long-range molecular order
- **Unit Cells** are basic repeating structural units of a crystalline solid
- Ionic Crystals
  - Cations and anions as lattice points
  - Held together by electrostatic attraction
  - Hard, brittle, high m.p.
  - Poor conductor of heat and electricity
- Covalent Crystals
  - Atoms as lattice points
  - Held together by covalent bonds
  - Hard, high m.p.
  - Poor conductor of electricity (except graphite)
  - Diamond v.s. Graphite
    - Diamond
      - Hard due to rigid covalent bonds
      - Poor conductor of heat and electricity
    - Graphite
      - Soft and slippery due to layered structure with LDF
      - Good conductor of heat and electricity due to delocalized electrons
- Molecular Crystals
  - Molecules as lattice points

- Held together by IMF
- Soft, low m.p.
- Poor conductor of heat and electricity
- Metallic Crystals
  - Metal atoms as lattice points
  - Held together by metallic bonds
  - Can be soft/hard, with low/high m.p.
  - Good conductors of heat and electricity

## Stoichiometry

### Stoichiometry

- Derived from στοιχεῖον (element) and μέτρον (measure)
- Area of study that examines the quantities of substances consumed and produced in chemical reactions
- Built on understanding of atomic mass, chemical formulas, and the Law of Conservation of Mass

### Chemical Equations

- Reactant(s) → Product(s)
  - **Reactants:** starting substances, left of the arrow
  - +: read as "reacts with"
  - →: read as "produces" or "yields"
  - **Products:** substances produced, right of arrow
  - **Coefficients:** numbers in front of formulas, relative numbers of molecules
    - Should use simplest whole-number ratio for coefficients
  - (g) for gas, (l) for liquid, (s) for solid, (aq) for aqueous solution, Δ above the arrow for a heated reaction
- Equations have to be balanced, meaning that they need an equal number of atoms of each element on each side

### Simple Patterns

- **(Direct) Combinations Reactions/Syntheses** is when 2+ substances react to form 1 product
  - A + B → C
- **Decomposition Reaction/Analyses** is when 1 substance breaks down into 2+ substances
  - C → A + B

- **Combustion Reactions** are rapid reactions that produce a flame, most involve O<sub>2</sub> as a reactant
  - X + O<sub>2</sub> → CO<sub>2</sub> + H<sub>2</sub>O

### Chemical Reactions of 1A Elements (Alkali Metals)

- Reaction with water
  - Metal fizzes and floats and disappears
  - The test tube becomes warm
  - Indicator (pH tester) turns pink
    - Formation of an alkali (XOH)
  - Reactivity of alkali metals increase down the group
  - Word equation: sodium + water → sodium hydroxide + hydrogen
  - Chemical equation: 2Na + 2H<sub>2</sub>O → 2NaOH + H<sub>2</sub>
- Reaction with Chlorine
  - Burst into flame, burns bright, forms chloride
  - Word equation: sodium + chlorine → sodium chloride
  - Chemical equation: 2Na + Cl<sub>2</sub> → 2NaCl
- Reaction with Oxygen
  - Burst into flame, burns fiercely to form oxides
  - Word equation: sodium + oxygen → sodium oxide
  - Chemical equation: 4Na + O<sub>2</sub> → 2Na<sub>2</sub>O