



NOTE: This is an official document by Indexademics. Unless otherwise stated, this document may not be accredited to individuals or groups other than the club IDX, nor should this document be distributed, sold, or modified for personal use in any way.

2.1 Rate of Change and Limits2.2 Limits Involving Infinity2.3 Continuity2.4 Rate of Change and Tangent Lines3.1 Derivative of a Function3.2 Differentiability3.3 Rules for Differentiation3.5 Derivatives of Trigonometric Functions3.6 Chain Rule3.7 Implicit Differentiation**2.1 Rate of Change and Limits**

For any $\varepsilon > 0$, there exists a $\delta > 0$, such that whenever $0 < |x - c| < \delta$,

$$|f(x) - L| < \varepsilon$$

Then we say f has a limit L as x approaches c . for notation, we write

$$\lim_{x \rightarrow c} f(x) = L$$

Properties of Limits

If L, M, c, k are real numbers, and $\lim_{x \rightarrow c} f(x) = L$, $\lim_{x \rightarrow c} g(x) = M$

- $\lim_{x \rightarrow c} k = k$
- $\lim_{x \rightarrow c} x = c$

- $\lim_{x \rightarrow c} (f(x) + g(x)) = L + M$
- $\lim_{x \rightarrow c} (f(x) - g(x)) = L - M$
- $\lim_{x \rightarrow c} (k \cdot f(x)) = k \cdot L$
- $\lim_{x \rightarrow c} (f(x) \cdot g(x)) = L \cdot M$
- If $M \neq 0$, then $\lim_{x \rightarrow c} \frac{f(x)}{g(x)} = \frac{L}{M}$
- If r and s are integers and $s \neq 0$, $L^{\frac{r}{s}}$ is a real number, then $\lim_{x \rightarrow c} (f(x))^{\frac{r}{s}} = L^{\frac{r}{s}}$

Left-handed and right-handed limits

$$\lim_{x \rightarrow c^-} f(x) \text{ or } \lim_{x \rightarrow c^+} f(x)$$

- $\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1$

Sandwich Theorem (Squeeze Theorem)

If $g(x) \leq f(x) \leq h(x)$ for all $x \neq c$ in the neighborhood of c , and

$$\lim_{x \rightarrow c} g(x) = \lim_{x \rightarrow c} h(x) = L$$

Then

$$\lim_{x \rightarrow c} f(x) = L$$

Average and Instantaneous Speed

Average speed: $\frac{\Delta y}{\Delta t}$

Instantaneous speed: $\lim_{\Delta t \rightarrow 0} \frac{\Delta y}{\Delta t}$

2.2 Limits Involving Infinity

Finite Limits as $x \rightarrow \pm \infty$

Horizontal Asymptote:

The line $y = b$ is a horizontal asymptote of the graph of a function $y = f(x)$ if either

$$\lim_{x \rightarrow \infty} f(x) = b \text{ or } \lim_{x \rightarrow -\infty} f(x) = b$$

Note: Rational functions have at most one horizontal asymptote, while other functions have at most two horizontal asymptotes.

Vertical Asymptote:

The line $x = a$ is a vertical asymptote of the graph of a function $y = f(x)$ if either

$$\lim_{x \rightarrow a^-} f(x) = \infty \text{ or } -\infty$$

Or

$$\lim_{x \rightarrow a^+} f(x) = \infty \text{ or } -\infty$$

Note: all properties that apply to limits apply to limits involving infinity, including the sandwich theorem.

End Behavior Model

The function g is

(a) a right end behavior model for $f \Leftrightarrow \lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = 1$

(b) a left end behavior model for $f \Leftrightarrow \lim_{x \rightarrow -\infty} \frac{f(x)}{g(x)} = 1$

If one function provides both a left and right end behavior model, it is simply called end behavior model.

2.3 Continuity

Continuity at a Point

Interior point:

A function $y = f(x)$ is continuous at an interior point c of its domain if

$$\lim_{x \rightarrow c} f(x) = f(c)$$

Endpoint:

A function $y = f(x)$ is continuous at left endpoint a of its domain if

$$\lim_{x \rightarrow a^+} f(x) = f(a)$$

A function $y = f(x)$ is continuous at right endpoint b of its domain if

$$\lim_{x \rightarrow b^-} f(x) = f(b)$$

Types of Discontinuity

Removable discontinuity: $f(x) \neq \lim_{x \rightarrow c} f(x)$

Jump discontinuity: $\lim_{x \rightarrow c^-} f(x) \neq \lim_{x \rightarrow c^+} f(x) \Rightarrow \lim_{x \rightarrow c} f(x) DNE$

Infinite discontinuity: $\lim_{x \rightarrow c^-} f(x) = \pm \infty$ or $\lim_{x \rightarrow c^+} f(x) = \pm \infty \Rightarrow \lim_{x \rightarrow c} f(x) DNE$

Oscillating discontinuity

Continuous functions

Functions are continuous on an interval iff it is continuous at every point of its domain

Properties of Continuous Functions

If the functions f and g are continuous at $x = c$, then the following combinations are continuous at $x = c$,

1. sums: $f + g$
2. differences: $f - g$
3. products: $f \cdot g$
4. constant multiples: $k \cdot f$, for any number k
5. quotients: $\frac{f}{g}$, provided $g(c) \neq 0$

If the function f is continuous at c and g is continuous at $f(c)$, then the composite $g \circ f$ is continuous at c .

Intermediate Value Theorem

A function $y = f(x)$ that is continuous on a closed interval $[a, b]$ takes on every value between $f(a)$ and $f(b)$

Corollary 1: Extreme value theorem (EVT)

A function $y = f(x)$ that is continuous on a closed interval $[a, b]$ must have its maximum and minimum, and hence takes on every value between its maximum and minimum values that occur on $[a, b]$.

Corollary 2: Zero's Theorem

If a function $y = f(x)$ that is continuous on a closed interval $[a, b]$, and if

$$(f(a) \cdot f(b)) < 0$$

Then there exists $c \in (a, b)$ such that $f(c) = 0$

2.4 The Rate of Change of Tangent Lines

Difference Quotient

The difference quotient for f at a is given by the expression

$$\frac{f(a+h) - f(a)}{h}$$

Slope of a Curve

The slope of a curve $y = f(x)$ at a point $P(a, f(a))$ is the number

$$m = \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}$$

Provided the limit exists.

- The tangent to the curve at P is the line through P with this slope.

Normal to a Curve

Normal line: the line perpendicular to the tangent at a point

Rates of Change

The average rate of change of a function over an interval is

$$\frac{\text{amount of change in function value}}{\text{length of the interval}} = \frac{\Delta y}{\Delta x} = \frac{f(x_2) - f(x_1)}{x_2 - x_1}$$

The instantaneous rate of change of a function at a point $(x, f(x))$ is

$$\lim_{\Delta x \rightarrow 0} \frac{\Delta y}{\Delta x} = \lim_{x_2 \rightarrow x_1} \frac{f(x_2) - f(x_1)}{x_2 - x_1}$$

3.1 Derivative of a Function

Definition of Derivatives

Derivative of a point:

The derivative of the function f at $x = a$ is defined by

$$f'(a) = \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}$$

Provided the limit exists.

Derivative as a function with respect to x :

The derivative of the function f with respect to x is the function f' whose value at x is

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

Provided the limit exists.

Alternate definition: the derivative of the function f at $x = a$ is limit

$$f'(a) = \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a}$$

Provided the limit exists.

Notations

$$y', \frac{dy}{dx}, \frac{df}{dx}, \frac{d}{dx} f(x), f'(x)$$

One-sided Derivatives

$$\text{Right-handed derivative: } f'_+(a) = \lim_{h \rightarrow 0^+} \frac{f(a+h) - f(a)}{h}$$

$$\text{Left-handed derivative: } f'_-(a) = \lim_{h \rightarrow 0^-} \frac{f(a+h) - f(a)}{h}$$

If $f'(x) > 0$, the slope of the tangent line is positive, so f is increasing.

If $f'(x) < 0$, the slope of the tangent line is negative, so f is decreasing.

3.2 Differentiability

Types of Nonexistence of $f'(a)$

If $f'(a)$ exists, then

1. $f(a)$ is defined
2. $\lim_{h \rightarrow 0^+} \frac{f(a+h) - f(a)}{h} = \lim_{h \rightarrow 0^-} \frac{f(a+h) - f(a)}{h}$

Nonexistence:

1. Corner: $f'_+(a)$ and $f'_-(a)$ exist but differ
2. Vertical tangent: $f'_+(a) = \infty$, $f'_-(a) = \infty$ or $f'_+(a) = -\infty$, $f'_-(a) = -\infty$
3. Cusp: $f'_+(a) = \infty$, $f'_-(a) = -\infty$ or $f'_+(a) = -\infty$, $f'_-(a) = \infty$
4. Discontinuity

Derivatives on a Calculator

Symmetric difference quotient (numerical derivative of f):

$$NDER(f(x), a) = \frac{f(a+h) - f(a-h)}{2h}$$

Where default $h = 0.001$, but still can be changed

Continuity \nRightarrow Differentiability

Differentiability \Rightarrow Continuity

Note: IVT also applies for derivatives

If a and b are any two points in an interval on which f is differentiable, then f' takes on every value between $f'(a)$ and $f'(b)$

3.3 Rules for Differentiation

Rule 1 Derivative of a Constant

If c is a constant, then

$$\frac{d}{dx}(c) = 0$$

Rule 2 Power Rules for Positive Integer Powers of x

If n is a positive integer, then

$$\frac{d}{dx}(x^n) = nx^{n-1}$$

Rule 3 The Constant Multiple Rule

If u is a differentiable function of x and c is a constant, then

$$\frac{d}{dx}(cu) = c \frac{du}{dx}$$

Rule 4 The Sum and Difference Rule

If u and v are differentiable functions of x , then their sum and difference are differentiable at every point where u and v are differentiable. At each points,

$$\frac{d}{dx}(u \pm v) = \frac{du}{dx} \pm \frac{dv}{dx}$$

Rule 5 the Products Rule

The product of two differentiable function u and v is differentiable, and

$$\frac{d}{dx}(uv) = v \frac{du}{dx} + u \frac{dv}{dx}$$

Rule 6 The Quotient Rule

At a point where $v \neq 0$, the quotient u/v of two differentiable functions is differentiable, and

$$\frac{d}{dx}\left(\frac{u}{v}\right) = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$$

Rule 7 Power Rules for Negative Integer Powers of x

If n is a negative integer and $x \neq 0$, then

$$\frac{d}{dx}(x^n) = nx^{n-1}$$

Second and Higher Order Derivatives

Second derivative: derivative of y'

$$f''(x) = y'' = \frac{dy'}{dx} = \frac{d}{dx}\left(\frac{dy}{dx}\right) = \frac{d^2y}{dx^2}$$

Third derivative:

$$f'''(x) = y''' = \frac{dy''}{dx} = \frac{d}{dx}\left(\frac{d^2y}{dx^2}\right) = \frac{d^3y}{dx^3}$$

The n -th derivative:

$$f^{(n)}(x) = y^{(n)} = \frac{dy^{(n-1)}}{dx} = \frac{d^n y}{dx^n}$$

3.5 Derivative of Trigonometric Functions

Derivatives of Sine and Cosine Functions

$$\frac{d}{dx}(\sin x) = \cos x$$

$$\frac{d}{dx}(\cos x) = -\sin x$$

Derivatives of Other Trigonometric Functions

$$\frac{d}{dx}(\tan x) = \sec^2 x$$

$$\frac{d}{dx}(\cot x) = -\csc^2 x$$

$$\frac{d}{dx}(\sec x) = \sec x \cdot \tan x$$

$$\frac{d}{dx}(\csc x) = -\csc x \cdot \cot x$$

These can be derived by differentiation of quotient rule.

3.6 Chain Rule

If f is differentiable at the point $u = g(x)$, and g is differentiable at x , then the composite function $(f \circ g)(x) = f(g(x))$ is differentiable at x , and

$$(f \circ g)'(x) = f'(g(x)) \cdot g'(x)$$

Alternatively:

If $y = f(u)$ and $u = g(x)$, the

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx}$$

Where $\frac{dy}{du}$ is evaluated at $u = g(x)$.

3.7 Implicit Differentiation

Implicit Defined Function

An equation that defines several functions implicitly (hidden inside the equation), without giving explicit formulas. A obvious example would be the function $y = e^{x+y} + 2$.

Implicit Differentiation Process

1. Differentiate both sides of the equation with respect to x (or another destined variable)
2. Collect the terms with $\frac{dy}{dx}$ on one side of the equation
3. Factor out $\frac{dy}{dx}$
4. Solve for $\frac{dy}{dx}$

Power Rule for Rational Powers

If n is any rational number, then

$$\frac{d}{dx}(x^n) = nx^{n-1}$$

If $n < 1$, then the derivative does not exist at $x = 0$ due to the existence of a infinite discontinuity at the point $x=0$