



**IDX G10 Biology H**  
**Study Guide S1 Midterm**  
**By Arianna, Edited by Amy**

**NOTE: This is an official document by Indexademics. Unless otherwise stated, this document may not be accredited to individuals or groups other than the club IDX, nor should this document be distributed, sold, or modified for personal use in any way.**

**Contents: Include a table of contents at the beginning of each guide. Follow the formatting here.**

1. 7.3 Cell transport
2. 8.1 Energy and Life
3. 8.2 Photosynthesis: An Overview
4. 8.3 Reactions of Photosynthesis

### 7.3 Cell transport

- Solution (e.g. cytoplasm): liquid mixture with  $\geq 2$  substances
- Concentration  $\rightarrow$  mass/volume, mole/litre
- Passive transport  $\rightarrow$  no need energy  $\rightarrow$  diffusion, facilitated diffusion
- active transport

#### 1) Diffusion

- molecules tend to move from high concentration to low concentration until reaching equilibrium
- cause substance to move across cell membrane but does not require energy
- equilibrium: concentration on both side is the same
  - no further changes in concentration
  - solutes continue to move in both directions in equal number (no net movement)
- higher temperature, faster rate of diffusion
- greater concentration difference, faster rate of diffusion

#### 2) Facilitated Diffusion

- movement of specific molecules across cell membranes through transport protein
- no energy consumes
- high to low concentration
- temperature, concentration difference, number of carrier or channel proteins in membrane increase, rate increases
- Osmosis
  - Involve only diffusion of water across selectively permeable membrane, which only some substances can pass through
  - water move from low to high solute concentration to dilute the concentrated solutions
  - water moves from more free water to less free water spaces
  - many cells contain water channel proteins named aquaporins that allow water to pass faster through cell membrane
  - isotonic solution: concentration of water and solute be the same on both sides of membrane
  - hypertonic solution: higher concentration of solute
  - hypotonic solution: lower concentration of solute

- Osmosis: water moves from hypotonic solution to hypertonic
- Osmotic pressure: pressure exerted by osmosis on the hypertonic side of selectively permeable membrane
- Plant cell and bacteria have cell wall to prevent cell from expanding
- Contractile vacuole: in some protozoans, pump excess water out of cell

### 3) Active Transport

- requires energy(ATP)
- protein pump: moves materials against concentration difference (from low to high concentration)
  - Eg: Na<sup>+</sup>-K<sup>+</sup> Pump: pumping Na<sup>+</sup> out and K<sup>+</sup> in against concentration gradients
- bulk transport: endocytosis and exocytosis, fuse of vesicle
  - transport large molecules or a large amount of small molecules/ions by movement of cell membrane
  - Endocytosis: process of taking material into cell by infoldings or pockets of cell membrane that forms a vesicle entering the cytoplasm
    - ① phagocytosis: process of large particles taken into cell by endocytosis
    - ② pinocytosis: process of solution taken into cell
  - Exocytosis: removal of large amount of material from cell, membrane of vesicles fuses with cell membrane

### 8.1 Energy and Life

- forms of energy include light, heat, electricity, and chemical energy
- Energy stored in ATP (Adenosine Triphosphate)→Adenine (contain nitrogen), ribose (5 carbon sugar), three phosphate group
- Adenosine diphosphate (ADP)→adenine, ribose, 2 phosphate groups
- Cells can store small amount of energy by adding a phosphate group to ADP
- ATP is used by all types of cell as basic energy source
- Release energy: subtract third phosphate group
- Using the energy
  - ① Active transport: sodium potassium pump uses 1 ATP to move 3 Na<sup>+</sup> out and 2 K<sup>+</sup> in
  - ② Motor protein move organelles along microtubules using ATP
  - ③ Protein synthesis

- ④ Responses to chemical signals at cell surface
- ⑤ Produce light
- Glucose store 90 times more energy than ATP
- cells regenerate ATP from ADP by using energy in food (e.g. glucose, protein, lipid)
- Autotroph: produce food by photosynthesis/chemosynthesis
- Animal(sea slug) can carry out photosynthesis
- Heterotroph: obtain energy from food that are consumed

## 8.2 Photosynthesis: An Overview

- Discovery of photosynthesis
  - 1643, Jan Van Helmont watered plant for 5 years and measured the mass of the soil. He found that most of the mass increase of plant came from water
  - 1711, Joseph Priestley used a candle, jar, mint plant to experiment. The mint released oxygen gas
  - 1779, Jan Ingenhousz, light is necessary for plants to produce oxygen gas
- In the presence of light, plants transform carbon dioxide and water into carbohydrates and release oxygen
- Light and pigment
  - white sunlight is a mixture of different colors
  - visible light made by visible spectrum
  - photosynthesis requires light and pigment, molecules in chloroplast
  - chlorophyll: principle pigment, includes chlorophyll a and b
  - chlorophyll absorbs well in violet-blue and red regions, but not well in green region
  - carotenoid: red, orange, yellow pigments, absorb light in violet and blue green regions
    - As temperature drops, chlorophyll break down first, leaving carotenoid pigments to see in the autumn
- Chloroplast
  - outer and inner membranes
  - thylakoid: saclike photosynthetic membrane containing chlorophyll
  - stroma: fluid portion, outside of a thylakoid
  - grana: stacks of thylakoid membrane
- Energy Collection
  - chlorophyll absorbs light

- the energy from light are collected in the electrons in chlorophyll--energy levels are raised--high energy electrons make photosynthesis work
- Electron Carriers
  - a compound that can accept a pair of high energy electrons and transfer them to another molecules
  - e.g. NADP+
- Oxidation and reduction
  - oxidized: substances lose electrons
  - reduction: gain electrons
- Light dependent reaction
  - thylakoid membrane
  - need light and water
  - release oxygen gas
  - produce ATP, NADPH
- Light independent reaction
  - stroma
  - need carbon dioxide
  - need ATP, NADPH
  - produce sugar

### 8.3 Reactions of Photosynthesis

#### Light dependent reaction

- Thylakoid membrane
- Photosystems
  - clusters of chlorophyll and proteins
  - contain light harvesting complex
    - accessory pigments
    - a matrix of protein
  - contain reaction center complex
    - a pair of chlorophyll a
    - a matrix of protein
    - a primary electron acceptor
- use solar energy to produce oxygen gas and convert ADP, NADP+ into ATP and NADPH
- photosystem I and II are connected by the transfer of electrons through electron transport chain

#### A. Photosystem II

- pigments(P680)
- energy--electron--ETC
- photolysis: water split to replenish the lost high energy electrons

#### B. Electron Transport Chain

- a series of electron carrier proteins that move the high energy electrons from PS II to PS I
- proteins in ETC use energy from electrons to pump hydrogen ions from stroma into thylakoid space
- plastoquinone(PQ): first electron carrier
- cytochrome complex: second carrier(pump hydrogen ions)
- plastocyanin(PC): third electron carrier

#### C. Photosystem I

- pigment uses solar energy to recognize electrons
- electrons pass down the second short ETC to protein ferredoxin(Fd)
- NADP<sup>+</sup> picks up 2 high energy electrons--NADPH
- NADP<sup>+</sup> reductase catalyzes the transfer of electrons from Fd to NADP<sup>+</sup>

#### D. Hydrogen ion movement

- in thylakoid space: more hydrogen ions(positively charged)
- in stroma: fewer hydrogen ions(negatively charged)
- gradient of hydrogen ions provide energy to make ATP

#### E. ATP Formation

- hydrogen ions pass through ATP synthase
- ATP synthase rotates and produces ATP
- chemiosmosis: process that uses energy stored in the proton gradient to make ATP

#### Calvin Cycle

- uses ATP and NADPH from light dependent reaction to produce high energy sugars in stroma
- ribulose biphosphate(RuBP): 5 carbon compound, combine with carbon dioxide

- glyceraldehyde-3 phosphate(G3P): 3 carbon compound

A. CO<sup>2</sup> enter the cycle

- $6\text{CO}_2 + 6\text{RuBP} \rightarrow 12\text{PGA}$
- catalyzed by an enzyme Rubisco

B. Energy Input

- 12 ATP and 12 NADPH provide energy to convert the 12 PGA into 12G3P

C. 6 carbon sugar produced

- two of 12 G3P convert into one 6-carbon sugar
- remaining 10 G3P converted to 6 RuBP

D. RuBP regenerated

- 10 G3P converted to 6 RuBP
- uses energy from 6 ATP
- 6 RuBP combine with 6 new carbon dioxide, new cycles begin

- Overall, 6 molecules of carbon dioxide is used to produce one 6-carbon sugar which stores long term energy, builds complex carbohydrate
- 12 NADPH and 18 ATP consumed

Measuring rate of photosynthesis

- oxygen gas production
- carbon dioxide consumption
- biomass increase of plant

Factors of Photosynthesis

- Water
  - raw material
  - plants have cuticle to prevent water loss
    - waxy waterproof layer on leaves
- Temperature
  - most enzymes work best between 0-35 degree celsius
  - too high temperature will denature the photosynthetic enzyme

- Light intensity
  - light intensity increase, rate increase
  - after maximum rate, reach a plateau
- Carbon dioxide
  - low to fairly high carbon dioxide concentrations: rate positively correlated to carbon dioxide concentration
  - very high carbon dioxide concentrations: reach plateau
  - very low carbon dioxide concentrations: no photosynthesis

#### C4 plants

- spatially separate capture of carbon dioxide and use of carbon dioxide
- PEP carboxylase has high affinity for carbon dioxide but low affinity for oxygen gas
- light dependent reaction happens in mesophyll cell
- bundle-sheath cell is a high carbon dioxide, low oxygen gas environment
- fix carbon dioxide into 4-carbon compounds instead of 3-carbon molecules
- capture at even very low levels of carbon dioxide
- enables photosynthesis to keep working under intense light and high temperatures
- E.g. Corn, sugarcane, sorghum

#### CAM plants

- adapted to dry climate
- keep stomata closed during the day, allow carbon dioxide to enter leaves only at night
- E.g. cacti, pineapple tree