

CLASSICAL FOURIER ANALYSIS: INTERPOLATION OF L^p SPACES

YANNIS BÄHNI

DEFINITION 1.1. *Let (X, μ) and (Y, ν) be measure spaces. Further let T be an operator defined on a linear space of complex-valued measurable functions on X and taking values in the set of all complex-valued, finite almost everywhere, measurable functions on Y . Then T is called linear if for all functions f and g in the domain of T and all $z \in \mathbb{C}$ holds*

$$T(f + g) = T(f) + T(g) \quad T(zf) = zT(f) \quad (1)$$

and quasi-linear if

$$|T(f + g)| \leq K(|T(f)| + |T(g)|) \quad |T(zf)| = |z| |T(f)| \quad (2)$$

holds for some real constant $K > 0$. If $K = 1$, T is called sublinear.

(Yannis Bähni) UNIVERSITY OF ZÜRICH, RÄMISTRASSE 71, 8006 ZÜRICH
E-mail address: yannis.baehni@uzh.ch.

Recall, that a complex-valued function f is said to be *holomorphic* in $\Omega \subseteq \mathbb{C}$ open, if $f'(z)$ exists for any $z \in \Omega$.

LEMMA 1.1. (Hadamard's three lines lemma) *Let F be a holomorphic function in the strip $S := \{z \in \mathbb{C} : 0 < \operatorname{Re} z < 1\}$, continuous and bounded on \overline{S} , such that $|F(z)| \leq B_0$ when $\operatorname{Re} z = 0$ and $|F(z)| \leq B_1$ when $\operatorname{Re} z = 1$, for some $0 < B_0, B_1 < \infty$. Then $|F(z)| \leq B_0^{1-\theta} B_1^\theta$ when $\operatorname{Re} z = \theta$, for any $0 \leq \theta \leq 1$.*

Proof. For $z \in \overline{S}$ define

$$G(z) := \frac{F(z)}{B_0^{1-z} B_1^z} \quad G_n(z) := G(z) e^{(z^2-1)/n}, \quad n \in \mathbb{N}_{>0}$$

$G(z)$ and $G_n(z)$ are holomorphic in S by

$$G'(z) = \frac{F'(z) - F(z) \log(B_1/B_0)}{B_0^{1-z} B_1^z} \quad G'_n(z) = G'(z) e^{(z^2-1)/n} + \frac{2}{n} z G_n(z)$$

and $e^z \neq 0$ for every $z \in \mathbb{C}$. Further, we have

$$|B_0^{1-z} B_1^z| = (B_0^{1-z} B_0^{1-\bar{z}} B_1^z B_1^{\bar{z}})^{1/2} = B_0^{1-\operatorname{Re} z} B_1^{\operatorname{Re} z}$$

Consider $0 \leq \operatorname{Re} z \leq 1$ and $B_0 \geq 1$. Then $B_0^{1-\operatorname{Re} z} \geq 1$ and $B_0^{1-\operatorname{Re} z} \geq B_0$ in the case $B_0 < 1$. Similarly, $B_1^{\operatorname{Re} z} \geq 1$ if $B_1 \geq 1$ and $B_1^{\operatorname{Re} z} \leq B_1$ if $B_1 < 1$. Hence

$$|B_0^{1-z} B_1^z| \geq \min\{1, B_0\} \min\{1, B_1\} > 0 \quad (3)$$

for all $z \in \overline{S}$. Since F is bounded on \overline{S} , we have $|F(z)| \leq L$ for some $L > 0$ and all $z \in \overline{S}$. Thus by (3)

$$|G(z)| = \frac{|F(z)|}{|B_0^{1-z} B_1^z|} \leq \frac{L}{\min\{1, B_0\} \min\{1, B_1\}} =: M$$

for every $z \in \overline{S}$. Fix $n \in \mathbb{N}_{>0}$ and write $z := x + iy \in \overline{S}$. Then

$$|G_n(z)| \leq M \left(e^{(x^2+2ixy-y^2-1)/n} e^{(x^2-2ixy-y^2-1)/n} \right)^{1/2} = M e^{-y^2/n} e^{(x^2-1)/n} \leq M e^{-y^2/n}$$

for $0 \leq x \leq 1$. Thus

$$\lim_{y \rightarrow \pm\infty} \sup\{|G_n(z)| : 0 \leq x \leq 1\} = 0$$

by the pinching-principle. Hence there exist $C_0, C_1 \in \mathbb{R}$, such that

$$\sup\{|G_n(z)| : 0 \leq x \leq 1\} \leq 1$$

when $y > C_0$ or $y < C_1$ and so there exists some $C(n) > 0$ such that $|G_n(z)| \leq 1$ for all $0 \leq x \leq 1$ when $|y| \geq C(n)$. Now consider the rectangle $R := (0, 1) \times (-C(n), C(n))$. We have $|G_n(z)| \leq 1$ on the lines $[0, 1] \times \{\pm C(n)\}$. By

$$|G_n(iy)| = |F(iy)| / B_0 \leq 1 \quad |G_n(1 + iy)| = |F(1 + iy)| / B_1 \leq 1$$

we have $|G_n(z)| \leq 1$ on the line $\{0\} \times [-C(n), C(n)]$ and $\{1\} \times [-C(n), C(n)]$. Thus $|G_n(z)| \leq 1$ on ∂R . Since $|G_n(z)|$ is continuous on \overline{R} , R is a bounded region and holomorphic in R , the maximum modulus theorem implies

$$|G_n(z)| \leq \sup \{|G_n(z)| : z \in \partial R\} \leq 1$$

for every $z \in R$. Therefore $|G_n(z)| \leq 1$ on \overline{R} and so $|G_n(z)| \leq 1$ on \overline{S} . Since inequalities are preserved by limits and the modulus is a continuous function, we have that $|G(z)| = \lim_{n \rightarrow \infty} |G_n(z)| \leq 1$ for $z \in \overline{S}$. We conclude by

$$|F(\theta + it)| = |G(\theta + it)| \left| B_0^{1-\theta-it} B_1^{\theta+it} \right| \leq B_0^{1-\theta} B_1^\theta$$

whenever $0 \leq \theta \leq 1, t \in \mathbb{R}$. □

THEOREM 1.1. (Riesz-Thorin Interpolation Theorem) *Let (X, μ) be a measure space, (Y, ν) a semifinite measure space and T be a linear operator defined on Σ_X and taking values in the set of measurable functions on Y . Let $1 \leq p_0, p_1, q_0, q_1 \leq \infty$ and assume that*

$$\|T(f)\|_{L^{q_0}} \leq M_0 \|f\|_{L^{p_0}} \quad \|T(f)\|_{L^{q_1}} \leq M_1 \|f\|_{L^{p_1}} \quad (4)$$

for all $f \in \Sigma_X$ and $M_0, M_1 < \infty$. Then for all $0 < \theta < 1$ we have

$$\|T(f)\|_{L^q} \leq M_0^{1-\theta} M_1^\theta \|f\|_{L^p} \quad (5)$$

for all $f \in \Sigma_X$, where

$$\frac{1}{p} = \frac{1-\theta}{p_0} + \frac{\theta}{p_1} \quad \frac{1}{q} = \frac{1-\theta}{q_0} + \frac{\theta}{q_1} \quad (6)$$

LEMMA 1.2. (Hadamard's three lines lemma, extension) *Let F be a holomorphic function in the strip $S := \{z \in \mathbb{C} : 0 < \operatorname{Re} z < 1\}$ and continuous on \bar{S} , such that for some $0 < A < \infty$ and $\tau_0 \in (0, \pi)$ we have $\log |F(z)| \leq A e^{\tau_0 |\operatorname{Im} z|}$ for every $z \in \bar{S}$. Then*

$$|F(z)| \leq \exp \left(\frac{\sin(\pi x)}{2} \int_{-\infty}^{\infty} \left[\frac{\log |F(it + iy)|}{\cosh(\pi t) - \cos(\pi x)} + \frac{\log |F(1 + it + iy)|}{\cosh(\pi t) + \cos(\pi x)} \right] d\lambda(t) \right)$$

whenever $z := x + iy \in S$.

DEFINITION 1.2. (Analytic family, admissible growth) *Let (X, μ) be a measure space, (Y, ν) be a semifinite measure spaces and $(T_z)_{z \in \bar{S}}$, where T_z is defined on Σ_X and taking values in the space of all measurable functions on Y such that*

$$\int_Y |T_z(\chi_A) \chi_B| d\nu \quad (7)$$

whenever $\mu(A), \nu(B) < \infty$. The family $(T_z)_{z \in \bar{S}}$ is said to be analytic if for all $f \in \Sigma_X, g \in \Sigma_Y$ we have that

$$z \mapsto \int_Y T_z(f) g d\nu \quad (8)$$

is analytic on S and continuous on \bar{S} . Further, an analytic family $(T_z)_{z \in \bar{S}}$ is called of admissible growth, if there is a constant $\tau_0 \in (0, \pi)$, such that for all $f \in \Sigma_X, g \in \Sigma_Y$ a constant $C(f, g)$ exists with

$$\log \left| \int_Y T_z(f) g d\nu \right| \leq C(f, g) e^{\tau_0 |\operatorname{Im} z|} \quad (9)$$

for all $z \in \overline{S}$.

THEOREM 1.2. (Stein's Theorem on Interpolation of Analytic Families of Operators) *Let $(T_z)_{z \in \overline{S}}$ be an analytic family of admissible growth, $1 \leq p_0, p_1, q_0, q_1 \leq \infty$ and suppose that M_0, M_1 are positive functions on the real line such that for some $\tau_1 \in (0, \pi)$*

$$\sup \left\{ e^{-\tau_1 |y|} \log M_0(y) : y \in \mathbb{R} \right\} < \infty \quad \sup \left\{ e^{-\tau_1 |y|} \log M_1(y) : y \in \mathbb{R} \right\} < \infty \quad (10)$$

Fix $0 < \theta < 1$ and define

$$\frac{1}{p} := \frac{1-\theta}{p_0} + \frac{\theta}{p_1} \quad \frac{1}{q} := \frac{1-\theta}{q_0} + \frac{\theta}{q_1} \quad (11)$$

Further suppose that for all $f \in \Sigma_X$ and $y \in \mathbb{R}$ we have

$$\|T_{iy}(f)\|_{L^{q_0}} \leq M_0(y) \|f\|_{L^{p_0}} \quad \|T_{1+iy}(f)\|_{L^{q_1}} \leq M_1(y) \|f\|_{L^{p_1}} \quad (12)$$

Then for all $f \in \Sigma_X$ we have

$$\|T_\theta(f)\|_{L^q} \leq M(\theta) \|f\|_{L^p}$$

where for $0 < x < 1$

$$M(x) = \exp \left(\frac{\sin(\pi x)}{2} \int_{-\infty}^{\infty} \left[\frac{\log M_0(t)}{\cosh(\pi t) - \cos(\pi x)} + \frac{\log M_1(t)}{\cosh(\pi t) + \cos(\pi x)} \right] d\lambda(t) \right)$$

THEOREM 1.3. (The Marcinkiewicz Interpolation Theorem) *Let (X, μ) be a σ -finite measure space, (Y, ν) another measure space and $0 < p_0 < p_1 \leq \infty$. Further let T be a sublinear operator defined on*

$$L^{p_0} + L^{p_1} := \{f_0 + f_1 : f_0 \in L^{p_0}(X, \mu), f_1 \in L^{p_1}(X, \mu)\}$$

and taking values in the space of measurable functions on Y . Assume that there exist $A_0, A_1 < \infty$ such that

$$\forall f \in L^{p_0}(X, \mu) \quad \|T(f)\|_{L^{p_0, \infty}} \leq A_0 \|f\|_{L^{p_0}} \quad (13)$$

$$\forall f \in L^{p_1}(X, \mu) \quad \|T(f)\|_{L^{p_1, \infty}} \leq A_1 \|f\|_{L^{p_1}} \quad (14)$$

Then for all $p_0 < p < p_1$ and for all $f \in L^p(X, \mu)$ we have the estimate

$$\|T(f)\|_{L^p} \leq A \|f\|_{L^p} \quad (15)$$

where

$$A := 2 \left(\frac{p}{p-p_0} + \frac{p}{p_1-p} \right)^{1/p} A_0^{\frac{\frac{1}{p}-\frac{1}{p_1}}{\frac{1}{p_0}-\frac{1}{p_1}}} A_1^{\frac{\frac{1}{p_0}-\frac{1}{p}}{\frac{1}{p_0}-\frac{1}{p_1}}} \quad (16)$$