

TORIC SYMPLECTIC MANIFOLDS

SIMON GRÜNING

(DARBOUX THEOREM)

(!) We follow closely the exposition of Yannis gathered of material from Lee and Salomon's books. We take inspiration from Anna's stuff.

1. PREREQUISITES

REMARK 1.1. *Must be even dimensional*

All manifolds are smooth unless noted otherwise as per ana's conventions

chart centered on x means $\phi(x) = 0$.

F diffeomorphism is linear?

*precomposition definition of F^**

PROPOSITION 1.1 (Cartan's Magic Formula). (*! ref lee*)

Fix a manifold M , a vector field $X \in \mathfrak{X}(M)$, and an $\omega \in \Omega^k(M)$ for some $k \in \mathbb{N}$. Then

$$\mathcal{L}_X \omega = i_X(d\omega) + d(i_X \omega).$$

THEOREM 1.1. *Canonical Form theorem*

DEFINITION 1.1. *time dependent vector field*

DEFINITION 1.2. *time dependent flow of X*

DEFINITION 1.3. *time dependent differential k -form*

PROPOSITION 1.2 (Fisherman's Formula). (*!ref lee*) *Fix a manifold M . If $X : J \times M \rightarrow TM$ is a time-dependent vector field with time-dependent flow $\Psi : \mathcal{D} \rightarrow M$ then for any $\omega \in \Omega^k(M)$, $(t_1, t_0, x) \in \mathcal{D}$ we have that*

$$\left. \frac{d}{dt} \right|_{t=t_1} \Psi_{t,t_0}^* \omega = \Psi_{t_1,t_0}^* \left(\mathcal{L}_{X_{t_1}} \omega \right)$$

with $X_{t_1} := X(t_1, \cdot) \in \mathfrak{X}(M)$.

PROPOSITION 1.3 (Fisherman's Formula Adapted). *Fix a manifold M . If $X : J \times M \rightarrow TM$ is a time-dependent vector field with time-dependent flow $\Psi : \mathcal{D} \rightarrow M$, and further, $\omega : J \times M \rightarrow \Lambda^k T^*M$ is a time-dependent differential k -form, then for any $\omega \in \Omega^k(M)$, $(t_1, t_0, x) \in \mathcal{D}$ we have that*

$$\left. \frac{d}{dt} \right|_{t=t_1} \Psi_{t,t_0}^* \omega_t = \Psi_{t_1,t_0}^* \left(\mathcal{L}_{X_{t_1}} \omega_{t_1} + \left. \frac{d}{dt} \right|_{t=t_1} \omega_t \right)$$

with $X_{t_1} := X(t_1, \cdot) \in \mathfrak{X}(M)$.

Proof. todo chain rule amazing □

LEMMA 1.1. *Let M be a manifold, $x \in M$ with basis (e_i) for $T_x M$. Then there exists a chart (U, x^1, \dots, x^n) centered on x such that for any $i = 1, \dots, n$:*

$$\left. \frac{\partial}{\partial x^i} \right|_x = e_i$$

DEFINITION 1.4 (Tubular Neighbourhood). (*! ref lee*)

PROPOSITION 1.4 (Homotopy Formula). (*!ref canas*). Fix U , a tubular neighborhood of a submanifold S embedded in M . If $\omega \in \Omega^k(U)$ is closed and $i^*\omega = 0$ for some $i : S \hookrightarrow U$, then there exists an $\eta \in \Omega^{k-1}(U)$ with $\omega = d\eta$ and $\forall x \in S : \eta_x = 0$.

Proof. todo □

2. MOSER TRICK

THEOREM 2.1. *Moser Trick*

THEOREM 2.2 (Moser Isotopy). (*!salomon*) Fix as M a $2n$ -dimensional manifold and as $S \subseteq M$ a compact submanifold. If $\omega_0, \omega_1 \in \Omega^2(M)$ are close and

- (1) $\forall x \in S : \omega_0|_x = \omega_1|_x$
- (2) $\forall x \in S : \omega_0|_x, \omega_1|_x$ are nondegenerate.

then there exist neighbourhoods U_0, U_1 of S in M and a diffeomorphism $F : U_0 \rightarrow U_1$ with

$$\begin{aligned} F|_S &= id_S \\ F^*(\omega_1|_{U_1}) &= \omega_0|_{U_0}. \end{aligned}$$

Proof. asf □

3. DARBOUX THEOREM

THEOREM 3.1 (Darboux's Theorem). Fix (M, ω) a $2n$ -dimensional symplectic manifold, $x \in M$. Then there exists a chart $(U, x^1, \dots, x^n, y^1, \dots, y^n)$ centered on x such that:

$$\omega|_U = \sum_{i=1}^n dx^i \wedge dy^i$$

.

Proof. The canonical form theorem for symplectic tensors 1.1 provides us a basis $(a_1, \dots, a_n, b_1, \dots, b_n)$ for $T_x M$ such that for its dual basis $(a^1, \dots, a^n, b^1, \dots, b^n)$ we have

$$\omega_x = \sum_{i=1}^n da^i \wedge db^i.$$

By proposition 1.1 we further have a chart $(U, \tilde{\varphi})$ centered on x with associated coordinates $(\tilde{x}^1, \dots, \tilde{x}^n, \tilde{y}^1, \dots, \tilde{y}^n)$ such that for $i = 1, \dots, n$

$$\begin{aligned} \left. \frac{\partial}{\partial \tilde{x}^i} \right|_x &= a_i \\ \left. \frac{\partial}{\partial \tilde{y}^i} \right|_x &= b_i \end{aligned}$$

Combining the previous two results and traversing again into the dual basis we have

$$\omega_x = \sum_{i=1}^n d\tilde{x}^i|_x \wedge d\tilde{y}^i|_x.$$

Define:

$$\begin{aligned}\omega_0 &:= \omega|_U \\ \omega_1 &:= \sum_{i=1}^n d\tilde{x}^i \wedge d\tilde{y}^i.\end{aligned}$$

Then ω_0, ω_1 are symplectic forms on U (!).

Application of the Moser isotopy 2.2 to the compact submanifold $x \subseteq U$ given ω_0, ω_1 provides the existence of neighbourhoods U_0, U_1 of x in U and a diffeomorphism $F : U_0 \rightarrow U_1$ with

$$\begin{aligned}F(x) &= x \\ F^*\omega_1 &= \omega_0.\end{aligned}$$

Define now another chart (U_0, φ) with $\varphi := \tilde{\varphi}|_{U_1} \circ F$. By construction (!) the associated coordinates are

$$\begin{aligned}x^i &= \tilde{x}^i \circ F \\ y^i &= \tilde{y}^i \circ F.\end{aligned}$$

It then follows (!) that $\varphi(x) = \tilde{\varphi}(x) = 0$. The remaining property of our chart (U_0, φ) follows by:

$$\begin{aligned}\omega|_{U_0} &= \omega_0|_{U_0} \\ &= F^*(\omega_1|_{U_1}) \\ &= F^*\left(\sum_{i=1}^n d\tilde{x}^i \wedge d\tilde{y}^i\right) \\ &= \sum_{i=1}^n F^*(d\tilde{x}^i \wedge d\tilde{y}^i) \\ &= \sum_{i=1}^n F^*(d\tilde{x}^i) \wedge F^*(d\tilde{y}^i) \\ &= \sum_{i=1}^n dF^*(\tilde{x}^i) \wedge dF^*(\tilde{y}^i) \\ &= \sum_{i=1}^n d(\tilde{x}^i \circ F) \wedge d(\tilde{y}^i \circ F) \\ &= \sum_{i=1}^n dx^i \wedge dy^i.\end{aligned}$$

□

4. DISCUSSION / APPLICATIONS

bla

APPENDIX A. BASICS

- (1) symplectic manifold
- (2) smooth manifold
- (3) einstein summation convention
- (4) $T_x M$ et al.
- (5) basis of above and dual basis
- (6) coordinates associated to chart?
- (7) time dependent vector field and flow
- (8) time dep differential k-form

LEMMA A.1 (refyan E.). *For a smooth function F from manifolds M to N and $\omega, \eta \in \Omega(N)$ we have*

$$F^*(\omega \wedge \eta) = F^*\omega \wedge F^*\eta$$

LEMMA A.2 (refyan E.203). *For a smooth function F from manifolds M to N and $\omega \in \Omega(M)$ we have*

$$F^*(d\omega) = d(F^*\omega).$$