

eMeasure Title	Ischemic Vascular Disease (IVD): Complete Lipid Profile and LDL-C Control (<100 mg/dL)		
eMeasure Identifier (Measure Authoring Tool)	182	eMeasure Version number	6.0.000
NQF Number	Not Applicable	GUID	500e4792-7f94-4e34-8546-ee71c56fe463
Measurement Period	January 1, 20XX through December 31, 20XX		
Measure Steward	National Committee for Quality Assurance		
Measure Developer	National Committee for Quality Assurance		
Endorsed By	None		
Description	Percentage of patients 18 years of age and older who were diagnosed with acute myocardial infarction (AMI), coronary artery bypass graft (CABG) or percutaneous coronary interventions (PCI) in the 12 months prior to the measurement period, or who had a diagnosis of ischemic vascular disease (IVD) during the measurement period, and who had a complete lipid profile performed during the measurement period and whose most recent Low-density Lipoprotein (LDL-C) was adequately controlled (< 100 mg/dL)		
Copyright	Physician Performance Measure (Measures) and related data specifications were developed by the National Committee for Quality Assurance (NCQA). The Measures are copyrighted but can be reproduced and distributed, without modification, for noncommercial purposes (eg, use by healthcare providers in connection with their practices). Commercial use is defined as the sale, licensing, or distribution of the Measures for commercial gain, or incorporation of the Measures into a product or service that is sold, licensed or distributed for commercial gain. Commercial use of the Measures requires a license agreement between the user and NCQA. NCQA is not responsible for any use of the Measures. (C) 2008-2016 National Committee for Quality Assurance. All Rights Reserved. Limited proprietary coding is contained in the Measure specifications for user convenience. Users of proprietary code sets should obtain all necessary licenses from the owners of the code sets. NCQA disclaims all liability for use or accuracy of any CPT or other codes contained in the specifications. CPT(R) contained in the Measure specifications is copyright 2004-2015 American Medical Association. LOINC(R) copyright 2004-2015 Regenstrief Institute, Inc. This material contains SNOMED Clinical Terms(R) (SNOMED CT [R]) copyright 2004-2015 International Health Terminology Standards Development Organisation. ICD-10 copyright 2015 World Health Organization. All Rights Reserved.		
Disclaimer	These performance Measures are not clinical guidelines and do not establish a standard of medical care, and have not been tested for all potential applications. THE MEASURES AND SPECIFICATIONS ARE PROVIDED "AS IS" WITHOUT WARRANTY OF ANY KIND. Due to technical limitations, registered trademarks are indicated by (R) or [R] and unregistered trademarks are indicated by (TM) or [TM].		
Measure Scoring	Proportion		
Measure Type	Process		
Stratification	None		
Risk Adjustment	None		
Rate Aggregation	None		
Rationale	A 10 percent decrease in total cholesterol levels (population wide) may result in an estimated 30 percent reduction in the incidence of coronary heart disease (CHD) (Centers for Disease Control and Prevention 2000). Based on data from the Third Report of the Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults: *Less than half of persons who qualify for any kind of lipid-modifying treatment for CHD risk reduction are receiving it *Less than half of even the highest-risk persons, those who have symptomatic CHD, are receiving lipid-lowering treatment *Only about a third of treated patients are achieving their LDL goal; less than 20 percent of CHD patients are at their LDL goal (National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Pressure 2002) According to data from the Behavioral Risk Factor Surveillance System (BRFSS) from 1991 - 2003, the prevalence of cholesterol screening during the preceding 5 years increased from 67.3 percent in 1991 to 73.1 percent in 2003 (Centers for Disease Control and Prevention 2005). Between 1988-94 and 1999-2002, the age-adjusted mean total serum cholesterol level of adults 20 years of age and older decreased from 206 mg/dL to 203 mg/dL, and LDL cholesterol levels decreased from 129 mg/dL to 123 mg/dL. The mean level of LDL cholesterol for American adults age 20 and older is 123 mg/dL (Carroll et al. 2005). However, even given this decrease, there is still a significant amount of room for improvement.		
Clinical Recommendation Statement	Third report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). (2002) AND Implications of recent clinical trials for the National Cholesterol Education Program Adult Treatment Panel III guidelines (2004) In high-risk persons, the recommended LDL-C goal is < 100 mg/dL. * An LDL-C goal of < 70 mg/dL is a therapeutic option on the basis of available clinical trial evidence, especially for patients at very high risk. * If LDL-C is > 100 mg/dL, an LDL-lowering drug is indicated simultaneously with lifestyle changes. * If baseline LDL-C is < 100 mg/dL, institution of an LDL-lowering drug to achieve an LDL-C level < 70 mg/dL is a therapeutic option on the basis of available clinical trial evidence. * If a high-risk person has high triglycerides or low HDL-C, consideration can be given to combining a fibrate or nicotinic acid with an LDL-lowering drug. When triglycerides are > 200 mg/dL, non-HDL-C is a secondary target of therapy, with a goal 30 mg/dL higher than the identified LDL-C goal. The U.S. Preventive Services Task Force (USPSTF) strongly recommends screening men aged 35 and older for lipid disorders and recommends screening men aged 20 to 35 for lipid disorders if they are at increased risk for coronary heart disease. The USPSTF also strongly recommends screening women aged 45 and older for lipid disorders if they are at increased risk for coronary heart disease and recommends screening women aged 20 to 45 for lipid disorders if they are at increased risk for coronary heart disease.		
Improvement Notation	Higher score indicates better quality		
Reference	Centers for Disease Control and Prevention (CDC). 2005. "Trends in cholesterol screening and awareness of high		

	blood cholesterol - United States, 1991-2003." MMWR 54:865-870.
Reference	Carroll, M.D., D.A. Lacher, P.D. Sorlie, J.I. Cleeman, D.J. Gordon, M. Wolz, S.M. Grundy, C.L. Johnson. 2005. "Trends in serum lipids and lipoproteins of adults. 1960-2002." JAMA 294:1773-1781.
Reference	Centers for Disease Control and Prevention (CDC). 2000. "State-specific cholesterol screening trends - United States, 1991-1999." MMWR 49:750-755.
Reference	National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). 2002. "Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III) final report." Circulation 106(25):3143-421.
Reference	Grundy SM, Cleeman JI, Merz CN, et al. Implications of recent clinical trials for the National Cholesterol Education Program Adult Treatment Panel III guidelines. Circulation 2004; 110: 227-39.
Reference	U.S. Preventive Services Task Force. 2008. "Screening for lipid disorders in adults" (June) http://www.uspreventiveservicestaskforce.org/uspstf/uspchol.htm
Definition	None
Guidance	Only patients who were diagnosed with acute myocardial infarction (AMI), coronary artery bypass graft (CABG) or percutaneous coronary interventions (PCI) are included in this measure
Transmission Format	None
Initial Population	Patients 18 years of age and older with a visit during the measurement year who had an AMI, CABG, or PCI during the 12 months prior to the measurement year or who had a diagnosis of IVD during the measurement year
Denominator	Equals Initial Population
Denominator Exclusions	None
Numerator	Numerator 1: Patients with a complete lipid profile performed during the measurement period Numerator 2: Patients whose most recent LDL-C level performed during the measurement period is <100 mg/dL
Numerator Exclusions	Not Applicable
Denominator Exceptions	None
Supplemental Data Elements	For every patient evaluated by this measure also identify payer, race, ethnicity and sex

Table of Contents

- [Population Criteria](#)
- [Data Criteria \(QDM Variables\)](#)
- [Data Criteria \(QDM Data Elements\)](#)
- [Supplemental Data Elements](#)
- [Risk Adjustment Variables](#)

Population Criteria

----- Population Criteria 1 -----

- **Initial Population =**
 - AND: Age >= 18 year(s) at: "Measurement Period"
 - AND:
 - OR: "Diagnosis: Acute Myocardial Infarction" < 12 month(s) starts before start of "Measurement Period"
 - OR: "Diagnosis: Ischemic Vascular Disease" overlaps "Measurement Period"
 - OR: Union of:
 - "Procedure, Performed: Percutaneous Coronary Interventions"
 - "Procedure, Performed: Coronary Artery Bypass Graft"
 - < 12 month(s) ends before start of "Measurement Period"
 - AND: Union of:
 - "Encounter, Performed: Office Visit"
 - "Encounter, Performed: Face-to-Face Interaction"
 - "Encounter, Performed: Preventive Care Services - Established Office Visit, 18 and Up"
 - "Encounter, Performed: Preventive Care Services-Initial Office Visit, 18 and Up"
 - "Encounter, Performed: Home Healthcare Services"
 - "Encounter, Performed: Annual Wellness Visit"
 - during "Measurement Period"
- **Denominator =**
 - AND: Initial Population
- **Denominator Exclusions =**
 - None
- **Numerator =**
 - AND: "Laboratory Test, Performed: Total Cholesterol Laboratory Test (result)" during "Measurement Period"
 - AND: "Laboratory Test, Performed: HDL-C Laboratory Test (result)" during "Measurement Period"
 - AND: "Laboratory Test, Performed: LDL-c (result)" during "Measurement Period"
 - AND: "Laboratory Test, Performed: Triglycerides Laboratory Test (result)" during "Measurement Period"
- **Numerator Exclusions =**
 - None
- **Denominator Exceptions =**
 - None
- **Stratification =**
 - None

----- Population Criteria 2 -----

- **Initial Population =**
 - AND: Age >= 18 year(s) at: "Measurement Period"
 - AND:
 - OR: "Diagnosis: Acute Myocardial Infarction" < 12 month(s) starts before start of "Measurement Period"
 - OR: "Diagnosis: Ischemic Vascular Disease" overlaps "Measurement Period"
 - OR: Union of:
 - "Procedure, Performed: Percutaneous Coronary Interventions"
 - "Procedure, Performed: Coronary Artery Bypass Graft"
 - < 12 month(s) ends before start of "Measurement Period"
 - AND: Union of:
 - "Encounter, Performed: Office Visit"
 - "Encounter, Performed: Face-to-Face Interaction"

- "Encounter, Performed: Preventive Care Services - Established Office Visit, 18 and Up"
 - "Encounter, Performed: Preventive Care Services-Initial Office Visit, 18 and Up"
 - "Encounter, Performed: Home Healthcare Services"
 - "Encounter, Performed: Annual Wellness Visit"
 - during "Measurement Period"
- **Denominator =**
 - AND: Initial Population
- **Denominator Exclusions =**
 - None
- **Numerator =**
 - AND: "Laboratory Test, Performed: LDL-c" satisfies all:
 - Most Recent: (result) during "Measurement Period"
 - (result < 100 mg/dL)
- **Numerator Exclusions =**
 - None
- **Denominator Exceptions =**
 - None
- **Stratification =**
 - None

Data Criteria (QDM Variables)

- None

Data Criteria (QDM Data Elements)

- "Diagnosis: Acute Myocardial Infarction" using "Acute Myocardial Infarction Grouping Value Set (2.16.840.1.113883.3.464.1003.104.12.1001)"
- "Diagnosis: Ischemic Vascular Disease" using "Ischemic Vascular Disease Grouping Value Set (2.16.840.1.113883.3.464.1003.104.12.1003)"
- "Encounter, Performed: Annual Wellness Visit" using "Annual Wellness Visit Grouping Value Set (2.16.840.1.113883.3.526.3.1240)"
- "Encounter, Performed: Face-to-Face Interaction" using "Face-to-Face Interaction Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1048)"
- "Encounter, Performed: Home Healthcare Services" using "Home Healthcare Services Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1016)"
- "Encounter, Performed: Office Visit" using "Office Visit Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1001)"
- "Encounter, Performed: Preventive Care Services - Established Office Visit, 18 and Up" using "Preventive Care Services - Established Office Visit, 18 and Up Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1025)"
- "Encounter, Performed: Preventive Care Services-Initial Office Visit, 18 and Up" using "Preventive Care Services-Initial Office Visit, 18 and Up Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1023)"
- "Laboratory Test, Performed: HDL-C Laboratory Test" using "HDL-C Laboratory Test Grouping Value Set (2.16.840.1.113883.3.464.1003.104.12.1012)"
- "Laboratory Test, Performed: LDL-c" using "LDL-c LOINC Value Set (2.16.840.1.113883.3.117.1.7.1.215)"
- "Laboratory Test, Performed: Total Cholesterol Laboratory Test" using "Total Cholesterol Laboratory Test Grouping Value Set (2.16.840.1.113883.3.464.1003.104.12.1013)"
- "Laboratory Test, Performed: Triglycerides Laboratory Test" using "Triglycerides Laboratory Test Grouping Value Set (2.16.840.1.113883.3.464.1003.104.12.1014)"
- "Procedure, Performed: Coronary Artery Bypass Graft" using "Coronary Artery Bypass Graft Grouping Value Set (2.16.840.1.113883.3.464.1003.104.12.1002)"
- "Procedure, Performed: Percutaneous Coronary Interventions" using "Percutaneous Coronary Interventions Grouping Value Set (2.16.840.1.113883.3.464.1003.104.12.1010)"

Supplemental Data Elements

- "Patient Characteristic Ethnicity: Ethnicity" using "Ethnicity CDCREC Value Set (2.16.840.1.114222.4.11.837)"
- "Patient Characteristic Payer: Payer" using "Payer SOP Value Set (2.16.840.1.114222.4.11.3591)"
- "Patient Characteristic Race: Race" using "Race CDCREC Value Set (2.16.840.1.114222.4.11.836)"
- "Patient Characteristic Sex: ONC Administrative Sex" using "ONC Administrative Sex AdministrativeGender Value Set (2.16.840.1.113762.1.4.1)"

Risk Adjustment Variables

- None

Measure Set	None
-------------	------