

eClinicalWorks

PQRS 226 (NQF 0028) (CMS 138) - Preventive Care and Screening: Tobacco Use - Screening and Cessation Intervention

Description

Percentage of patients aged 18 years and older who were screened for tobacco use one or more times within 24 months AND who received cessation counseling intervention if identified as a tobacco user.

Reporting Frequency

Once per reporting period.

Denominator

Patients are included in this denominator if they meet ALL of the following criteria:

Denominator Criteria	Area to Document within eClinicalWorks
They are 18 years of age or older	Patient Information
They have had an appropriate office visit	Progress Notes > Billing Note: The following E&M codes are considered appropriate encounters: 90791, 90792, 90832, 90834, 90837, 90845, 92002, 92004, 92012, 92014, 92521, 92522, 92523, 92524, 92540, 92557, 92625, 96150, 96151, 96152, 97003, 97004, 99201, 99202, 99203, 99204, 99205, 99212, 99213, 99214, 99215, 99406, 99407, G0438, G0439

Numerator

Patients in the denominator are included in this numerator if they meet ALL of the following criteria:

Numerator Criteria	Area to Document within eClinicalWorks
They have been screened for tobacco use at least once in the past 24 months	Record this information as structured data from one of the following locations: ■ Progress Notes > Smart Form > Tobacco Status ■ Progress Notes > Social History > Tobacco Use > Tobacco Use/Smoking > Smart Form > Tobacco Status ■ Progress Notes > Social History > Tobacco Use > Tobacco Use other than smoking > Q: Are you an other tobacco user? ■ Progress Notes > Social History > Tobacco Use > Smoking > Additional Findings: Tobacco User ■ Progress Notes > Social History > Tobacco Use > Smoking > Additional Findings: Tobacco Non-User
If they were identified as a tobacco user, enter at least ONE of the following:	
■ They have had a tobacco cessation intervention recorded as structured data	Record this information as structured data from one of the following locations: ■ Progress Notes > Preventive Medicine > Counseling > Tobacco Use ◆ Q1: Type of tobacco use cessation • select any answer ◆ Q2: Patient counseled on the dangers of tobacco use and urged to quit • enter the date they were counseled ■ Progress Notes > Preventive Medicine > Counseling > Communication to patient ◆ Q1: Counseled the patient on tobacco use; cessation provided • enter the date they were counseled ◆ Q2: Counseled the patient on pregnancy smoking effects • enter the date they were counseled ◆ Q3: Counseled the patient on smoking cessation; education provided • enter the date they were counseled ◆ Q4: Referral given to patient to stop smoking clinic • enter the date on which they were referred ◆ Q5: Smoking cessation assistance provided • enter the date on which assistance was provided ◆ Q6: Referral to smoking cessation advisor was given to patient • enter the date on which they were referred

Numerator Criteria	Area to Document within eClinicalWorks
They have had a tobacco cessation medication prescribed	Record the prescription of a cessation medication from one of the following locations: - Progress Notes > Treatment > Add Medication - Progress Notes > Current Medication > Add Medication - Telephone Encounter > Rx tab - Telephone Encounter > Virtual Visit tab > Current Medication > Add Medication - Telephone Encounter > Virtual Visit tab > Treatment > Add Medication
They have had a tobacco cessation intervention recorded as a procedure	A procedure is recorded from the following location: Progress Notes > Billing
	Note: The following CPT II codes satisfy the numerator: 4004F - Screened, Tobacco user and received cessation intervention 1036F - Current tobacco non user The following CPT II codes do NOT satisfy the numerator: 4004F with 8P modifier - Tobacco screening or cessation not performed, no reason specified
	Note: In the event that a patient is screened for tobacco use and identified as a user but did not receive tobacco cessation intervention, report 4004F with 8P.
Note: CMS defines the following terms thusly: Tobacco Use – Includes use of any type of tobacco. Tobacco Cessation Intervention – Includes brief counseling (three minutes or less), and/or pharmacotherapy.	


Numerator Exclusion

Patients are excluded from this numerator if they meet at least ONE of the following criteria:

Exclusion Criteria	Area to Document within eClinicalWorks
Structured data has been recorded indicating that tobacco screening was not performed	To satisfy this numerator exclusion criteria, record this information as structured data from the following location: ■ Progress Notes > Social History > Tobacco Use > Screening Not Performed > Q1: Reason ◆ O1: Medical reason ◆ O2: Limited life expectancy ■ Progress Notes > Preventive Medicine > Counseling > Tobacco Use > Q2: Patient not counseled ◆ O1: Patient not eligible
	Note: Answering O3: <i>No reason specified</i> to the first question or O2: <i>No reason specified</i> to the second question does NOT exclude patients from the numerator.
A procedure has been recorded indicating that tobacco screening was not performed	Record this information from the following location: ■ Progress Notes > Billing
	Note: The following CPT II code excludes patients from the numerator: ■ 4004F with 1P modifier The following CPT II codes do NOT exclude patients from the numerator: ■ 4004F with 8P modifier - Tobacco screening or cessation not performed, no reason specified

Setup

Structured data must be mapped to community items from the Mapper window:



Access this feature from: Community > Mappings > Structured Data