
Residuals vs Fitted
Residuals
Fitted values
lm(volume ~ girth + height)
0.6
0.4
0.2
0.0
−0.2
−0.4
−0.6
1
2
3
4
5
31
2
18


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(volume ~ girth + height)
31
2
18
−2
−1
0
1
2


Scale−Location
31
18
2
1.5
1.0
0.5
0.0
√|Standardized residuals|
1
2
3
4
5
Fitted values
lm(volume ~ girth + height)


Residuals vs Leverage
Standardized residuals
Leverage
lm(volume ~ girth + height)
Cook's distance
0.5
0.5
31
3
18
3
2
1
0
−1
−2
0.00
0.05
0.10
0.15
0.20


Residuals vs Fitted
Residuals
Fitted values
lm(volume ~ girth + height)
0.6
0.4
0.2
0.0
−0.2
−0.4
−0.6
1
2
3
4
5
31
2
18


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(volume ~ girth + height)
31
2
18
−2
−1
0
1
2


Scale−Location
31
18
2
1.5
1.0
0.5
0.0
√|Standardized residuals|
1
2
3
4
5
Fitted values
lm(volume ~ girth + height)


Residuals vs Leverage
Standardized residuals
Leverage
lm(volume ~ girth + height)
Cook's distance
0.5
0.5
31
3
18
3
2
1
0
−1
−2
0.00
0.05
0.10
0.15
0.20


Residuals vs Fitted
Residuals
Fitted values
lm(volume ~ girth + height)
0.6
0.4
0.2
0.0
−0.2
−0.4
−0.6
1
2
3
4
5
31
2
18


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(volume ~ girth + height)
31
2
18
−2
−1
0
1
2


Scale−Location
31
18
2
1.5
1.0
0.5
0.0
√|Standardized residuals|
1
2
3
4
5
Fitted values
lm(volume ~ girth + height)


Residuals vs Leverage
Standardized residuals
Leverage
lm(volume ~ girth + height)
Cook's distance
0.5
0.5
31
3
18
3
2
1
0
−1
−2
0.00
0.05
0.10
0.15
0.20