

---

## Computations supporting Example 3.5 in the paper Matricial Gaussian Quadrature Rules: Nonsingular Case by Aljaž Zalar and Igor Zbovič.

This example shows a sequence  $S$  with  $k > 0$ , where  $k = \text{rank}(HH - K) - \text{rank } HH$ , where  $HH$  is the  $(x - t)$ -localizing matrix. It exhibits two distinct  $((n + 1)p)$ -atomic  $R$ -representing matrix measures for  $S$ . In both cases, the measures include an atom at  $t$  with the largest multiplicity allowed by Theorem 3.1, namely  $m = p - k$ , demonstrating that a representing measure for  $S$  that contains an atom  $t$  of multiplicity  $m$  is not unique if  $m < p$ .

Let  $p = 2$ ,  $n = 1$  and  $t = 0$ . Let  $S = (S_0, S_1, S_2)$ , where  $S_0 := \{\{18, 10\}, \{10, 7\}\}$ ,  $S_1 := \{\{2, 2\}, \{2, 2\}\}$ ,  $S_2 := \{\{50, 26\}, \{26, 14\}\}$ . Let  $M := M(1)$  the corresponding moment matrix.

```
In[ ]:= S0 = {{18, 10}, {10, 7}};
MatrixForm[S0]
S1 = {{2, 2}, {2, 2}};
MatrixForm[S1]
S2 = {{50, 26}, {26, 14}};
MatrixForm[S2]
M = ArrayFlatten[{{S0, S1}, {S1, S2}}];
MatrixForm[M]

Out[ ]//MatrixForm=

$$\begin{pmatrix} 18 & 10 \\ 10 & 7 \end{pmatrix}$$


Out[ ]//MatrixForm=

$$\begin{pmatrix} 2 & 2 \\ 2 & 2 \end{pmatrix}$$


Out[ ]//MatrixForm=

$$\begin{pmatrix} 50 & 26 \\ 26 & 14 \end{pmatrix}$$


Out[ ]//MatrixForm=

$$\begin{pmatrix} 18 & 10 & 2 & 2 \\ 10 & 7 & 2 & 2 \\ 2 & 2 & 50 & 26 \\ 2 & 2 & 26 & 14 \end{pmatrix}$$

```

We check that  $M$  is positive definite by computing its eigenvalues.

```
In[ ]:= N[Eigenvalues[M]]

Out[ ]:=
{63.9786, 23.6284, 1.17717, 0.215786}
```

Let  $HH = S1$  be the  $(x - t)$ -localizing matrix and let  $K = S2$ . We will compute  $k := \text{rank}(HH \ K) - \text{rank}(HH)$ .

```
In[ ]:= HH = S1;
        K = S2;
        HHK = ArrayFlatten[{{HH, K}}];
        MatrixForm[HHK]
        MatrixRank[HHK]
        MatrixRank[HH]
        k = MatrixRank[HHK] - MatrixRank[HH]

Out[ ]//MatrixForm=

$$\begin{pmatrix} 2 & 2 & 50 & 26 \\ 2 & 2 & 26 & 14 \end{pmatrix}$$


Out[ ]=
2

Out[ ]=
1

Out[ ]=
1
```

We write  $K = (K1 \ K2)$ . We check that  $\text{rank}(HH \ K) = \text{rank}(HH \ K1)$ .

```
In[ ]:= K1 = K[[All, 1 ;; k]];
        MatrixForm[K1]
        K2 = K[[All, k + 1 ;;]];
        MatrixForm[K2]
        HHK1 = ArrayFlatten[{{HH, K1}}];
        MatrixForm[HHK1]
        MatrixRank[HHK1]

Out[ ]//MatrixForm=

$$\begin{pmatrix} 50 \\ 26 \end{pmatrix}$$


Out[ ]//MatrixForm=

$$\begin{pmatrix} 26 \\ 14 \end{pmatrix}$$


Out[ ]//MatrixForm=

$$\begin{pmatrix} 2 & 2 & 50 \\ 2 & 2 & 26 \end{pmatrix}$$


Out[ ]=
2
```

We define  $J := \{-1/2, \{1\}, \{1/2\}\}$ , and we check that  $K2 = (HH \ K1)J$ .

```
In[ ]:= J = {{-1/2}, {1}, {1/2}};
MatrixForm[J]
MatrixForm[HHK1.J]
```

```
Out[ ]//MatrixForm=
```

$$\begin{pmatrix} -\frac{1}{2} \\ 1 \\ \frac{1}{2} \end{pmatrix}$$

```
Out[ ]//MatrixForm=
```

$$\begin{pmatrix} 26 \\ 14 \end{pmatrix}$$

We will now construct two different matrices Z1 and Z2, which will represent S3.

```
In[ ]:= Z11 = 2
Z12 = ArrayFlatten[{{Transpose[K1], Z11}}].J; MatrixForm[Z12]
Z13 = ArrayFlatten[{{Transpose[K2], Z12}}].J; MatrixForm[Z13]
Z1 = ArrayFlatten[{{Z11, Z12}, {Transpose[Z12], Z13}}]; MatrixForm[Z1]
Z21 = 98
Z22 = ArrayFlatten[{{Transpose[K1], Z21}}].J; MatrixForm[Z22]
Z23 = ArrayFlatten[{{Transpose[K2], Z22}}].J; MatrixForm[Z23]
Z2 = ArrayFlatten[{{Z21, Z22}, {Transpose[Z22], Z23}}];
MatrixForm[Z2]
```

```
Out[ ]:=
```

2

```
Out[ ]//MatrixForm=
```

$$(2)$$

```
Out[ ]//MatrixForm=
```

$$(2)$$

```
Out[ ]//MatrixForm=
```

$$\begin{pmatrix} 2 & 2 \\ 2 & 2 \end{pmatrix}$$

```
Out[ ]:=
```

98

```
Out[ ]//MatrixForm=
```

$$(50)$$

```
Out[ ]//MatrixForm=
```

$$(26)$$

```
Out[ ]//MatrixForm=
```

$$\begin{pmatrix} 98 & 50 \\ 50 & 26 \end{pmatrix}$$

We will now obtain two pairs of matrix coefficients (H10, H11) and (H20, H11) which will compose the corresponding generating polynomials.

```
In[ ]:= H1 = Inverse[M].ArrayFlatten[{{S2}, {Z1}}]; MatrixForm[H1]
H10 = H1[[1 ;; 2]]; MatrixForm[H10]
H11 = H1[[3 ;; 4]]; MatrixForm[H11]
polH1[x_] := x^2 IdentityMatrix[2] - (H10 + x H11)
polH1[x_] // Simplify // MatrixForm
H2 = Inverse[M].ArrayFlatten[{{S2}, {Z2}}]; MatrixForm[H2]
H20 = H2[[1 ;; 2]]; MatrixForm[H20]
H21 = H2[[3 ;; 4]]; MatrixForm[H21]
polH2[x_] := x^2 IdentityMatrix[2] - (H20 + x H21)
polH2[x_] // Simplify // MatrixForm
```

```
Out[ ]//MatrixForm=

$$\begin{pmatrix} 3 & \frac{3}{2} \\ 0 & 0 \\ 2 & \frac{1}{2} \\ -4 & -1 \end{pmatrix}$$

```

```
Out[ ]//MatrixForm=

$$\begin{pmatrix} 3 & \frac{3}{2} \\ 0 & 0 \end{pmatrix}$$

```

```
Out[ ]//MatrixForm=

$$\begin{pmatrix} 2 & \frac{1}{2} \\ -4 & -1 \end{pmatrix}$$

```

```
Out[ ]//MatrixForm=

$$\begin{pmatrix} -3 - 2x_+ + x_+^2 & \frac{1}{2}(-3 - x_+) \\ 4x_+ & x_+(1 + x_+) \end{pmatrix}$$

```

```
Out[ ]//MatrixForm=

$$\begin{pmatrix} 3 & \frac{3}{2} \\ 0 & 0 \\ 6 & \frac{5}{2} \\ -8 & -3 \end{pmatrix}$$

```

```
Out[ ]//MatrixForm=

$$\begin{pmatrix} 3 & \frac{3}{2} \\ 0 & 0 \end{pmatrix}$$

```

```
Out[ ]//MatrixForm=

$$\begin{pmatrix} 6 & \frac{5}{2} \\ -8 & -3 \end{pmatrix}$$

```

```
Out[ ]//MatrixForm=

$$\begin{pmatrix} -3 - 6x_+ + x_+^2 & \frac{1}{2}(-3 - 5x_+) \\ 8x_+ & x_+(3 + x_+) \end{pmatrix}$$

```

We compute the determinants from which we get the atoms.

```

In[ ]:= Factor[Det[polH1[x]]]
        Solve[Det[polH1[x]] == 0, x]
        Factor[Det[polH2[x]]]
        Solve[Det[polH2[x]] == 0, x]

Out[ ]:=

$$(-1 + x) x (-3 + x^2)$$


Out[ ]:=

$$\{\{x \rightarrow 0\}, \{x \rightarrow 1\}, \{x \rightarrow -\sqrt{3}\}, \{x \rightarrow \sqrt{3}\}\}$$


Out[ ]:=

$$(-3 + x) (-1 + x) x (1 + x)$$


Out[ ]:=

$$\{\{x \rightarrow -1\}, \{x \rightarrow 0\}, \{x \rightarrow 1\}, \{x \rightarrow 3\}\}$$


```

We will now find the corresponding masses. First we compute the Vandermonde matrices and their inverses.

```
In[*]:= V1 =
  Table[root^i, {i, 0, 3}, {root, x /. Solve[Det[polH1[x]] == 0, x]}] /. Indeterminate -> 1;
MatrixForm[V1]
InvV1 = Inverse[V1]; MatrixForm[InvV1]
V2 =
  Table[root^i, {i, 0, 3}, {root, x /. Solve[Det[polH2[x]] == 0, x]}] /. Indeterminate -> 1;
MatrixForm[V2]
InvV2 = Inverse[V2];
MatrixForm[InvV2]
```

Power: Indeterminate expression  $0^0$  encountered. [i](#)

Out[\*]//MatrixForm=

$$\begin{pmatrix} 1 & 1 & 1 & 1 \\ 0 & 1 & -\sqrt{3} & \sqrt{3} \\ 0 & 1 & 3 & 3 \\ 0 & 1 & -3\sqrt{3} & 3\sqrt{3} \end{pmatrix}$$

Out[\*]//MatrixForm=

$$\begin{pmatrix} 1 & -1 & -\frac{1}{3} & \frac{1}{3} \\ 0 & \frac{3}{2} & 0 & -\frac{1}{2} \\ 0 & \frac{3-3\sqrt{3}}{12\sqrt{3}} & \frac{1}{6} & \frac{-3+\sqrt{3}}{12\sqrt{3}} \\ 0 & \frac{-3-3\sqrt{3}}{12\sqrt{3}} & \frac{1}{6} & \frac{3+\sqrt{3}}{12\sqrt{3}} \end{pmatrix}$$

Power: Indeterminate expression  $0^0$  encountered. [i](#)

Out[\*]//MatrixForm=

$$\begin{pmatrix} 1 & 1 & 1 & 1 \\ -1 & 0 & 1 & 3 \\ 1 & 0 & 1 & 9 \\ -1 & 0 & 1 & 27 \end{pmatrix}$$

Out[\*]//MatrixForm=

$$\begin{pmatrix} 0 & -\frac{3}{8} & \frac{1}{2} & -\frac{1}{8} \\ 1 & -\frac{1}{3} & -1 & \frac{1}{3} \\ 0 & \frac{3}{4} & \frac{1}{2} & -\frac{1}{4} \\ 0 & -\frac{1}{24} & 0 & \frac{1}{24} \end{pmatrix}$$

We compute the corresponding masses by  $\text{InvV1} \{S0, S1, S2, Z1\}$  and  $\text{InvV2} \{S0, S1, S2, Z2\}$ .

```
In[*]:= masses1 = Table[InvV1[[r]].{S0, S1, S2, Z1}, {r, 1, Length[InvV1]}];
MatrixForm[masses1]
masses1Simplified = FullSimplify[ArrayFlatten[Transpose[{masses1}]]];
MatrixForm[masses1Simplified]
masses2 = Table[InvV2[[r]].{S0, S1, S2, Z2}, {r, 1, Length[InvV2]}];
MatrixForm[masses2]
masses2Simplified = FullSimplify[ArrayFlatten[Transpose[{masses2}]]];
MatrixForm[masses2Simplified]
```

Out[\*]//MatrixForm=

$$\begin{pmatrix} \begin{pmatrix} 0 \\ 0 \end{pmatrix} & \begin{pmatrix} 0 \\ 1 \end{pmatrix} \\ \begin{pmatrix} 2 \\ 2 \end{pmatrix} & \begin{pmatrix} 2 \\ 2 \end{pmatrix} \\ \begin{pmatrix} \frac{25}{3} + \frac{3-3\sqrt{3}}{6\sqrt{3}} + \frac{-3+\sqrt{3}}{6\sqrt{3}} \\ \frac{13}{3} + \frac{3-3\sqrt{3}}{6\sqrt{3}} + \frac{-3+\sqrt{3}}{6\sqrt{3}} \end{pmatrix} & \begin{pmatrix} \frac{13}{3} + \frac{3-3\sqrt{3}}{6\sqrt{3}} + \frac{-3+\sqrt{3}}{6\sqrt{3}} \\ \frac{7}{3} + \frac{3-3\sqrt{3}}{6\sqrt{3}} + \frac{-3+\sqrt{3}}{6\sqrt{3}} \end{pmatrix} \\ \begin{pmatrix} \frac{25}{3} + \frac{-3-3\sqrt{3}}{6\sqrt{3}} + \frac{3+\sqrt{3}}{6\sqrt{3}} \\ \frac{13}{3} + \frac{-3-3\sqrt{3}}{6\sqrt{3}} + \frac{3+\sqrt{3}}{6\sqrt{3}} \end{pmatrix} & \begin{pmatrix} \frac{13}{3} + \frac{-3-3\sqrt{3}}{6\sqrt{3}} + \frac{3+\sqrt{3}}{6\sqrt{3}} \\ \frac{7}{3} + \frac{-3-3\sqrt{3}}{6\sqrt{3}} + \frac{3+\sqrt{3}}{6\sqrt{3}} \end{pmatrix} \end{pmatrix}$$

Out[\*]//MatrixForm=

$$\begin{pmatrix} 0 & 0 \\ 0 & 1 \\ 2 & 2 \\ 2 & 2 \\ 8 & 4 \\ 4 & 2 \\ 8 & 4 \\ 4 & 2 \end{pmatrix}$$

Out[\*]//MatrixForm=

$$\begin{pmatrix} \begin{pmatrix} 12 \\ 6 \end{pmatrix} & \begin{pmatrix} 6 \\ 3 \end{pmatrix} \\ \begin{pmatrix} 0 \\ 0 \end{pmatrix} & \begin{pmatrix} 0 \\ 1 \end{pmatrix} \\ \begin{pmatrix} 2 \\ 2 \end{pmatrix} & \begin{pmatrix} 2 \\ 2 \end{pmatrix} \\ \begin{pmatrix} 4 \\ 2 \end{pmatrix} & \begin{pmatrix} 2 \\ 1 \end{pmatrix} \end{pmatrix}$$

Out[\*]//MatrixForm=

$$\begin{pmatrix} 12 & 6 \\ 6 & 3 \\ 0 & 0 \\ 0 & 1 \\ 2 & 2 \\ 2 & 2 \\ 4 & 2 \\ 2 & 1 \end{pmatrix}$$

In[\*]:=

In[\*]:=

*In[ ]:=*

*In[ ]:=*